

See discussions, stats, and author profiles for this publication at:  
<https://www.researchgate.net/publication/245757347>

# Computing the Minimum Fill-In is NP-Complete

Article *in* SIAM Journal on Algebraic and Discrete Methods · March 1981

DOI: 10.1137/0602010

---

CITATIONS

501

---

READS

379

1 author:



Mihalis Yannakakis

Columbia University

267 PUBLICATIONS 25,496 CITATIONS

SEE PROFILE

All content following this page was uploaded by Mihalis Yannakakis on 27 August 2014.

The user has requested enhancement of the downloaded file.

## COMPUTING THE MINIMUM FILL-IN IS NP-COMPLETE

MIHALIS YANNAKAKIS†

**Abstract.** We show that the following problem is NP-complete. Given a graph, find the minimum number of edges (fill-in) whose addition makes the graph chordal. This problem arises in the solution of sparse symmetric positive definite systems of linear equations by Gaussian elimination.

**1. Introduction and terminology.** A graph is a pair  $G = (N, E)$ , where  $N$  is a finite set of *nodes* and  $E$ , a set of unordered pairs  $(u, v)$  of distinct nodes, is a set of *edges*. Two nodes  $u$  and  $v$  are *adjacent* if  $(u, v) \in E$ . The *neighborhood*  $\Gamma(v)$  of a node  $v$  is the set of nodes that are adjacent to  $v$ . The *degree*  $d(v)$  of  $v$  is the number of nodes adjacent to  $v$ . A graph is a *clique* if every two nodes are adjacent. A set of nodes is *independent* if no two of them are adjacent.

If  $S \subseteq N$  is a subset of nodes, the *subgraph* of  $G$  induced by  $S$ , denoted as  $\langle S \rangle$ , is the graph  $(S, E_S)$ , where  $E_S = \{(u, v) \in E \mid u, v \in S\}$ . The graph  $G - S$ , formed by deleting a subset  $S \subseteq N$  of nodes from  $G$ , is  $\langle N - S \rangle$ . A graph  $G = (N, E)$  is *bipartite* if  $N$  can be partitioned into two sets  $P, Q$  of independent nodes; we will write the bipartite graph as  $(P, Q, E)$ . The bipartite graph  $(P, Q, E)$  is a *chain graph* if the neighborhoods of the nodes in  $P$  form a chain; i.e., there is a bijection  $\pi: \{1, 2, \dots, |P|\} \leftrightarrow P$  (an *ordering* of  $P$ ) such that  $\Gamma(\pi(1)) \supseteq \Gamma(\pi(2)) \supseteq \dots \supseteq \Gamma(\pi(|P|))$ . It is easy to see [Y] that then the neighborhoods of the nodes in  $Q$  form also a chain, and thus the definition is unambiguous.

A graph is *chordal* (or *triangulated*) if every cycle of length  $\geq 4$  has a *chord*, i.e., an edge connecting two nonconsecutive nodes of the cycle. Chordal graphs are important in connection with the solution of sparse symmetric positive definite systems of linear equations by Gaussian elimination [R]. From the symmetric  $n \times n$  matrix  $M = (m_{ij})$  of coefficients of such a system we can construct a graph  $G = (N, E)$  with  $n$  nodes, where node  $v_i$  corresponds to the  $i$ th row and column of  $M$  and  $(v_i, v_j) \in E$  iff  $m_{ij} \neq 0$ . The *elimination* of node  $v_i$  from  $G$  is performed by (1) adding edges so that  $\Gamma(v_i)$  becomes a clique, and (2) deleting  $v_i$  from the augmented graph. The added edges correspond to the new nonzero elements that are created when we eliminate the  $i$ th variable, assuming no lucky cancellations. (See [R] for a detailed exposition of this graph-theoretic modeling.) If  $\pi$  is an ordering of  $N$ , the *fill-in*  $F(\pi)$  produced by  $\pi$  is the set of new edges that are added when we eliminate  $\pi(1)$  from  $G$ , then eliminate  $\pi(2)$  from the resulting graph,  $\pi(3)$  from the new graph, etc. The ordering  $\pi$  is a *perfect elimination ordering* if  $F(\pi) = \emptyset$ . Chordal graphs come into the picture because of the following two properties [R]. (1) A graph has a perfect elimination ordering if and only if it is chordal. Thus, "chordal" is a *hereditary* property (i.e., deleting nodes from a chordal graph does not violate the property), and every chordal graph has a node  $v$  such that  $\langle \Gamma(v) \rangle$  is a clique;  $v$  is called a *simplicial* node. (2) If  $\pi$  is an elimination ordering of a graph  $G = (N, E)$ , then the augmented graph  $G_\pi = (N, E \cup F(\pi))$  is chordal:  $\pi$  is a perfect elimination ordering of  $G_\pi$ .

In this paper we examine the problem of finding an elimination ordering which produces a minimum fill-in, or equivalently, finding the minimum set of edges whose addition renders the graph chordal. We shall show that this problem is NP-complete.

\* Received by the editors June 26, 1980.

† Bell Laboratories, Murray Hill, New Jersey 07974.

(For an exposition of NP-completeness see [GJ].) The NP-completeness of the minimum fill-in problem was conjectured in [RTL] and [RT], but a proof had not been found, and it is one of the open problems in [GJ]. The version of the problem on directed graphs was shown to be NP-complete in [RT].

**2. The reduction.** We will make use of chain graphs. Two edges  $(u, v)$ ,  $(x, y)$  are said to be *independent* in a graph  $G$  if the nodes  $u, v, x, y$  are distinct and the subgraph of  $G$  induced by them consists of exactly these two edges. The following lemma from [Y] is easy to prove.

**LEMMA 1.** *A bipartite graph is a chain graph if and only if it does not contain a pair of independent edges.*

Let  $G = (P, Q, E)$  be a bipartite graph. From  $G$  we construct another graph  $C(G) = (N, E')$  by making  $P$  and  $Q$  cliques; i.e.,  $E' = E \cup \{(u, v) | u, v \in P\} \times \cup \{(u, v) | u, v \in Q\}$ .

**LEMMA 2.** *Let  $G$  be a bipartite graph.  $C(G)$  is chordal if and only if  $G$  is a chain graph.*

*Proof (only if).* Suppose that  $G$  is not a chain graph. Then it has two independent edges  $(u, v)$  and  $(x, y)$  by Lemma 1. Suppose without loss of generality that  $u, x \in P$  and  $v, y \in Q$ . Then these two edges together with  $(u, x)$  and  $(v, y)$  form a chordless cycle of length 4 in  $C(G)$ .

*(if).* Suppose that  $G$  is a chain graph, and let  $\pi$  be an ordering of  $P$  such that  $\Gamma(\pi(1)) \supseteq \Gamma(\pi(2)) \supseteq \dots \supseteq \Gamma(\pi(p))$ , where  $p = |P|$ . Since the property of being a chain graph is hereditary, it suffices to show that  $C(G)$  has a simplicial node. The neighborhood of  $\pi(p)$  in  $C(G)$  is  $\Gamma'(\pi(p)) = \Gamma(\pi(p)) \cup [P - \pi(p)]$ . In  $C(G)$  the subgraphs  $\langle P - \pi(p) \rangle$  and  $\langle \Gamma(\pi(p)) \rangle$  are cliques, the latter because  $\Gamma(\pi(p)) \subseteq Q$  and  $\langle Q \rangle$  is a clique. Also, since  $\Gamma(\pi(p)) \subseteq \Gamma(v)$  for every  $v \in P$ , all nodes of  $P$  are adjacent to all nodes of  $\Gamma(\pi(p))$ . Therefore  $\langle \Gamma'(\pi(p)) \rangle$  is a clique, and  $\pi(p)$  is a simplicial node of  $C(G)$ .  $\square$

**LEMMA 3.** *It is NP-complete to find the minimum number of edges whose addition to a bipartite graph  $G = (P, Q, E)$  gives a chain graph.*

*Proof.* The reduction is from the Optimal Linear Arrangement Problem. A linear arrangement of a graph  $G = (N, E)$  is an ordering  $\pi$  of  $N$ . For an edge  $e = (u, v)$  of  $G$ , let  $\delta(e, \pi) = |\pi^{-1}(u) - \pi^{-1}(v)|$ . The cost  $c(\pi)$  of the linear arrangement  $\pi$  is  $c(\pi) = \sum_{e \in E} \delta(e, \pi)$ . The optimal linear arrangement problem is to decide, given a graph  $G$  and an integer  $k$ , whether there exists a linear arrangement  $\pi$  of  $G$  with cost  $c(\pi) \leq k$ . This problem was shown to be NP-complete in [GJS].

Let  $(G = (N, E); k)$  be an instance of the optimal linear arrangement problem. We construct a bipartite graph  $G' = (P, Q, E')$  as follows.  $P$  has one node for every node of  $G$  (i.e.,  $P = N$ );  $Q$  has two nodes  $e_1, e_2$  for every edge  $e$  of  $G$ , and a set  $R(v)$  of  $n - d(v)$  nodes for every node  $v$  of  $N$ , where  $n = |N|$  and  $d(v)$  is the degree of  $v$  in  $G$ . If  $e = (u, v)$  is an edge of  $G$ , then the nodes  $e_1, e_2$  that correspond to  $e$  are adjacent to  $u$  and  $v$ . The nodes in  $R(v)$  are adjacent to  $v$ . In Fig. 1 we show an example of this construction.




FIG. 1

Let  $l(G)$  be the minimum cost of a linear arrangement of  $G$ , and  $h(G')$  the minimum number of edges whose addition to  $G'$  gives a chain graph. We claim that

$$(1) \quad h(G') = l(G) + \frac{n^2(n-1)}{2} - 2m,$$

where  $n, m$  are respectively the numbers of nodes and edges of  $G$ . Thus,  $l(G) \leq k$  iff  $h(G') \leq k + (n^2(n-1)/2) - 2m$ .

First observe that an ordering  $\pi$  of  $N$  specifies uniquely a minimal set  $H(\pi)$  of edges whose addition makes  $G'$  a chain graph with the neighborhoods of the nodes in  $P(=N)$  ordered according to  $\pi$ . For every node  $x$  in  $Q$ , let  $\sigma(x) = \max \{i | (x, \pi(i)) \in E'\}$ . Then  $H(\pi) = \{(x, \pi(j)) | x \in Q, j < \sigma(x)\} - E'$ . Conversely, suppose that  $F$  is a set of edges such that  $G'(F) = (P, Q, E' \cup F)$  is a chain graph and let  $\pi$  be an ordering of the nodes in  $P$  according to their neighborhoods in  $G'(F)$ . It is easy to see that  $F \supseteq H(\pi)$ , and therefore if  $F$  is a minimal augmentation then  $F = H(\pi)$ . Let  $h(\pi) = |H(\pi)|$ . In order to show (1), it suffices thus to show that for every ordering  $\pi$  of  $N$ ,  $h(\pi) = c(\pi) + (n^2(n-1)/2) - 2m$ , where  $c(\pi)$  is the cost of the linear arrangement  $\pi$  of  $G$ .

Let  $\pi$  be an ordering of  $N$ . For every  $v \in N$  and  $x \in R(v)$ ,  $H(\pi)$  contains  $\pi^{-1}(v) - 1$  edges incident to  $x$ . Let  $e = (u, v)$  be an edge of  $G$ , and suppose without loss of generality that  $\pi^{-1}(u) < \pi^{-1}(v)$ . The number of edges of  $H(\pi)$  incident to each of the two nodes  $e_1, e_2$  that correspond to  $e$  is  $\pi^{-1}(v) - 2 = \pi^{-1}(u) + [\pi^{-1}(v) - \pi^{-1}(u)] - 2 = \pi^{-1}(u) + \delta(e, \pi) - 2$ ; thus, the number of edges of  $H(\pi)$  incident to  $e_1$  and  $e_2$  is  $\pi^{-1}(v) + \pi^{-1}(u) + \delta(e, \pi) - 4$ . Consequently,

$$\begin{aligned} h(\pi) &= \sum_{v \in N} \sum_{x \in R(v)} [\pi^{-1}(v) - 1] + \sum_{e=(u,v) \in E} [\pi^{-1}(v) + \pi^{-1}(u) + \delta(e, \pi) - 4] \\ &= \sum_{v \in N} (n - d(v))(\pi^{-1}(v) - 1) + \sum_{v \in N} d(v)\pi^{-1}(v) + \sum_{e \in E} \delta(e, \pi) - 4m \\ &= \sum_{v \in N} n[\pi^{-1}(v) - 1] + \sum_{v \in N} d(v) + c(\pi) - 4m \\ &= c(\pi) + \frac{n^2(n-1)}{2} - 2m, \end{aligned}$$

since  $\sum_{v \in N} d(v) = 2m$ , and

$$\sum_{v \in N} [\pi^{-1}(v) - 1] = 0 + 1 + 2 + \cdots + (n-1) = \frac{n(n-1)}{2}. \quad \square$$

**THEOREM 1.** *The minimum fill-in problem is NP-complete.*

*Proof.* Follows from Lemmas 2 and 3.  $\square$

## REFERENCES

- [GJ] M. R. GAREY AND D. S. JOHNSON, *Computers and Intractability: A Guide to the Theory of NP-completeness*, W. H. Freeman, San Francisco, 1979.
- [GJS] M. R. GAREY, D. S. JOHNSON AND L. STOCKMEYER, *Some simplified NP-complete graph problems*, Theoret. Comp. Sci., 1 (1976), pp. 237–267.
- [R] D. J. ROSE, *A graph-theoretic study of the numerical solution of sparse positive definite systems of linear equations*, in Graph Theory and Computing, R. Read, ed., Academic Press, New York, 1973, pp. 183–217.
- [RT] D. J. ROSE AND R. E. TARJAN, *Algorithmic aspects of vertex elimination on directed graphs*, SIAM J. Appl. Math., 34 (1978), pp. 176–197.
- [RTL] D. J. ROSE, R. E. TARJAN AND G. S. LUEKER, *Algorithmic aspects of vertex elimination on graphs*, SIAM J. Comput., 5 (1976), pp. 266–283.
- [Y] M. YANNAKAKIS, *Node-deletion problems on bipartite graphs*, SIAM J. Comput., 10 (1981).