
Path Integral Study Note

Refer to U.Mosel, Path Integrals in Field Theory

Tae Geun Kim

2018-07-27

The Path Integral in Quantum Theory

1.1 Propagator of the Schrödinger Equation

The non-relativistic Schrödinger equation with one dimensional potential $V(x)$:

$$H\psi(x, t) = -\frac{\hbar^2}{2m} \frac{\partial^2 \psi(x, t)}{\partial x^2} + V(x)\psi(x, t) = i\hbar \frac{\partial \psi(x, t)}{\partial t} \quad (1)$$

What can obtain from schrödinger equation?

- the wave function $\psi(x, t)$ if we know $\psi(x, t_0)$ at the earlier time $t_0 < t$.

Rewrite Schrödinger eq :

$$\left(i\hbar \frac{\partial}{\partial t} - H \right) \psi(x, t) = 0 \quad (2)$$

Next, we consider the function $K(x, t; x_i, t_i)$ which is defined as a solution of the equation :

$$\left(i\hbar \frac{\partial}{\partial t} - H \right) K(x, t; x_i, t_i) = i\hbar \delta(x - x_i) \delta(t - t_i) \quad (3)$$

What is K ?

- the **Green's function** of the Schrödinger equation
- or often called the **propagator**

Initial condition of K :

$$K(x, t_i + 0; x_i, t_i) = \delta(x - x_i) \quad (4)$$

The solution for $t > t_i$ (Huygen's principle) of the Schrödinger equation :

$$\psi(x, t) = \int K(x, t; x_i, t_i) \psi(x_i, t_i) dx_i \quad (5)$$