

COMP 353 Databases

Design Theory for Relational Databases
Functional Dependencies
Schema Refinement (Decomposition)
Normal Forms

1

Functional Dependencies (FDs)

- A **functional dependency** (FD) is a kind of **constraint**
- Suppose R is a relation schema and $X, Y \subseteq R$.
A FD on R is a statement of the form $X \rightarrow Y$, which asserts:
"For every "legal/valid" instance r of R , and for all pairs of tuples t_1 and t_2 in r , if t_1 and t_2 **agree** on the values in X , then t_1 and t_2 **agree** also on the values in Y ."
In symbols: $\forall t_1, t_2 \in r: t_1[X] = t_2[X] \rightarrow t_1[Y] = t_2[Y]$.
- We read $X \rightarrow Y$ as:
 X (functionally) **determines** Y (or Y is **determined by** X)
- We say that the FD: $X \rightarrow Y$ is **relevant to** R if $XUY \subseteq R$.

2

Functional Dependencies

- Consider the relation schema:
Star (name, SIN, street, city, postalCode, phone)
- Since we know the semantics of this relation from the design phase, we can answer the following question:
 - What are the functional dependencies on **Star**?
- Note that in general, FDs on a relation R may not be determined based on a given instance of R !

3

Functional Dependencies

- Consider the relation:
Movie (title, year, length, filmType)
- What are the FD's on the **Movie** relation?
We use the semantics of this relation to answer.

4

Keys

- The concept of FD generalizes the concept of key. How?
 - Let $X \subseteq R$. Then X is a key of R iff $X \rightarrow R$
- X is a **(candidate) key** of R (or a key, for short) if
 1. $X \rightarrow R$. That is, attributes in X functionally determine **all** the attributes of R
 2. No proper subset of X is key, i.e., a candidate key must be **minimal**
- Is {title, year, filmType} a key for relation **Movie**?
- A set of attributes that contains a key is called a **superkey** (that is, a **superset of a key**)
 - Note that every key is a **superkey**, but not vice versa

5

Functional Dependencies

- $X \rightarrow Y$ is called a **functional dependency** because, in principle, there is a function that takes a list of values, one for each attribute in X , and returns at most one value (i.e., a *unique* value or no value at all) for the attributes in Y

6

Functional Dependencies

- Consider the relation:
Movie (title, year, length, filmType, studioName, starName)
- What are the functional dependencies?
 - $\{title, year\} \rightarrow length$
 - $\{title, year\} \rightarrow filmType$
 - $\{title, year\} \rightarrow studioName$
 - $\rightarrow \{title, year\} \rightarrow \{length, filmType, studioName\}$
- Note: $\{title, year\} \rightarrow starName$ does not hold
- What is the key of the **Movie** relation?

7

Trivial FD's

- An FD $X \rightarrow Y$ is said to be **trivial** if $Y \subseteq X$.
 - For example: $\{title, year\} \rightarrow title$ is a trivial FD
- Otherwise, the FD is called **nontrivial**
 - For example: $\{title, year\} \rightarrow length$ is a nontrivial FD

8

Functional Dependencies

- Why are we interested in functional dependencies?

9

Redundancy Problem

- Redundancy – a “piece” of information is unnecessarily repeated in different tuples in a relation
- Recall that **redundancy** is the main source of problems:
 - Storage waste
 - Some information stored repeatedly
 - Update anomalies
 - If a copy of such information is updated, an inconsistency may arise unless all its copies are updated
 - Insertion anomalies
 - Unless we allow nulls, it may not be possible to store some information unless we have all the information to store
 - Deletion anomalies
 - Deleting some information may result in losing some other information (which we don't want to lose)

10

Is this a good design for relation R?

Name	SSN	Phone
Fred	123-321-99	(201) 555-1234
Fred	123-321-99	(201) 572-4312
Joe	909-438-44	(908) 464-0028
Mary	938-401-54	(201) 555-1234

The only FD on R is: **SSN \rightarrow Name**
Therefore, the only key of R is: **{SSN, Phone}**

11

What about this design, replacing R with R1 and R2?

R1	SSN	Name
	123-321-99	Fred
	909-438-44	Joe
	938-401-54	Mary


R2	SSN	Phone
	123-321-99	(201) 555-1234
	123-321-99	(201) 572-4312
	909-438-44	(908) 464-0028
	938-401-54	(201) 555-1234

$D = \{R1(SSN, Name), R2(SSN, Phone)\}$
FD's on R1 and R2?

12

Another Example

- Suppose each star has a representing *firm* and each *firm* has one *spokes person*



Relation Instance:

Name	Address	RepresentingFirm	SpokesPerson
Carrie Fisher	123 Maple	Star One	Joe Smith
Harrison Ford	789 Palm rd.	Star One	Joe Smith
Mark Hamill	456 Oak rd.	Movies & Co	Mary Johns

13

Redundancy Problem

What is the role of FDs in detecting redundancy?

- Consider the relation scheme $R(A, B, C)$
 - Suppose no (nontrivial) FD holds on R
 - There is no redundancy in any instance r of R .
 - Now suppose FD: $A \rightarrow B$ holds on R
 - If several tuples have the same A value \rightarrow they must all have the same B value; otherwise this FD is violated
- Presence of some FDs in a relation suggests possibility of redundancy

14

Implications of FDs and Reasoning

- Consider relation $R(A, B, C)$ with the set of FDs: $F = \{A \rightarrow B, B \rightarrow C\}$
- We can deduce from F that $A \rightarrow C$ also holds on R .
How? Apply the definition...
- To detect possible data redundancy, is it **necessary** to consider "all" the FDs (implicit and explicit)?
 - As shown above, there might be some additional hidden (nontrivial) FDs "implied" by a given set of FD's

15

Implications of FDs

- Defn: If a relation instance r satisfies every FD in a given set F of FD's, then we say that r satisfies F .
 - In this case, we also say that r is a *legal/valid instance*.
- Given $\langle R, F \rangle$, we say that F **implies** a FD $X \rightarrow Y$, if *every* instance r of R that satisfies F also satisfies $X \rightarrow Y$.
Formally, we express this as: $F \models X \rightarrow Y$.
We may also say that $X \rightarrow Y$ follows from F .
- To show $F \not\models X \rightarrow Y$, we may give a counter-example, i.e., an instance r of R that satisfies F but not $X \rightarrow Y$.

16

FDs Implication (Cont'd)

- Consider $R(A_1, A_2, A_3, A_4, A_5)$ with FDs: $F = \{A_1 \rightarrow A_2, A_2 \rightarrow A_3, A_2 A_3 \rightarrow A_4, A_2 A_3 A_4 \rightarrow A_5\}$
- Prove that $F \not\models A_5 \rightarrow A_1$
- Solution method:** Provide a counter-example; give a relation instance r of R that satisfies every FD in F but not $A_5 \rightarrow A_1$
- A desired instance r of R :**

	A_1	A_2	A_3	A_4	A_5
t_1 :	0	1	1	1	1
t_2 :	1	1	1	1	1

17

Closure of a set F of FDs

- Defn: The **closure** of F , denoted by F^+ , is the set of every FD: $X \rightarrow Y$ that is implied by F .
- How can we determine F^+ ?
 - Clearly, F^+ includes F and possibly some more FDs
 - To answer the question we need to *reason* about FDs

18

Equivalence of two sets of FD's

- Let R be a relation schema, and S, T be sets of FDs on R .
 - Defn: we say S **covers** T ($S \models T$) if for every instance r of R , whenever r satisfies (every FD in) S , r also satisfies T .
 - Defn: T and S are **equivalent** ($S \equiv T$) iff $S \models T$ and $T \models S$.
 - Note: F and F^+ are equivalent.
- Example: Suppose $R = \{A, B, C\}$, and
 $S = \{A \rightarrow B, B \rightarrow C, A \rightarrow C\}$
 $T = \{A \rightarrow B, B \rightarrow C\}$
 We can show that $S \equiv T$.

19

Armstrong's Axioms [1974]

- R is a relation schema, and X, Y, Z are subsets of R .
- Reflexivity**
 - If $Y \subseteq X$, then $X \rightarrow Y$ (trivial FDs)
- Augmentation**
 - If $X \rightarrow Y$, then $XZ \rightarrow YZ$, for every Z
- Transitivity**
 - If $X \rightarrow Y$ and $Y \rightarrow Z$, then $X \rightarrow Z$
- These are **sound** and **complete** inference rules for FDs

20

Additional rules / axioms

Other useful rules that follow from Armstrong Axioms:
 Suppose X, Y, Z , and W are sets of attributes.

- Union (Combining) Rule**
 - If $X \rightarrow Y$ and $X \rightarrow Z$, then $X \rightarrow YZ$
- Decomposition (Splitting) Rule**
 - If $X \rightarrow YZ$, then $X \rightarrow Y$ and $X \rightarrow Z$
- Pseudotransitivity Rule**
 - If $X \rightarrow Y$ and $WY \rightarrow Z$, then $XW \rightarrow Z$

21

Example – Discovering hidden FD's

- Consider $R = \{A, B, C, G, H, I\}$ with the FDs:
 $F = \{A \rightarrow B, A \rightarrow C, CG \rightarrow H, CG \rightarrow I, B \rightarrow H\}$
 - Using Armstrong's rules, we can derive more FDs
 - Since $A \rightarrow B$ and $B \rightarrow H$, then $A \rightarrow H$, by **transitivity**
 - Since $CG \rightarrow H$ and $CG \rightarrow I$, then $CG \rightarrow HI$, by **union**
 - Since $A \rightarrow C$ then $AG \rightarrow CG$, by **augmentation**
- Now, since $AG \rightarrow CG$ and $CG \rightarrow I$, then $AG \rightarrow I$, by **transitivity** (and in a similar way, we get $F \models AG \rightarrow H$)
- Many trivial dependencies can be derived(!) by **augmentation**

22

Implication Problem

- Given a set F of FDs, does $X \rightarrow Y$ follow from F ?
 - In other words: is $X \rightarrow Y$ in the closure of F ?
- (In symbols, does $F \models X \rightarrow Y$ hold, or is $X \rightarrow Y \in F^+$ true?)
- How to answer this question?
 - Compute the closure of F & check if it includes $X \rightarrow Y$
 - What is the problem with this approach?
 - Computing F^+ is **expensive!** Is there a better solution?

23

Closure of a Set of Attributes

- Given $\langle R, F \rangle$; Let $X \subseteq R$.
- The **closure of X under F** is the set of all attributes Y in R that are determined by X . This yields $X \rightarrow Y$, i.e., every valid instance of R (that satisfies F) also satisfies $X \rightarrow Y$
- We denote the **closure of a set of attributes X under F** by X^+_F
 - When F is known, we simply write X^+ (and omit F)
 - Closure of $\{A_1, A_2, \dots, A_n\}$ is denoted $\{A_1, A_2, \dots, A_n\}^+$
 - Note that $X \subseteq X^+$, for any set X of attributes (because $X \rightarrow X$)

24

Computing the Closure of Attributes

- Given a set F of FD's and a set X of attributes, how to compute the closure of X w.r.t. F ?
 - Starting with set $X^+ = X$, we repeatedly expand X^+ by adding the RHS Z for every FD: $W \rightarrow Z$ in F , if the LHD W is already in X^+ .
 - This process terminates when X^+ could not be expanded further.
 - This process is expressed as an algorithm in the next slide.

25

An Algorithm to Compute X^+ under F

```

 $X^+ \leftarrow X$  (initialization step)
repeat
  for each FD  $W \rightarrow Z$  in  $F$  do:
    if  $W \subseteq X^+$  then
       $X^+ \leftarrow X^+ \cup Z$  // add  $Z$  to the result
until  $X^+$  does not change
    
```

Complexity? In the worst case, how many times the "repeat" statement may be executed?

26

Examples

- Consider a relation schema $R = \{A, B, C, D, E, H\}$ with the FD's $F = \{AB \rightarrow C, BC \rightarrow AD, D \rightarrow E, CH \rightarrow B\}$
- Suppose $X = \{A, B\}$. Compute X^+
- Execution result at each iteration:
 - Initially, $X^+ = \{A, B\}$
 - Using $AB \rightarrow C$, we get $X^+ = \{A, B, C\}$
 - Using $BC \rightarrow AD$, we get $X^+ = \{A, B, C, D\}$
 - Using $D \rightarrow E$, we get $X^+ = \{A, B, C, D, E\}$
 - No more change to X^+ is possible.
- $\rightarrow X^+ = \{A, B\}^+ = \{A, B, C, D, E\}$
- Does the order in which FD's appear in F affects the computation?

27

Implication Problem Revisited

- Given a set of FD's F , does an FD: $X \rightarrow Y$ follow from F ?
 - That is, is FD $X \rightarrow Y$ in F^+ ?
- To answer this, we can compute X^+ under F , and check if Y is in X^+ or not
 - If yes, then the answer is positive! ($F \models X \rightarrow Y$ 😊)
 - Otherwise, it is negative ($F \not\models X \rightarrow Y$ ☹️)

28

Example

- Consider $\langle R, F \rangle$ where $R = \{A, B, C, D, E, H\}$ and $F = \{AB \rightarrow C, BC \rightarrow AD, D \rightarrow E, CH \rightarrow B\}$
- Does $AB \rightarrow D$ follow from F ?
- Two steps:
 - Compute $\{A, B\}^+ = \{A, B, C, D, E\}$
 - Check if $D \in \{A, B\}^+$
- So, here we conclude that $AB \rightarrow D$ is implied by F

29

Example

- Consider a relation schema $R = \{A, B, C, D, E, H\}$ with FDs: $F = \{AB \rightarrow C, BC \rightarrow AD, D \rightarrow E, CH \rightarrow B\}$
- True/False: Does $D \rightarrow A$ follow from F ?
- Two steps:
 - Compute $\{D\}^+ = \{D, E\}$
 - Check if $A \in \{D\}^+$
- Since $A \notin \{D, E\}$, the answer is NO, i.e., $F \not\models D \rightarrow A$

30

Closures and Keys

- Consider a case where X^+ includes all the attributes of a relation R
 - Clearly, X is a (super) key of R
 - To check if X is a candidate key of R , we should check 2 things:
 1. If X^+ is a superkey R , i.e., when $X^+ = R$, and
 2. If no proper subset of X is a key, i.e., $\forall A \in X: (X - \{A\})^+ \neq R$
- To find the keys of a relation, we can use the algorithm on slide 26
- This would be exponential in the number of attributes! Can do better?
- Knowledge about keys is essential to understand "Normal forms."