

Assignment 1: Abstract and Introduction Summary

Abstract :

Tiny non-coding RNAs generated by eukaryotes serve as specificity determinants for various gene-regulatory complexes. MicroRNAs (miRNAs), endogenous short interfering RNAs (siRNAs), and Piwi-associated RNAs are examples of these (piRNAs). Small RNA-seq, a variant RNA-seq approach based on highly parallel sequencing, was used to annotate and quantify the data. The alignment of small RNA-seq data to a reference genome is a crucial step in the research process. Due to their small size (20-30 nts depending on the organism and sub-type) and proclivity to stick together, Reads that align equally well to multiple genomic locations are very common, particularly if they come from multi-gene families or repeated regions. Typical approaches to dealing with multi-mapped narrow RNA-seq reads trade off precision for sensitivity. The method 'butter' combines accuracy and sensitivity by using an iterative approach to place multi-mapped reads, where the choice between possible positions is made each time. Dictated by the densities of more confidently matched reads in the surrounding area. In comparison to other small RNA-seq aligners, Butter performs well.

Introduction :

Small RNA-seq is a type of RNA-seq that uses directional RNA ligation, cDNA synthesis, and sequencing to capture small RNAs, usually in the 15-40 nt range. It's an effective tool for finding, annotating, and quantifying small RNA molecules. Multi-mapped reads, known as reads with more than one equally probable alignment location, are a common problem in RNA-seq data alignment, particularly for small data sets. Some of solution is to (completely disregard multi-mapped reads, place multi-mapped reads randomly at one of the possible positions. simply report all possible alignment positions).

Larger RNA precursors are converted into piRNAs, endogenous siRNAs, and miRNAs. Processing is heterogeneous to differing degrees, resulting in several

distinct small RNAs that, when properly mapped to the reference genome, provide a cluster of alignments at the origin locus.

I present butter (Bowtie Using iterative placement of Repetitive small rnas), a technique that employs an iterative approach based on uniquely and low-level multi-mapped reads to improve the precision of small RNA-seq genomic alignments.