1. The probability of a leap year selected at random contain 53 Sunday is:					
(a) 53/366 (b) 1/7 (c) 2/7 (d) 53/36	65				
2. A bag contains 3 red and 2 blue marbles. A marble is					
random. The probability of drawing a black ball is :					
(a) 3/5 (b) 2/5 (c) 0/5 (d) 1/5					
3. The probability that it will rain tomorrow is 0.85. What	at is the				
probability that it will not rain tomorrow					
(a) 0.25 (b) 0.145 (c) 3/20 (d) none	of these				
4. What is the probability that a number selected from t	he numbers				
(1, 2, 3,,15) is a multiple of 4?					
(a) 1/5 (b) 4/5 (c) 2/15 (d) 1/3					
5. What are the total outcomes when we throw three co	oins?				
(a) 4 (b) 5 (c) 8 (d) 7					
6. The probability that a prime number selected at rand	lom from the				
numbers (1,2,3,35) is :					
(a) 12/35 (b) 11/35 (c) 13/35 (d) nor					
7. The sum of the probability of an event and non event					
(a) 2 (b) 1 (c) 0 (d) none of these					
8. The following probabilities are given; choose the cor	rect answer				
for that which is not possible.					
(a) 0.15 (b) 2/7 (c) 7/5 (d) none					
9. If three coins are tossed simultaneously, than the pro	obability of				
getting at least two heads, is:					
(a) $1/4$ (b) $3/8$ (c) $\frac{1}{2}$ (d) $1/4$					
10. A letter is chosen at random from the letters of the					
♦ ASSASSINATION ♦ . The probability that the letter che					
(a) 6/13 (b) 7/13 (c) 1 (d) no	one of these.				
	_				
11. A dice is thrown. Find the probability of getting an ev	ven number.				
(A) 2/3 (B) 1 (C) 5/6 (D) 1/2					
12. Two coins are thrown at the same time. Find the probability of getting both heads.					
(A) 3/4 (B) 1/4 (C) 1/2 (D) 0					
13. Two dice are thrown simultaneously. The probability	of getting a				

sum of 9 is:

1

(A) 1/10	(B) 3/10	(C) 1/9	(D) 4/	9			
14. 100 cards are numbered from 1 to 100. Find the probability of getting a prime number.							
(A) 3/4	(B) 27/50	(C) 1/4	(D)	29/100			
_	a blue ball is o	double that of a		f the probability In the number of			
16. A box of 600 bulbs contains 12 defective bulbs. One bulb is taken out at random from this box. Then the probability that it is non-defective bulb is: (A) 143/150 (B) 147/150 (C) 1/25 (D) 1/50							
17. Cards n	narked with nu oughly. One ca lity that the nu	umbers 2 to 10 ard is drawn fro umber on card (C) 3/10	1 are placed om this box r is a perfect s	in a box and and andomly, then square.			
18. What is (A) 1/7	the probabili (B) 53/366	ty of getting 53 (C) 2/7	•	• •			
19. A card is drawn from a well shuffled deck of 52 cards. Find the probability of getting a king of red suit. (A) $1/26$ (B) $3/26$ (C) $7/52$ (D) $1/13$							
equally like 1,2,312	ly to come to ,then the prob	onsists of spinn rest pointing to pability that it w (C) 7/12	o one of the r	number n odd number is:			
its outcome result i.e. the probability	e each time. A nree heads or	ryan wins if all three tails and I lose the game	the tosses g loses otherw e.				

22. Riya and Kajal are friends. Probability that both will have the same birthday is the same birthday is:							
•		(C) 1/365	(D) 1/133225				
23. A number x is chosen at random from the numbers -2, -1, 0, 1, 2. Then the probability that $x^2 < 2$ is? (A) $1/5$ (B) $2/5$ (C) $3/5$ (D) $4/5$							
a marble is dr red is 2/3, the	24. A jar contains 24 marbles. Some are red and others are white. If a marble is drawn at random from the jar, the probability that it is red is 2/3, then the number of white marbles in the jar is: (A) 10 (B) 6 (C) 8 (D) 7						
Then the prob	25. A number is selected at random from first 50 natural numbers. Then the probability that it is a multiple of 3 and 4 is: (A) $7/50$ (B) $4/25$ (C) $1/25$ (D) $2/25$						
with n dots sh	26. Consider a dice with the property that that probability of a face with n dots showing up is proportional to n. The probability of face showing 4 dots is?						
a) ¹ / ₇	b) $\frac{5}{42}$	c) $\frac{1}{21}$	d) $\frac{4}{21}$				
			hes are 50, 70, 82,				
	he standard devia b) 25.49	c) 25.29	d) 25.69				
28. Find median and mode of the messages received on 9 consecutive days 15, 11, 9, 5, 18, 4, 18, 13, 17.							
a) 13, 15			d) 13, 16				
29. A coin is tossed up 4 times. The probability that tails turn up in 3 cases is							
a) $\frac{1}{2}$ 30. X is a vari	b) $^1\!/_3$ ate between 0 an	$\frac{\text{c) }^{1}/_{4}}{\text{cd 3. The value of }}$					
a) 8	b) 7	c) 27 d) 9				
31. The random variables X and Y have variances 0.2 and 0.5 respectively. Let Z= 5X-2Y. The variance of Z is?							

probability ⁴	?	•	n one is not poss	sible in
a) $P(x) = 1$ c) $P(x) = 0$.	b) ∑ x 5 <mark>d) P(</mark> 3	P(x) = 3 x) = -0.5		
33.If E(x) = a) 2	= 2 and E(z) = 4 b) 6	I, then E(z - c) 0		sufficient data
34.The cov	ariance of two	independe	nt random variak	ole is
a) 1	b) 0	c) - 1	d) Ur	ndefined
35.If Σ P(x a) 0) = k² - 8 then, b) 1	, the value o <mark>c) 3</mark>		sufficient data
• •	0.5 and x = 4, b) 0.5	• •	? d) 2	
37.In a discission always?	crete probabilit	ty distributio	on, the sum of al	l probabilities
a) 0	b) Infinite	c) 1	d) Un	defined
-	robability of hi	tting the tar	get is 0.4, find n	nean and
variance. a) 0.4, 0.24	b) 0.6,	0.24	c) 0.4, 0.16	d) 0.6, 0.16
-	% and if 10 bo		pped from a place opped, find mear 0.4, 0.16	
40. Find the a) 2	e mean of toss b) 4	ing 8 coins. c) 8		,

c) 5

d) 7

a) 3

b) 4

-				•		nd variance and variand	
					s given b (2) - (E(X	•	d) (E(X))2
					iven by 2) - (E(X)		d) (E(X))2
44.N a) 0	lean of	a consta <mark>b) a</mark>	nt 'a' is _	c) a/2		d) 1	
45.Variance of a constant 'a' is . a) 0							
46.Find the mean and variance of X?							
	Х	0	1	2	3	4	
	f(x)	1/9	2/9	3/9	2/9	1/9	
a) 2,	4/3	b)	3, 4/3		c) 2, 2/3		d) 3, 2/3
47.F	ind the	expectat	ion of a r	andom	variable)	(?	

	Х	0	1	2	3	
	f(x)	1/6	2/6	2/6	1/6	
a) ().5		b) 1.5		c) 2.5	d) 3.5

48. In a Binomial Distribution, if p, q and n are probability of success, failure and number of trials respectively then variance is given by

b) npq

c) np2q

d) npq2

- 49. If 'X' is a random variable, taking values 'x', probability of success and failure being 'p' and 'q' respectively and 'n' trials being conducted, then what is the probability that 'X' takes values 'x'? Use **Binomial Distribution.**
- a) P(X = x) = nCx px qx
- b) P(X = x) = nCx px q(n-x)
- c) P(X = x) = xCn qx p(n-x)
- d) P(x = x) = xCn pn qx
- 50. If 'p', 'q' and 'n' are probability pf success, failure and number of trials respectively in a Binomial Distribution, what is its Standard **Deviation?**

- a) \sqrt{np} b) \sqrt{pq} c) (np)2 d) \sqrt{npq}