

Thm: If f is differentiable at $x=c$,
then f must be continuous at c .

Proof: Since f is differentiable at c ,
 $f'(c) = \lim_{h \rightarrow 0} \frac{f(c+h) - f(c)}{h}$ exists.

$$= \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c} \quad (\text{put } x - c = h)$$

Now, $f(x) = [f(x) - f(c)] + f(c)$
 $= \left[\frac{f(x) - f(c)}{x - c} \right] (x - c) + f(c)$
 for any $x \neq c$

$$\therefore \lim_{x \rightarrow c} f(x) = \left[\lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c} \right] \lim_{x \rightarrow c} (x - c) + f(c)$$

$$= f'(c) \cdot 0 + f(c) = f(c)$$

$\Rightarrow f$ is continuous at c .

Remark: The converse of the previous theorem is not true.

Example: $f(x) = |x|$ is continuous at $x=0$ but not differentiable at $x=0$.

$$\frac{f(x) - f(0)}{x - 0} = \frac{|x|}{x} = \begin{cases} 1 & \text{if } x > 0 \\ -1 & \text{if } x < 0 \end{cases}$$

$$\lim_{x \rightarrow 0^+} \frac{f(x) - f(0)}{x - 0} = 1$$

$$\lim_{x \rightarrow 0^-} \frac{f(x) - f(0)}{x - 0} = -1$$

$\Rightarrow \lim_{x \rightarrow 0} \frac{f(x) - f(0)}{x - 0}$ does not exist
 $\Rightarrow f$ is not diff. at $x=0$.

Exercise: Check the continuity and differentiability of the following functions at $x=0$.

$$(i) \quad f(x) = \begin{cases} x \sin\left(\frac{1}{x}\right), & x \neq 0 \\ 0, & x = 0 \end{cases}$$

$$(ii) \quad g(x) = \begin{cases} x^2 \sin\left(\frac{1}{x}\right), & x \neq 0 \\ 0, & x = 0 \end{cases}$$

Some rules of differentiability:

$$(i) \quad (f+g)'(x) = f'(x) + g'(x)$$

$$(ii) \quad (f-g)'(x) = f'(x) - g'(x)$$

$$(iii) \quad (cf)'(x) = cf'(x)$$

(iv) Product rule:

$$(fg)'(x) = f'(x)g(x) + f(x)g'(x)$$

(v) Quotient rule:

$$\left(\frac{f}{g}\right)'(x) = \frac{f'(x)g(x) - f(x)g'(x)}{(g(x))^2}$$

Created with Doceri

(vi) Chain rule:

$$(f \circ g)'(x) = f'(g(x)) \cdot g'(x)$$

Example: $f(x) = \sin(x^2)$

$$f'(x) = \cos(x^2) \cdot 2x$$

$$(vii) \quad (f^{-1})'(x) = \frac{1}{f'(f^{-1}(x))}$$

$$\text{Let } g(x) = f^{-1}(x)$$

$$\text{Then } f(g(x)) = x$$

Differentiating both sides, we get

Created with Doceri

$$f'(g(x)) \cdot g'(x) = 1$$

$$\Rightarrow g'(x) = \frac{1}{f'(g(x))}$$

$$\Rightarrow (f^{-1})'(x) = \frac{1}{f'(f^{-1}(x))}$$

Example: Derivative of $\ln x$:


Let $f(x) = e^x$. Then $f^{-1}(x) = \ln x$

$$\therefore \frac{d}{dx}[\ln x] = (f^{-1})'(x) = \frac{1}{f'(f^{-1}(x))} = \frac{1}{e^{\ln x}} = \frac{1}{x}$$

Created with Doceri



Local minima and local maxima:



x_1 & x_3 are local max. of f
 x_2 & x_4 are local min. of f .

Created with Doceri



Defn: We say $x=a$ is a local minimum of f if there exists $\delta > 0$ s.t.
 $f(a) \leq f(x) \quad \forall x \in (a-\delta, a+\delta)$


Similarly $x=a$ is a local maximum if $\exists \delta > 0$ s.t.
 $f(a) \geq f(x) \quad \forall x \in (a-\delta, a+\delta)$

Created with Doceri



Theorem: If $x=a$ is a local min. or local max. of f and if f is differentiable at $x=a$, then $\boxed{f'(a) = 0}$.

$f'(0) = 0$ for $f(x) = x^3$
 But $x=0$ is neither a local min. nor a local max. of f .




Created with Doceri



Rolle's Theorem :

Suppose $f(x)$ is continuous on $[a, b]$
and it is differentiable on (a, b) .
If $f(a) = f(b)$, then $\exists c \in (a, b)$
such that $f'(c) = 0$



Created with Doceri



Created with Doceri

