

Sante Speech Health Summary

Report ID: sum_f332a1711822 | Generated: 2026-02-15 14:22 UTC

Session: conversation | Duration: 0.0s

Executive Summary

Data quality is limited; treat outputs as coarse exploratory estimates only.

- Perceived Vocal Age / Gender Proxy: exploratory-moderate (score 68, confidence 58%)
- Slurred Speech / Intoxication Likelihood: exploratory-moderate (score 61, confidence 58%)
- Voice Strain / Respiratory Effort Indicator: exploratory-low (score 49, confidence 58%)

Estimates

Mood / Depression Risk Signal | exploratory-low | score 32 | confidence 58%

Suggestion: Track trend across multiple days and pair with self-reported mood scales.

Evidence: Mean pause length 1.24s; Speaking-rate proxy 1.08 peaks/s; Disfluency ratio 0.29

Limitations: Speech-only estimate; mood state can be situational.; Exploratory speech biomarkers are probabilistic, not diagnostic outcomes.

Aphasia-like Language Pattern Flag | exploratory-low | score 36 | confidence 58%

Suggestion: If persistent, run structured language tasks and formal aphasia batteries.

Evidence: Phoneme coverage 277 tokens; Flagged disfluencies 79; Voiced segment rate 0.71/s

Limitations: Cannot subtype aphasia from this short speech sample.; Exploratory speech biomarkers are probabilistic, not diagnostic outcomes.

Perceived Vocal Age / Gender Proxy | exploratory-moderate | score 68 | confidence 58%

Suggestion: Use only as demographic context; avoid identity or eligibility decisions.

Evidence: F0 mean 26.83 st; Jitter 0.015

Limitations: Only estimates broad vocal age band, not chronological age.; Exploratory speech biomarkers are probabilistic, not diagnostic outcomes.

Slurred Speech / Intoxication Likelihood | exploratory-moderate | score 61 | confidence 58%

Suggestion: Confirm with behavioral checks and contextual history before acting.

Evidence: Jitter 0.015; Shimmer 1.176 dB; Mean pause length 1.24s

Limitations: Medication, fatigue, and microphone clipping can mimic this pattern.; Exploratory speech biomarkers are probabilistic, not diagnostic outcomes.

Cognitive Load / Fatigue Proxy | exploratory-low | score 47 | confidence 58%

Suggestion: Repeat under controlled conditions and compare with baseline sessions.

Evidence: Speech pacing proxy 1.08 peaks/s; Pause duration 1.24s; Disfluency ratio 0.29

Limitations: Single-session fatigue score is highly context-dependent.; Exploratory speech biomarkers are probabilistic, not diagnostic outcomes.

Voice Strain / Respiratory Effort Indicator | exploratory-low | score 49 | confidence 58%

Suggestion: Consider hydration, rest, and ENT/SLP follow-up if trend worsens.

Evidence: HNR 6.84 dB; Shimmer 1.176 dB; Jitter 0.015

Limitations: Breathing pattern cannot be directly measured from this sample.; Exploratory speech biomarkers are probabilistic, not diagnostic outcomes.

Transcript Snapshot

Oh, I'm feeling good, Dante. How are you? Oh yeah, I've been coding up a hackathon project. The project has been going good so far. I have been integrating multiple features and solving multiple bugs. Let's see. Coordination and

solving multiple integration bugs. Yeah, that has been the most challenging part. The coordination with the team is through just mobile communication and what tasks have been assigned and solving them. Thank you.

Research-only estimate report. It is not a medical diagnosis. Use in combination with formal clinical assessment and environmental quality checks.