

Sante Speech Health Summary

Report ID: sum_17e8ebba7951 | Generated: 2026-02-15 14:40 UTC

Session: conversation | Duration: 0.0s

Executive Summary

Data quality is limited; treat outputs as coarse exploratory estimates only.

- Perceived Vocal Age / Gender Proxy: exploratory-moderate (score 68, confidence 58%)
- Slurred Speech / Intoxication Likelihood: exploratory-moderate (score 56, confidence 58%)
- Voice Strain / Respiratory Effort Indicator: exploratory-low (score 48, confidence 58%)

Estimates

Mood / Depression Risk Signal | exploratory-low | score 24 | confidence 58%

Suggestion: Track trend across multiple days and pair with self-reported mood scales.

Evidence: Mean pause length 0.97s; Speaking-rate proxy 1.20 peaks/s; Disfluency ratio 0.38

Limitations: Speech-only estimate; mood state can be situational.; Exploratory speech biomarkers are probabilistic, not diagnostic outcomes.

Aphasia-like Language Pattern Flag | exploratory-low | score 41 | confidence 58%

Suggestion: If persistent, run structured language tasks and formal aphasia batteries.

Evidence: Phoneme coverage 332 tokens; Flagged disfluencies 127; Voiced segment rate 0.81/s

Limitations: Cannot subtype aphasia from this short speech sample.; Exploratory speech biomarkers are probabilistic, not diagnostic outcomes.

Perceived Vocal Age / Gender Proxy | exploratory-moderate | score 68 | confidence 58%

Suggestion: Use only as demographic context; avoid identity or eligibility decisions.

Evidence: F0 mean 27.87 st; Jitter 0.014

Limitations: Only estimates broad vocal age band, not chronological age.; Exploratory speech biomarkers are probabilistic, not diagnostic outcomes.

Slurred Speech / Intoxication Likelihood | exploratory-moderate | score 56 | confidence 58%

Suggestion: Confirm with behavioral checks and contextual history before acting.

Evidence: Jitter 0.014; Shimmer 1.166 dB; Mean pause length 0.97s

Limitations: Medication, fatigue, and microphone clipping can mimic this pattern.; Exploratory speech biomarkers are probabilistic, not diagnostic outcomes.

Cognitive Load / Fatigue Proxy | exploratory-low | score 45 | confidence 58%

Suggestion: Repeat under controlled conditions and compare with baseline sessions.

Evidence: Speech pacing proxy 1.20 peaks/s; Pause duration 0.97s; Disfluency ratio 0.38

Limitations: Single-session fatigue score is highly context-dependent.; Exploratory speech biomarkers are probabilistic, not diagnostic outcomes.

Voice Strain / Respiratory Effort Indicator | exploratory-low | score 48 | confidence 58%

Suggestion: Consider hydration, rest, and ENT/SLP follow-up if trend worsens.

Evidence: HNR 7.17 dB; Shimmer 1.166 dB; Jitter 0.014

Limitations: Breathing pattern cannot be directly measured from this sample.; Exploratory speech biomarkers are probabilistic, not diagnostic outcomes.

Transcript Snapshot

I'm feeling good, Santé. How are you? I've been coding a hackathon project. We are making medical voice assistants to detect voice biomarkers. Well, obviously to reduce healthcare delivery time, voice assistance can be a good use case

for doctors. The development is going pretty good. We have resolved many bugs, integrated multiple complex features. It's multiple coordination with the teammates and keeping up with the deadline. It's just mobile communication.

Research-only estimate report. It is not a medical diagnosis. Use in combination with formal clinical assessment and environmental quality checks.