

Sante Speech Health Summary

Report ID: sum_153098cf079a | Generated: 2026-02-15 19:06 UTC

Session: conversation | Duration: 0.0s

Executive Summary

Data quality is limited; treat outputs as coarse exploratory estimates only.

- Aphasia-like Language Pattern Flag: exploratory-low (score 13)
- Sick / Tired Voice Pattern Signal: exploratory-low (score 12)
- Intoxicated Speech Pattern Signal: exploratory-low (score 8)

Estimates

Intoxicated Speech Pattern Signal | exploratory-low | score 8

Suggestion: Low signal in this sample. Monitor trend across repeated sessions and retest with a matched microphone/device/codec setup before drawing conclusions.

Evidence: Short-pause index 1.00; Slow-rate index 1.00; Loudness mean 0.000; Jitter 0.000

Limitations: Sleep deprivation, stress, and microphone clipping can mimic this pattern.; Exploratory speech biomarkers are probabilistic, not diagnostic outcomes.

Sick / Tired Voice Pattern Signal | exploratory-low | score 12

Suggestion: Low signal in this sample. Monitor trend across repeated sessions and retest with a matched microphone/device/codec setup before drawing conclusions.

Evidence: Long-pause index 0.00; Energy-drop index 0.70; Low-loudness index 1.00; HNR 0.00 dB

Limitations: Cannot separate respiratory causes from fatigue without clinical context.; Exploratory speech biomarkers are probabilistic, not diagnostic outcomes.

Stress Activation Speech Signal | exploratory-low | score 3

Suggestion: Low signal in this sample. Monitor trend across repeated sessions and retest with a matched microphone/device/codec setup before drawing conclusions.

Evidence: Speech-rate proxy 0.00 peaks/s; Short-pause index 1.00; Pressured-tempo index 0.22; Fast-rate index 0.00

Limitations: High arousal can also reflect urgency, excitement, or speaking style.; Exploratory speech biomarkers are probabilistic, not diagnostic outcomes.

Aphasia-like Language Pattern Flag | exploratory-low | score 13

Suggestion: Low signal in this sample. Monitor trend across repeated sessions and retest with a matched microphone/device/codec setup before drawing conclusions.

Evidence: Phoneme coverage 0 tokens; Flagged disfluencies 0; Voiced segment rate 0.00/s; Lexical diversity 0.00

Limitations: Cannot subtype aphasia from this short speech sample.; Exploratory speech biomarkers are probabilistic, not diagnostic outcomes.

Transcript Snapshot

(no transcription captured)

Research-only estimate report. It is not a medical diagnosis. Use in combination with formal clinical assessment and environmental quality checks.