

Sante Speech Health Summary

Report ID: sum_ad176244fbf8 | Generated: 2026-02-15 09:17 UTC

Session: conversation | Duration: 89.0s

Executive Summary

Data quality is limited; treat outputs as coarse exploratory estimates only.

- Perceived Vocal Age / Gender Proxy: exploratory-moderate (score 69, confidence 57%)
- Voice Strain / Respiratory Effort Indicator: exploratory-moderate (score 64, confidence 58%)
- Aphasia-like Language Pattern Flag: exploratory-low (score 45, confidence 58%)

Estimates

Mood / Depression Risk Signal | exploratory-low | score 41 | confidence 58%

Suggestion: Track trend across multiple days and pair with self-reported mood scales.

Evidence: Mean pause length 0.00s; Speaking-rate proxy 0.00 peaks/s; Disfluency ratio 0.00

Limitations: Speech-only estimate; mood state can be situational.; Exploratory speech biomarkers are probabilistic, not diagnostic outcomes.

Aphasia-like Language Pattern Flag | exploratory-low | score 45 | confidence 58%

Suggestion: If persistent, run structured language tasks and formal aphasia batteries.

Evidence: Phoneme coverage 0 tokens; Flagged disfluencies 0; Voiced segment rate 0.00/s

Limitations: Cannot subtype aphasia from this short speech sample.; Exploratory speech biomarkers are probabilistic, not diagnostic outcomes.

Perceived Vocal Age / Gender Proxy | exploratory-moderate | score 69 | confidence 57%

Suggestion: Use only as demographic context; avoid identity or eligibility decisions.

Evidence: F0 mean 0.00 st; Jitter 0.000

Limitations: Only estimates broad vocal age band, not chronological age.; Exploratory speech biomarkers are probabilistic, not diagnostic outcomes.

Slurred Speech / Intoxication Likelihood | exploratory-low | score 18 | confidence 58%

Suggestion: Confirm with behavioral checks and contextual history before acting.

Evidence: Jitter 0.000; Shimmer 0.000 dB; Mean pause length 0.00s

Limitations: Medication, fatigue, and microphone clipping can mimic this pattern.; Exploratory speech biomarkers are probabilistic, not diagnostic outcomes.

Cognitive Load / Fatigue Proxy | exploratory-low | score 39 | confidence 58%

Suggestion: Repeat under controlled conditions and compare with baseline sessions.

Evidence: Speech pacing proxy 0.00 peaks/s; Pause duration 0.00s; Disfluency ratio 0.00

Limitations: Single-session fatigue score is highly context-dependent.; Exploratory speech biomarkers are probabilistic, not diagnostic outcomes.

Voice Strain / Respiratory Effort Indicator | exploratory-moderate | score 64 | confidence 58%

Suggestion: Consider hydration, rest, and ENT/SLP follow-up if trend worsens.

Evidence: HNR 0.00 dB; Shimmer 0.000 dB; Jitter 0.000

Limitations: Breathing pattern cannot be directly measured from this sample.; Exploratory speech biomarkers are probabilistic, not diagnostic outcomes.

Transcript Snapshot

(no transcription captured)

Research-only estimate report. It is not a medical diagnosis. Use in combination with formal clinical assessment and environmental

quality checks.