

Comparison of particle trajectories and collision operators for collisional transport in nonaxisymmetric plasmas

Matt Landreman, (University of Maryland, MIT, ORISE)

Håkan Smith, Per Helander (IPP Greifswald)

arXiv: 1312.6058 (2013)



Max-Planck-Institut
für Plasmaphysik



Outline

- Overview of new stellarator drift-kinetic code SFINCS.
- Variants of the drift-kinetic equation, differing in the E_r terms.
- Results: the different kinetic equations lead to indistinguishable predictions when $E_r < E_r^{\text{res}} / 3$, but can differ substantially otherwise.
- Comparison of collision operators.

SFINCS (Stellarator Fokker-Planck Iterative Neoclassical Conservative Solver)

- Solves time-independent drift-kinetic equations for $f_s(\theta, \zeta, v_\perp, v_\parallel)$.

SFINCS (Stellarator Fokker-Planck Iterative Neoclassical Conservative Solver)

- Solves time-independent drift-kinetic equations for $f_s(\theta, \zeta, v_\perp, v_\parallel)$.
- Several options for terms in the kinetic equation involving E_r – “effective particle trajectories.”

SFINCS (Stellarator Fokker-Planck Iterative Neoclassical Conservative Solver)

- Solves time-independent drift-kinetic equations for $f_s(\theta, \zeta, v_\perp, v_\parallel)$.
- Several options for terms in the kinetic equation involving E_r – “effective particle trajectories.”
- Several options for collision operator, including full linearized Fokker-Planck.

SFINCS (Stellarator Fokker-Planck Iterative Neoclassical Conservative Solver)

- Solves time-independent drift-kinetic equations for $f_s(\theta, \zeta, v_\perp, v_\parallel)$.
- Several options for terms in the kinetic equation involving E_r – “effective particle trajectories.”
- Several options for collision operator, including full linearized Fokker-Planck.
- Continuum discretization, with mix of finite-difference, spectral, and pseudospectral methods.

SFINCS (Stellarator Fokker-Planck Iterative Neoclassical Conservative Solver)

- Solves time-independent drift-kinetic equations for $f_s(\theta, \zeta, v_\perp, v_\parallel)$.
- Several options for terms in the kinetic equation involving E_r – “effective particle trajectories.”
- Several options for collision operator, including full linearized Fokker-Planck.
- Continuum discretization, with mix of finite-difference, spectral, and pseudospectral methods.
- General nested flux surface geometry allowed, with interface to equilibrium data.

SFINCS (Stellarator Fokker-Planck Iterative Neoclassical Conservative Solver)

- Solves time-independent drift-kinetic equations for $f_s(\theta, \zeta, v_\perp, v_\parallel)$.
- Several options for terms in the kinetic equation involving E_r – “effective particle trajectories.”
- Several options for collision operator, including full linearized Fokker-Planck.
- Continuum discretization, with mix of finite-difference, spectral, and pseudospectral methods.
- General nested flux surface geometry allowed, with interface to equilibrium data.
- Multiple species.

SFINCS (Stellarator Fokker-Planck Iterative Neoclassical Conservative Solver)

- Solves time-independent drift-kinetic equations for $f_s(\theta, \zeta, v_\perp, v_\parallel)$.
- Several options for terms in the kinetic equation involving E_r – “effective particle trajectories.”
- Several options for collision operator, including full linearized Fokker-Planck.
- Continuum discretization, with mix of finite-difference, spectral, and pseudospectral methods.
- General nested flux surface geometry allowed, with interface to equilibrium data.
- Multiple species.
- Uses preconditioned GMRES solver (via PETSc library).

SFINCS (Stellarator Fokker-Planck Iterative Neoclassical Conservative Solver)

- Solves time-independent drift-kinetic equations for $f_s(\theta, \zeta, v_\perp, v_\parallel)$.
- Several options for terms in the kinetic equation involving E_r – “effective particle trajectories.”
- Several options for collision operator, including full linearized Fokker-Planck.
- Continuum discretization, with mix of finite-difference, spectral, and pseudospectral methods.
- General nested flux surface geometry allowed, with interface to equilibrium data.
- Multiple species.
- Uses preconditioned GMRES solver (via PETSc library).
- Closely related to the tokamak code PERFECT for finite-orbit-width neoclassical calculations in tokamak pedestals: *arXiv:1312.2148 (2013)*.

Several choices are available for the E_r terms

1. “DKES trajectories” (Incompressible ExB drift, *van Rij & Hirshman (1989)*):

$$\left(v_{\parallel} \mathbf{b} + \frac{c}{\langle B^2 \rangle} \frac{d\Phi}{d\psi} \mathbf{B} \times \nabla \psi \right) \cdot \nabla f_1 - \frac{(1 - \xi^2)}{2B} v (\nabla_{\parallel} B) \frac{\partial f_1}{\partial \xi} - C\{f_1\} = -\mathbf{v}_m \cdot \nabla \psi \frac{\partial f_M}{\partial \psi}$$

$$\xi = v_{\parallel} / v$$

Several choices are available for the E_r terms

1. “DKES trajectories” (Incompressible ExB drift, *van Rij & Hirshman (1989)*):

$$\left(v_{\parallel} \mathbf{b} + \frac{c}{\langle B^2 \rangle} \frac{d\Phi}{d\psi} \mathbf{B} \times \nabla \psi \right) \cdot \nabla f_1 - \frac{(1 - \xi^2)}{2B} v (\nabla_{\parallel} B) \frac{\partial f_1}{\partial \xi} - C\{f_1\} = -\mathbf{v}_m \cdot \nabla \psi \frac{\partial f_M}{\partial \psi}$$

$$\xi = v_{\parallel} / v$$

2. “Partial trajectories” (Correct ExB drift):

$$\left(v_{\parallel} \mathbf{b} + \frac{c}{B^2} \frac{d\Phi}{d\psi} \mathbf{B} \times \nabla \psi \right) \cdot \nabla f_1 - \frac{(1 - \xi^2)}{2B} v (\nabla_{\parallel} B) \frac{\partial f_1}{\partial \xi} - C\{f_1\} = -\mathbf{v}_m \cdot \nabla \psi \frac{\partial f_M}{\partial \psi}$$

Several choices are available for the E_r terms

1. “DKES trajectories” (Incompressible ExB drift, *van Rij & Hirshman (1989)*):

$$\left(v_{\parallel} \mathbf{b} + \frac{c}{\langle B^2 \rangle} \frac{d\Phi}{d\psi} \mathbf{B} \times \nabla \psi \right) \cdot \nabla f_1 - \frac{(1 - \xi^2)}{2B} v (\nabla_{\parallel} B) \frac{\partial f_1}{\partial \xi} - C\{f_1\} = -\mathbf{v}_m \cdot \nabla \psi \frac{\partial f_M}{\partial \psi}$$

$$\xi = v_{\parallel} / v$$

2. “Partial trajectories” (Correct ExB drift):

$$\left(v_{\parallel} \mathbf{b} + \frac{c}{B^2} \frac{d\Phi}{d\psi} \mathbf{B} \times \nabla \psi \right) \cdot \nabla f_1 - \frac{(1 - \xi^2)}{2B} v (\nabla_{\parallel} B) \frac{\partial f_1}{\partial \xi} - C\{f_1\} = -\mathbf{v}_m \cdot \nabla \psi \frac{\partial f_M}{\partial \psi}$$

3. “Full trajectories” (Including other terms required to conserve μ):

$$\left(v_{\parallel} \mathbf{b} + \frac{c}{B^2} \frac{d\Phi}{d\psi} \mathbf{B} \times \nabla \psi \right) \cdot \nabla f_1 + \left[-\frac{(1 - \xi^2)}{2B} v (\nabla_{\parallel} B) + \frac{c\xi(1 - \xi^2)}{2B^3} \frac{d\Phi}{d\psi} \mathbf{B} \times \nabla \psi \cdot \nabla B \right] \frac{\partial f_1}{\partial \xi} + \frac{cv}{2B^3} (1 + \xi^2) \frac{d\Phi}{d\psi} (\mathbf{B} \times \nabla \psi \cdot \nabla B) \frac{\partial f_1}{\partial v} - C\{f_1\} = -\mathbf{v}_m \cdot \nabla \psi \frac{\partial f_M}{\partial \psi}$$

Several choices are available for the E_r terms

1. “DKES trajectories” (Incompressible ExB drift, *van Rij & Hirshman (1989)*):

$$\left(v_{\parallel} \mathbf{b} + \frac{c}{\langle B^2 \rangle} \frac{d\Phi}{d\psi} \mathbf{B} \times \nabla \psi \right) \cdot \nabla f_1 - \frac{(1 - \xi^2)}{2B} v (\nabla_{\parallel} B) \frac{\partial f_1}{\partial \xi} - C\{f_1\} = -\mathbf{v}_m \cdot \nabla \psi \frac{\partial f_M}{\partial \psi}$$

$$\xi = v_{\parallel} / v$$

2. “Partial trajectories” (Correct ExB drift):

$$\left(v_{\parallel} \mathbf{b} + \frac{c}{B^2} \frac{d\Phi}{d\psi} \mathbf{B} \times \nabla \psi \right) \cdot \nabla f_1 - \frac{(1 - \xi^2)}{2B} v (\nabla_{\parallel} B) \frac{\partial f_1}{\partial \xi} - C\{f_1\} = -\mathbf{v}_m \cdot \nabla \psi \frac{\partial f_M}{\partial \psi}$$

3. “Full trajectories” (Including other terms required to conserve μ):

$$\begin{aligned} \left(v_{\parallel} \mathbf{b} + \frac{c}{B^2} \frac{d\Phi}{d\psi} \mathbf{B} \times \nabla \psi \right) \cdot \nabla f_1 + \left[-\frac{(1 - \xi^2)}{2B} v (\nabla_{\parallel} B) + \frac{c\xi(1 - \xi^2)}{2B^3} \frac{d\Phi}{d\psi} \mathbf{B} \times \nabla \psi \cdot \nabla B \right] \frac{\partial f_1}{\partial \xi} \\ + \frac{cv}{2B^3} (1 + \xi^2) \frac{d\Phi}{d\psi} (\mathbf{B} \times \nabla \psi \cdot \nabla B) \frac{\partial f_1}{\partial v} - C\{f_1\} = -\mathbf{v}_m \cdot \nabla \psi \frac{\partial f_M}{\partial \psi} \end{aligned}$$

These models are ordered from least to most accurate, in a sense, but care is required...

In the partial and full trajectory models, unphysical constraints will be imposed on f unless you are careful.

Example: partial trajectories:

$$\left(v_{\parallel} \mathbf{b} + \frac{c}{B^2} \frac{d\Phi}{d\psi} \mathbf{B} \times \nabla \psi \right) \cdot \nabla f_1 - \frac{(1 - \xi^2)}{2B} v (\nabla_{\parallel} B) \frac{\partial f_1}{\partial \xi} - C\{f_1\} = -\mathbf{v}_m \cdot \nabla \psi \frac{\partial f_M}{\partial \psi}$$



Consider the $\left\langle \int d^3v (...) \right\rangle$ moment:

$$\frac{d\Phi}{d\psi} \left\langle \int d^3v f_1 \mathbf{B} \times \nabla \psi \cdot \nabla \frac{1}{B^2} \right\rangle = 0$$

In the partial and full trajectory models, unphysical constraints will be imposed on f unless you are careful.

Example: partial trajectories:


$$\left(v_{\parallel} \mathbf{b} + \frac{c}{B^2} \frac{d\Phi}{d\psi} \mathbf{B} \times \nabla \psi \right) \cdot \nabla f_1 - \frac{(1 - \xi^2)}{2B} v (\nabla_{\parallel} B) \frac{\partial f_1}{\partial \xi} - C\{f_1\} = -\mathbf{v}_m \cdot \nabla \psi \frac{\partial f_M}{\partial \psi}$$



Consider the $\left\langle \int d^3v (...) \right\rangle$ moment:

$$\frac{d\Phi}{d\psi} \left\langle \int d^3v f_1 \mathbf{B} \times \nabla \psi \cdot \nabla \frac{1}{B^2} \right\rangle = 0$$

If $\frac{d\Phi}{d\psi} = 0$,



$0 = 0$. No problem.

In the partial and full trajectory models, unphysical constraints will be imposed on f unless you are careful.

Example: partial trajectories:

$$\left(v_{\parallel} \mathbf{b} + \frac{c}{B^2} \frac{d\Phi}{d\psi} \mathbf{B} \times \nabla \psi \right) \cdot \nabla f_1 - \frac{(1 - \xi^2)}{2B} v (\nabla_{\parallel} B) \frac{\partial f_1}{\partial \xi} - C\{f_1\} = -\mathbf{v}_m \cdot \nabla \psi \frac{\partial f_M}{\partial \psi}$$



Consider the $\left\langle \int d^3v (...) \right\rangle$ moment:

$$\frac{d\Phi}{d\psi} \left\langle \int d^3v f_1 \mathbf{B} \times \nabla \psi \cdot \nabla \frac{1}{B^2} \right\rangle = 0$$

If $\frac{d\Phi}{d\psi} = 0$,

$0 = 0$. No problem.

If $\frac{d\Phi}{d\psi} \neq 0$,

$$\left\langle \int d^3v f_1 \mathbf{B} \times \nabla \psi \cdot \nabla \frac{1}{B^2} \right\rangle = 0$$

Unphysical constraint on f_1 .

Solution for $\frac{d\Phi}{d\psi} = 0$ is very different from solution for $\frac{d\Phi}{d\psi} = \varepsilon$.

In the partial and full trajectory models, unphysical constraints will be imposed on f unless you are careful.

Example: partial trajectories:

$$\left(v_{\parallel} \mathbf{b} + \frac{c}{B^2} \frac{d\Phi}{d\psi} \mathbf{B} \times \nabla \psi \right) \cdot \nabla f_1 - \frac{(1 - \xi^2)}{2B} v (\nabla_{\parallel} B) \frac{\partial f_1}{\partial \xi} - C\{f_1\} = -\mathbf{v}_m \cdot \nabla \psi \frac{\partial f_M}{\partial \psi}$$



Consider the $\left\langle \int d^3 v (...) \right\rangle$ moment:

$$\frac{d\Phi}{d\psi} \left\langle \int d^3 v f_1 \mathbf{B} \times \nabla \psi \cdot \nabla \frac{1}{B^2} \right\rangle = 0$$

If $\frac{d\Phi}{d\psi} = 0$,

$0 = 0$. No problem.

If $\frac{d\Phi}{d\psi} \neq 0$,

$$\left\langle \int d^3 v f_1 \mathbf{B} \times \nabla \psi \cdot \nabla \frac{1}{B^2} \right\rangle = 0$$

Unphysical constraint on f_1 .

Solution for $\frac{d\Phi}{d\psi} = 0$ is very different from solution for $\frac{d\Phi}{d\psi} = \varepsilon$.

Similar problem for the $\left\langle \int d^3 v v^2 (...) \right\rangle$ moment, & for full trajectories.

The partial and full trajectory models become well-behaved if you include sources.

$$(\nu_{\parallel} \mathbf{b} + \mathbf{v}_E) \cdot \nabla f_1 + \dot{\xi} \frac{\partial f_1}{\partial \xi} - C_{\ell} \{f_1\} - S_p f_M \left(\frac{mv^2}{2T} - \frac{5}{2} \right) - S_h f_M \left(\frac{mv^2}{2T} - \frac{3}{2} \right) = -\mathbf{v}_d \cdot \nabla f_M$$

The partial and full trajectory models become well-behaved if you include sources.

$$(\nu_{\parallel} \mathbf{b} + \mathbf{v}_E) \cdot \nabla f_1 + \xi \frac{\partial f_1}{\partial \xi} - C_{\ell} \{f_1\} - S_p f_M \left(\frac{mv^2}{2T} - \frac{5}{2} \right) - S_h f_M \left(\frac{mv^2}{2T} - \frac{3}{2} \right) = -\mathbf{v}_d \cdot \nabla f_M$$

2 extra unknowns (S_p and S_h) require 2 extra equations.

$$\begin{array}{l} \text{Kinetic equation } \{ \\ \langle \int d^3v f_1 \rangle_{\psi} = 0 \{ \\ \langle \int d^3v f_1 v^2 \rangle_{\psi} = 0 \{ \end{array} \left(\begin{array}{ccc} \text{---} & \text{---} & \text{---} \\ \text{---} & 0 & 0 \\ \text{---} & 0 & 0 \end{array} \right) \underbrace{\begin{pmatrix} f_1 \\ S_p \\ S_h \end{pmatrix}}_{\text{Vector of unknowns}} = \begin{pmatrix} -\mathbf{v}_d \cdot \nabla f_M \\ 0 \\ 0 \end{pmatrix}$$

The partial and full trajectory models become well-behaved if you include sources.

$$(\nu_{\parallel} \mathbf{b} + \mathbf{v}_E) \cdot \nabla f_1 + \xi \frac{\partial f_1}{\partial \xi} - C_{\ell} \{f_1\} - S_p f_M \left(\frac{mv^2}{2T} - \frac{5}{2} \right) - S_h f_M \left(\frac{mv^2}{2T} - \frac{3}{2} \right) = -\mathbf{v}_d \cdot \nabla f_M$$

2 extra unknowns (S_p and S_h) require 2 extra equations.

$$\begin{array}{l} \text{Kinetic equation} \{ \\ \langle \int d^3v f_1 \rangle_{\psi} = 0 \{ \\ \langle \int d^3v f_1 v^2 \rangle_{\psi} = 0 \{ \end{array} \left(\begin{array}{ccc} \text{---} & \text{---} & \text{---} \\ \text{---} & 0 & 0 \\ \text{---} & 0 & 0 \end{array} \right) \underbrace{\begin{pmatrix} f_1 \\ S_p \\ S_h \end{pmatrix}}_{\text{Vector of unknowns}} = \begin{pmatrix} -\mathbf{v}_d \cdot \nabla f_M \\ 0 \\ 0 \end{pmatrix}$$


DKES trajectories: $S_p = 0$, $S_h = 0$.

Partial trajectories: $S_p \neq 0$, $S_h \neq 0$.

Full trajectories: $S_p = 0$, $S_h \neq 0$ except at the ambipolar E_r .



Example: W7-X edge.

Let's revisit the W7-X scenarios from
Turkin et al, *PoP* **18**, 022505 (2011):



Example: W7-X edge. Trajectory model has little impact on ambipolar E_r , modest effect on j_{bs}

— DKES trajectories
— Partial trajectories
— Full trajectories




Example: W7-X edge. Trajectory model has little impact on ambipolar E_r , modest effect on j_{bs}

— DKES trajectories
— Partial trajectories
— Full trajectories

----- -33 kV/m


$$E_r^{res} = B v_{th,i} tr / R = 100 \text{ kV/m}$$



For the next few slides, we will consider the ion transport matrix L_{ik}


$$\begin{pmatrix} \langle \Gamma \cdot \nabla \psi \rangle \\ \langle \mathbf{q} \cdot \nabla \psi \rangle \\ \langle V_{\parallel} B \rangle \end{pmatrix} \propto \underbrace{\begin{pmatrix} L_{11} & L_{12} & L_{13} \\ L_{21} & L_{22} & L_{23} \\ L_{31} & L_{32} & L_{33} \end{pmatrix}}_{\text{Transport matrix}} \begin{pmatrix} \frac{1}{n} \frac{dn}{d\psi} + \frac{e}{T} \frac{d\Phi}{d\psi} - \frac{3}{2T} \frac{dT}{d\psi} \\ \frac{1}{T} \frac{dT}{d\psi} \\ \langle E_{\parallel} B \rangle \end{pmatrix}$$

When $E_* = E_r / E_r^{\text{res}}$ is < 0.3 , the 3 models are nearly identical; otherwise differences can be significant.



$$\nu_{ii} R / \nu_{th,i} = 0.01, \quad E_r^{\text{res}} = B \nu_{th,i} r / R$$







The same pattern is evident for LHD



$$\nu_{ii}R / \nu_{th,i} = 0.01, \quad E_r^{res} = B\nu_{th,i}tr / R$$

SFINCS allows comparison between collision operators

- SFINCS: Fokker–Planck
- +— SFINCS: Pure pitch–angle scattering
- SFINCS: Momentum–conserving model
- + DKES: Pure pitch–angle scattering









$$\nu' = \nu_{ii} R / \nu_{th,i},$$

W7-X geometry,

$E_r = 0$ for these plots. 28

Similar patterns are apparent for LHD geometry

- SFINCS: Fokker–Planck
- SFINCS: Pure pitch–angle scattering
- ▲— SFINCS: Momentum–conserving model



$$v' = v_{ii} R / v_{th,i},$$

$E_r = 0$ for these plots.

Summary

- Our new code SFINCS allows a comparison of several variants of the drift-kinetic equation, differing in the E_r terms.
 - Below $\sim 1/3$ of the E_r resonance, the variants give nearly identical results.
 - For larger E_r/E_r^{res} , there are substantial differences, especially in the flows and j_{bs} .
 - The “full trajectory” model is probably the best of the 3 models considered here, but radial coupling could also be important.
- Momentum conservation in collisions is always important for parallel flow and current. The full linearized Fokker-Planck operator gives results quite close to a momentum-conserving model.

Extra slides

For the next few slides, we will consider the ion transport matrix L_{ik}

$$\begin{pmatrix} \frac{Ze(G + \iota I)}{ncTG} \left\langle \int d^3v f \mathbf{v}_d \cdot \nabla \psi \right\rangle \\ \frac{Ze(G + \iota I)}{ncTG} \left\langle \int d^3v f \frac{m v^2}{2T} \mathbf{v}_d \cdot \nabla \psi \right\rangle \\ \frac{1}{v_{th} B_0} \langle B V_{\parallel} \rangle \end{pmatrix} = \begin{pmatrix} L_{1,1} & L_{1,2} & L_{1,3} \\ L_{2,1} & L_{2,2} & L_{2,3} \\ L_{3,1} & L_{3,2} & L_{3,3} \end{pmatrix} \begin{pmatrix} \frac{GTc}{ZeB_0 v_{th}} \left[\frac{1}{n} \frac{dn}{d\psi} + \frac{Ze}{T} \frac{d\Phi}{d\psi} - \frac{3}{2} \frac{1}{T} \frac{dT}{d\psi} \right] \\ \frac{GTc}{ZeB_0 v_{th}} \frac{1}{T} \frac{dT}{d\psi} \\ \frac{Ze}{T} (G + \iota I) \frac{\langle E_{\parallel} B \rangle}{\langle B^2 \rangle} \end{pmatrix}$$

$$\mathbf{B} = \beta \nabla \psi + G(\psi) \nabla \zeta + I(\psi) \nabla \theta$$

New speed discretization is highly efficient

Spectral collocation method based on non-standard orthogonal polynomials in v , not v^2 :

$$\int_0^\infty dx P_i(x) P_j(x) e^{-x^2} \propto \delta_{ij}$$


Laguerre/Sonine polynomials lose accuracy because of nonanalytic $\sqrt{\cdot}$ in Jacobian at $x = 0$.

New speed discretization is highly efficient

Spectral collocation method based on non-standard orthogonal polynomials in v , not v^2 :

$$\int_0^\infty dx P_i(x) P_j(x) e^{-x^2} \propto \delta_{ij}$$

Laguerre/Sonine polynomials lose accuracy because of nonanalytic $\sqrt{\cdot}$ in Jacobian at $x = 0$.



SFINCS can use the full linearized Fokker-Planck collision operator.


$$C_i \{f_1\} = \underbrace{\left(\text{pitch-angle \& energy scattering} \right)}_{\text{test particle part}} + \underbrace{v_{ii} 3e^{-v^2/v_{th,i}^2} \left[f_1 - \frac{H}{2\pi v_{th,i}^2} + \frac{v^2}{2\pi v_{th,i}^4} \frac{\partial^2 G}{\partial v^2} \right]}_{\text{field particle part}}$$

$$\nabla_{\mathbf{v}}^2 H + 4\pi f_1 = 0$$


$$\nabla_{\mathbf{v}}^2 G - 2H = 0$$

SFINCS can use the full linearized Fokker-Planck collision operator.

$$C_i \{f_1\} = \underbrace{\left(\text{pitch-angle \& energy scattering} \right)}_{\text{test particle part}} + \underbrace{v_{ii} 3e^{-v^2/v_{th,i}^2} \left[f_1 - \frac{H}{2\pi v_{th,i}^2} + \frac{v^2}{2\pi v_{th,i}^4} \frac{\partial^2 G}{\partial v^2} \right]}_{\text{field particle part}}$$




Computed radial neoclassical diffusivity in the LHD stellarator



$$M = \frac{\nu_{th} E_r}{B}$$



Beidler et al,
Nucl Fusion
(2011)

- Unlike in tokamaks, collisionless trapped particle orbits are not generally confined.





- It's important to retain $\mathbf{v}_{\mathbf{E} \times \mathbf{B}}$ in $d\mathbf{r}/dt$; otherwise you lose poloidal precession & collisionless detrapping. A little bit of E_r ($v_E \ll v_{th}$) makes a big difference.


Helically trapped orbit



Toroidally trapped orbit



Comparison of particle trajectories and collision operators for collisional transport in nonaxisymmetric plasmas



Matt Landreman, (University of Maryland, MIT, ORISE)

Håkan Smith, Per Helander (IPP Greifswald)

arXiv: 1312.6058 (2013)



Max-Planck-Institut
für Plasmaphysik



OAK RIDGE INSTITUTE FOR
SCIENCE AND EDUCATION
Managed by ORAU for DOE

