THRIVEHEALTH360 · ENHANCED EDITION

7-DAY BLOOD SUGAR RESET

A Practical 7-Day Plan for Steadier Energy and Balanced Eating

IMPORTANT MEDICAL DISCLAIMER

All content on ThriveHealth360 is for general educational and informational purposes only and is not a substitute for professional medical advice, diagnosis, or treatment. Supplements, foods, and lifestyle interventions discussed here have not been evaluated by the FDA, MHRA, TGA, or Health Canada and are not intended to diagnose, treat, cure, or prevent any disease. Always consult your doctor or a qualified healthcare provider before making changes to your diet, exercise routine, medications, or supplement regimen — especially if you have diabetes or prediabetes or take any prescription medication. Individual results vary and are not guaranteed. Health regulations differ by country, so content may not apply in your jurisdiction. Never disregard professional medical advice or delay seeking it because of something you read in this guide.

3	7	10	8
Core Pillars	Day Reset	PubMed Refs	Improvements

Prevalence: According to the CDC, tens of millions of US adults have prediabetes, and most people with prediabetes do not know they have it. Source: CDC National Diabetes Statistics Report, 2024.

Welcome & How to Use This Guide

If you're reading this, you probably know what it feels like to hit that 2 PM energy crash, feel hungry after a "healthy" meal, or feel trapped on a blood sugar rollercoaster. You're not alone.

> **This Is a Starting Point, Not a Permanent Diet**
>
> The 7-day reset is designed to help you see how food, movement and sleep affect how you feel. Long-term success requires building sustainable habits over weeks and months. Think of Day 7 not as a finish line but as the beginning of a new metabolic baseline. The 30-Day Path Forward section provides continuation guidance.

What This Guide Includes

- Strategic Eating — Food choices that keep you stable and satisfied
- Optional Supplement Information — what the research shows, and its limits
- Lifestyle Habits — Simple daily practices that amplify everything

How to Get Maximum Results

- Read the entire guide first — Understanding the why makes the how actually stick.
- Follow the daily protocols — Each day builds on the previous one.
- Track how you feel — Energy, cravings, digestion, sleep.
- Start immediately — The best time was yesterday. The second-best is today.

> **⚠ ⚕ If You Take Diabetes Medication — Read This First**
>
> If you are currently taking metformin, insulin, sulfonylureas, or any other glucose-lowering medication: monitor your blood sugar closely throughout this protocol. Berberine may potentiate the glucose-lowering effect of your medication, which can cause hypoglycemia (low blood sugar). Discuss all supplement additions with your prescribing physician before beginning. Do NOT stop or reduce your prescribed medication without medical supervision.
>
> Do not use the supplements in this guide if you are pregnant, planning a pregnancy or breastfeeding, and do not give them to children. Berberine, alpha-lipoic acid, chromium, magnesium, inositol and cinnamon can interact with other medicines (including blood thinners, blood pressure medicines and immunosuppressants) and may not suit people with kidney or liver disease; magnesium supplements in particular need medical advice if you have kidney problems. Check with your doctor or pharmacist first.

The Blood Sugar Problem

And Why You Can't Just "Eat Less"

When blood sugar spikes, your body releases insulin. When insulin is constantly elevated, two things happen: your cells stop listening to insulin signals (insulin resistance), and your body stores excess energy as fat. Willpower is only part of the picture — **hormones** and food composition matter too.

For many people, cutting calories alone increases hunger and cravings. The solution isn't restriction — it's **stabilization**.

The Three Root Causes of Blood Sugar Chaos

1. Poor Meal Timing & Composition

Eating carbs alone causes rapid glucose spikes. Pairing them with protein, fat, and fiber keeps you stable.

2. Nutrient Gaps and Optional Support

Many people do not get enough magnesium from food. Some people also take supplements such as berberine, alpha-lipoic acid or chromium, but these are not essential nutrients and the evidence for them is mixed (see the references).

3. Lifestyle Patterns That Sabotage You

Chronic stress can raise cortisol (which can raise blood sugar), poor sleep disrupts hunger hormones, and sedentary behavior reduces muscle glucose uptake.

> **The Good News**
>
> These three areas can be changed. Some people notice differences in energy or appetite within a week, but results vary and blood sugar markers usually take longer to move. This guide draws on published research, including randomized trials, and says so where the evidence is limited or mixed.

The 3-Pillar System for Blood Sugar Success

Pillar 1: Strategic Eating (The Foundation)

Every meal should contain:

- Protein (25–35g) — Slows glucose absorption
- Healthy Fat (10–15g) — Helps with fullness and slows digestion
- Fiber (7–10g) — Buffers carbohydrate impact
- Carbs (30–50g) — Choose low-glycemic options only

✅ Foods to Prioritize

• Proteins: Salmon, chicken, eggs, Greek yogurt
• Fats: Avocado, olive oil, nuts, seeds
• Carbs: Sweet potato, oats, quinoa, berries
• Fiber: Leafy greens, broccoli, chia seeds

❌ Foods to Minimize

• Refined carbs (white bread, pasta, cereal)
• Sugar (soda, juice, candy, desserts)
• Deep-fried and heavily processed snack foods
• Ultra-processed foods, fast food

Pillar 2: Metabolic Support (The Accelerator)

Berberine: Start Gradually to Minimize GI Side Effects

Do NOT start at 1,500mg/day on Day 1.

• Days 1–2: 500mg once daily with your largest meal
• Days 3–4: 500mg twice daily if Day 2 was tolerated well
• Day 5+: Full 500mg three times daily protocol

Always take berberine WITH food. This gradual titration reduces GI side effects (nausea, cramping, diarrhea) that cause some people to quit.

Timing	Supplement	Dose	Purpose
Morning	**Berberine HCl**	**500mg (build to 3×/day)**	Studied for glucose lowering; can interact with medicines
Morning	**Alpha-Lipoic Acid**	**300–600mg**	Evidence mixed; studied mainly in diabetic neuropathy
Lunch	**Chromium Picolinate**	**200mcg**	Optional; evidence is weak outside diabetes
Dinner (30 min before bed)	**Magnesium Glycinate**	**up to 350mg elemental**	Low intake is linked to higher diabetes risk; modest effects in trials
Optional Daily	**Inositol**	**2–4g**	Limited evidence
Optional Daily	**Cinnamon Extract**	**500–1,000mg**	Evidence mixed

Pillar 3: Lifestyle Habits (The Multiplier)

Post-Meal Walks (15 min)

Working muscles take up glucose partly without insulin; short activity breaks lowered post-meal glucose in a small study of people with type 2 diabetes [Ref 9]

Stress Management (10 min)

Meditation or breathwork can help you manage stress, which can affect blood sugar

Hydration

Drink water regularly through the day; needs vary, so ask your doctor if you have heart or kidney conditions

Sleep (7–9 hrs)

Regular, sufficient sleep supports insulin sensitivity and appetite regulation

Movement (30 min/day)

Include resistance training — it builds muscle, a major site of glucose uptake

Morning Sunlight (10 min)

Supports a regular daily rhythm

DAY 1

Foundation Setting

Goal: Establish stable meal patterns and begin metabolic support

Morning Protocol

- A glass of water upon waking
- Wait 30 minutes before eating
- Breakfast: 3 scrambled eggs + 1 slice whole grain toast + ½ avocado + 1 cup spinach sautéed in olive oil

✦ **Take: Berberine 500mg (Day 1 — start here, once only) + Alpha-Lipoic Acid 300mg**

Mid-Morning

- Green tea (no sugar)
- Handful of almonds if hungry

Lunch (1 PM)

- Grilled chicken breast (6oz) + 1 medium sweet potato + roasted broccoli with olive oil

✦ **Take: Chromium Picolinate 200mcg**

Post-Lunch

- 15-minute walk — no phone, focus on breathing

Afternoon Snack (4 PM)

- Greek yogurt (6oz) + berries + 1 tbsp almond butter
- Herbal tea

Dinner (7 PM)

- Baked salmon (6oz) + quinoa (½ cup) + asparagus with olive oil
- Large mixed green salad with olive oil dressing

✦ **Take: Magnesium glycinate (up to 350 mg elemental magnesium)**

Evening

- No screens 30 min before bed
- 10 min deep breathing or meditation
- Set a consistent bedtime and keep it

Track Today

- Energy level (1–10) at wake, mid-afternoon, evening
- Sugar cravings: present? (yes/no)
- Any GI discomfort from Berberine? (note for tomorrow)

DAY 2

Strategic Eating

Goal: Master meal composition and recognize hunger vs. cravings

Morning Protocol

- A glass of water upon waking
- Breakfast: Veggie omelet (3 eggs, peppers, mushrooms, onions) + 2 slices whole grain toast + ½ banana

✦ **Take: Berberine 500mg (once or twice depending on Day 1 tolerance) + ALA 300mg**

Mid-Morning

- Coffee or tea (black, no sugar) + handful of walnuts

Lunch (1 PM)

- Turkey sandwich on whole grain bread with lettuce, tomato, avocado, mustard + side salad

✦ **Take: Chromium Picolinate 200mcg**

Post-Lunch

- 15-minute walk

Afternoon Snack (4 PM)

- Apple + 2 tbsp almond butter
- Water with lemon

Dinner (7 PM)

- Grass-fed beef (6oz) or turkey + sweet potato + roasted Brussels sprouts
- Side of sauerkraut or kimchi (natural probiotics)

✦ **Take: Magnesium glycinate (up to 350 mg elemental magnesium)**

Key Lesson

- True hunger = slight stomach rumble, gradual onset
- Cravings = immediate intense desire, emotionally triggered
- Notice which one is driving your food choices today

DAY 3

Metabolic Boost

Goal: Increase metabolic support and deepen food quality awareness

Morning Protocol

- Water + 10 minutes of morning daylight (supports a regular sleep–wake rhythm)
- Breakfast: Oatmeal (½ cup old-fashioned) with cinnamon, berries, walnuts, almond milk

✦ **Take: Berberine 500mg twice daily if Day 2 was well-tolerated + ALA 300mg + Inositol 2g (optional)**

Mid-Morning

- Green smoothie: spinach, frozen berries, protein powder, almond butter, almond milk
- Add 1 tbsp ground flaxseed for extra soluble fiber

Lunch (1 PM)

- Tuna salad (3oz in olive oil) on mixed greens + chickpeas + olive oil dressing

✦ **Take: Chromium Picolinate 200mcg**

Post-Lunch

- 20-minute walk — slightly longer to boost metabolism

Afternoon Snack (4 PM)

- Hummus + vegetable sticks (carrots, celery, peppers)
- Herbal tea

Dinner (7 PM)

- Baked white fish (6oz) + wild rice + roasted vegetables (zucchini, bell pepper, cauliflower)

✦ **Take: Magnesium glycinate (up to 350 mg elemental magnesium)**

Key Lesson

- Whole foods produce a different energy signature than processed foods
- Notice the absence of a crash or bloat after today's meals

DAY 4

Hormonal Rebalancing

Goal: Deepen insulin sensitivity and hormonal optimization

Morning Protocol

- Water + 5 min light stretching or yoga
- Breakfast: Greek yogurt (6oz) + granola (2 tbsp) + berries + honey (1 tsp) + walnuts

✦ **Take: Berberine 500mg + ALA 300mg**

Mid-Morning

- Black coffee + handful of almonds + 2–3 squares dark chocolate (70%+ cacao)

Lunch (1 PM)

- Chicken stir-fry: chicken + broccoli + bell peppers + snap peas in olive oil and garlic + brown rice (½ cup)

✦ **Take: Chromium Picolinate 200mcg**

Post-Lunch

- 15-minute walk

Afternoon Snack (4 PM)

- Cottage cheese (½ cup) + berries + cinnamon
- Green tea

Dinner (7 PM)

- Lamb or beef (6oz) + roasted sweet potato + leafy greens sautéed in garlic and olive oil
- Side of fermented vegetables

✦ **Take: Magnesium glycinate (up to 350 mg elemental magnesium) + Inositol 2g (optional)**

Key Lesson

- Habits are settling in. By Day 4, some people notice deeper sleep and fewer cravings, but experiences vary.
- Keep notes on how you feel — they are useful to share with your doctor.

DAY 5

Sustained Energy

Goal: Experience sustained energy without crashes

Morning Protocol

- Water + 10 min morning sunlight
- Breakfast: Smoked salmon on whole grain toast with cream cheese, cucumber, capers + fresh fruit

✦ **Take: Berberine 500mg — advance to full 3x/day if well-tolerated + ALA 300mg**

Mid-Morning

- Matcha green tea with almond milk (sustained caffeine + L-theanine = calm focus)

Lunch (1 PM)

- Beef or chicken + quinoa + roasted root vegetables (parsnips, carrots, beets) + olive oil

✦ **Take: Chromium Picolinate 200mcg**

Post-Lunch

- 20-minute walk + 5 min breathwork

Afternoon Snack (4 PM)

- Mixed nuts and seeds (1oz) + small orange or apple
- Water with lemon and ginger

Dinner (7 PM)

- Baked cod or halibut (6oz) + sweet potato (baked) + roasted cauliflower + mixed green salad

✦ **Take: Magnesium glycinate (up to 350 mg elemental magnesium)**

Key Lesson

- Check: Do you have the usual 2–3 PM energy crash today?
- Some people find their afternoon energy steadier by Day 5; others do not yet. Both are normal.
- Keep tracking — patterns take longer than a week to show.

DAY 6

Stabilization

Goal: Lock in new patterns and prepare for long-term success

Morning Protocol

- Water + 10 min stretching or yoga
- Breakfast: Veggie frittata (6 eggs, spinach, tomatoes, peppers, feta) + whole grain toast

✦ **Take: Berberine 500mg + ALA 300mg**

Mid-Morning

- Green juice: celery, cucumber, ginger, apple, lemon
- Brazil nuts — 1–2 (a rich source of selenium; do not eat large amounts regularly)

Lunch (1 PM)

- Turkey meatballs + zucchini noodles + quality tomato sauce + parmesan + side salad

✦ **Take: Chromium Picolinate 200mcg**

Post-Lunch

- 20-minute walk (outdoors if possible)

Afternoon Snack (4 PM)

- Nut butter with apple slices
- Herbal tea

Dinner (7 PM)

- Lamb steak or grass-fed beef (6oz) + roasted root vegetables + leafy greens
- Small amount of fermented food (sauerkraut or kimchi)

✦ **Take: Magnesium glycinate (up to 350 mg elemental magnesium) + Inositol 2g (optional)**

Evening Ritual

- 15 min meditation or journaling
- Consistent bedtime
- No screens 30 min prior

Key Lesson

- You've done this 6 days. Notice: these patterns now feel normal, not effortful.
- Your body is craving the right foods — that's biology, not willpower.

DAY 7

Measurement & Next Steps

Goal: Assess progress and create your sustainable forward plan

Morning Protocol

- Water + 10 min walk or yoga
- Breakfast: Your favorite balanced meal from the past 6 days

✦ **Take: Berberine 500mg + ALA 300mg**

Lunch (1 PM)

- Your favorite balanced lunch from the past 6 days

✦ **Take: Chromium Picolinate 200mcg**

Post-Lunch

- 20-minute walk + reflection time

Afternoon — Take Your Measurements

- Energy level (1–10): Did it improve vs. Day 1?
- Cravings: Gone? Significantly reduced?
- Sleep quality: Deeper, more restorative?
- Digestion: More stable, less bloating?
- Weight: changes vary from person to person; track it only if it helps you
- Mood and mental clarity: Sharper, more stable?
- Joint comfort and mental clarity: any change?

Dinner (7 PM)

- Your favorite balanced dinner from the past 6 days

✦ **Take: Magnesium glycinate (up to 350 mg elemental magnesium)**

Evening

- Write down your results and which meals you enjoyed most
- Plan which supplements you'll continue
- Read the 30-Day Path Forward section

Choosing Supplements (If You Decide To)

Supplement quality varies. If you decide to use any of these, look for products tested by an independent third party (such as USP, NSF or ConsumerLab) and check the label with your pharmacist.

Transparency Note

If a link appears below, it may be an affiliate link: we may earn a commission at no extra cost to you. We do not recommend specific brands in this edition, and nothing here replaces advice from your doctor or pharmacist.

Berberine HCL (500mg)

Berberine HCl is the form used in most studies. Not suitable during pregnancy or breastfeeding, and it can interact with medicines.

Alpha-Lipoic Acid (300mg)

Both R-ALA and the standard mixed form are sold. Evidence of benefit is mixed and mostly in diabetic neuropathy.

Chromium Picolinate (200mcg)

Picolinate is the form used in many trials. Evidence of benefit is limited outside diabetes.

Magnesium Glycinate (check elemental Mg on the label)

Glycinate is generally gentle on the stomach. No more than 350 mg/day of elemental magnesium from supplements, and ask your doctor first if you have kidney problems.

Your 30-Day Path Forward

You have finished the 7 days. How you feel, and what any blood tests show, will depend on your starting point and how consistently you keep the habits going. The next 30 days are about making the helpful ones routine.

Weeks 2–4: Lock It In

- Continue all supplements at current dosages
- Maintain balanced meal composition at every meal
- Keep daily movement and post-meal walks
- Protect your sleep schedule (consistent bedtime)
- Gradually reintroduce foods you missed — in balanced format

What to Track Over 30 Days

Energy
Is it steadier through the day?

Cravings
Easier or harder to manage than before?

Weight
Track only if it helps; changes vary widely

Mental Clarity
Any change in focus?

Sleep
Falling asleep and staying asleep

Mood
Any change you notice?

Long-Term Supplement Options

> **Option 1: Maintenance**
> Continue only the supplements you and your doctor decide are worthwhile. ~$50–80/month.

> **Option 2: Cycling**
> 3 months on, 1 month off. A common pattern, though evidence that cycling is needed is limited. ~$40–60/month.

> **Option 3: Core Only**
> Berberine + magnesium only, if you and your doctor decide to continue. ~$30–40/month.

Food and Lifestyle Come First

Meals, movement, sleep and stress habits are the foundation. Supplements are optional extras with modest and variable effects, and they cannot replace that foundation.

When to Seek Medical Attention

⚕ Stop the Protocol and Seek Immediate Medical Care If You Experience:

• Hypoglycemia symptoms: shaking, sweating, dizziness, confusion, extreme weakness, rapid heartbeat, or loss of consciousness. Call emergency services immediately.
• Severe gastrointestinal distress: prolonged vomiting, severe abdominal cramping, bloody stool, or diarrhea lasting more than 48 hours.
• Chest pain, shortness of breath, or severe headache — these are never normal responses to dietary change.
• Altered mental status, extreme fatigue, or unusual thirst with frequent urination (potential signs of DKA or severe hyperglycemia).
• Any allergic reaction symptoms: rash, swelling, difficulty breathing.

Managing Hypoglycemia on This Protocol

Berberine can lower blood glucose, especially combined with diabetes medications. If you experience mild hypoglycemia (shakiness, lightheadedness, sweating):

- Consume 15g of fast-acting carbohydrates immediately (4 oz juice, 3–4 glucose tablets, or 1 tbsp honey)
- Wait 15 minutes and recheck blood glucose if you have a glucometer
- If symptoms persist or worsen, seek emergency medical care
- Report the episode to your physician and discuss adjusting supplement or medication doses

Managing Berberine GI Side Effects

Nausea, bloating, and loose stools are the most common side effects and are usually dose-dependent.

- Always take berberine WITH food — never on an empty stomach
- Follow the gradual titration schedule in Pillar 2
- If GI distress persists beyond Day 3 at low dose, reduce to 250mg and consult your doctor
- Splitting the dose across meals often resolves symptoms without reducing efficacy

Reference List

Below are the 10 key sources behind this guide, each with a note on what it does and does not show. Evidence for several of the supplements is mixed or limited. References 1–9 were rechecked against PubMed, PMC or the publisher in October 2026; reference 10 links to the CDC page. Please read the original papers before relying on any single finding.

1. Yin J, Xing H, Ye J. (2008). Efficacy of Berberine in Patients with Type 2 Diabetes Mellitus. *Metabolism*. DOI: 10.1016/j.metabol.2008.01.013

PubMed · PMID 18442638

Small pilot study (36 people in the main comparison, plus a 48-person add-on study): berberine lowered blood glucose and HbA1c over three months and performed similarly to metformin in the head-to-head comparison. It was a pilot, and it did not include a rosiglitazone arm.

2. Xie W, Su F, Wang G, et al. (2022). Glucose-Lowering Effect of Berberine on Type 2 Diabetes: A Systematic Review and Meta-Analysis. *Frontiers in Pharmacology.* DOI: 10.3389/fphar.2022.1015045

PubMed · PMID PMC9709280

Meta-analysis of 37 randomized trials (3,048 people with type 2 diabetes): berberine lowered fasting glucose, HbA1c and post-meal glucose compared with control. This is evidence in people who already have diabetes, not prediabetes.

3. Anderson RA, Cheng N, Bryden NA, et al. (1997). Elevated Intakes of Supplemental Chromium Improve Glucose and Insulin Variables in Individuals with Type 2 Diabetes. *Diabetes.* DOI: 10.2337/diab.46.11.1786

PubMed · PMID 9356027

Placebo-controlled trial in 180 people with type 2 diabetes in China: chromium picolinate improved HbA1c and fasting glucose, with the larger effect at 1,000 mcg/day than at 200 mcg/day. Later reviews found the overall evidence mixed (see reference 4).

4. Balk EM, et al. (2007). Effect of Chromium Supplementation on Glucose Metabolism and Lipids: A Systematic Review of Randomized Controlled Trials. *Diabetes Care*.

PubMed · PMID 17519436

Systematic review: in people with type 2 diabetes, chromium lowered HbA1c by about 0.6 percentage points and fasting glucose by about 1.0 mmol/L, but no benefit was found in people without diabetes, and the authors noted limited study quality. Chromium is therefore optional and its evidence for prediabetes is weak.

5. Simental-Mendía LE, Sahebkar A, Rodríguez-Morán M, Guerrero-Romero F. (2016). A Systematic Review and Meta-Analysis of Randomized Controlled Trials on the Effects of Magnesium Supplementation on Insulin Sensitivity and Glucose Control. *Pharmacological Research.* DOI: 10.1016/j.phrs.2016.06.019

View Source

Meta-analysis of randomized trials: magnesium taken for four months or longer improved fasting glucose and insulin resistance (HOMA-IR) in people with and without diabetes; shorter trials did not show the same effect. A meta-analysis cannot show how magnesium works, only whether outcomes changed.

6. Xu L, Li X, Wang X, Xu M. (2022). Effects of Magnesium Supplementation on Hyperglycemia, Hypercholesterolemia, and Hypertension in Type 2 Diabetes. *Frontiers in Nutrition.* DOI: 10.3389/fnut.2022.1020327

PubMed · PMID PMC9889557

Analysis of 24 randomized trials in people with type 2 diabetes: effects on glucose, lipids and blood pressure depended on dose, duration and baseline magnesium status, and the authors describe magnesium as an add-on, not a replacement for treatment. The findings do not apply directly to prediabetes.

7. Baicus C, Purcarea A, von Elm E, Delcea C, Furtunescu FL. (2024). Alpha-Lipoic Acid for Diabetic Peripheral Neuropathy. *Cochrane Database of Systematic Reviews.* DOI: 10.1002/14651858.CD012967.pub2

PubMed · PMID 38205823

Cochrane review of oral alpha-lipoic acid for people who already have diabetic neuropathy: it probably has little or no effect on neuropathy symptoms at six months (moderate-certainty evidence) and may have little or no effect on nerve impairment (low-certainty). Doses studied were 600–1,800 mg/day. This is not evidence for prediabetes.

8. Vallianou N, Evangelopoulos A, Koutalas P. (2009). Alpha-Lipoic Acid and Diabetic Neuropathy. *Reviews in Diabetic Studies.* DOI: 10.1900/RDS.2009.6.230

PubMed · PMID 20043035

Review of alpha-lipoic acid in diabetic neuropathy. It is a neuropathy review, not a trial of blood sugar control; a 2019 meta-analysis of 10 trials found no difference from placebo in HbA1c or fasting glucose in people with uncomplicated type 2 diabetes (Ebada MA, et al., Iran J Pharm Res).

9. Dempsey PC, et al. (2016). Benefits for Type 2 Diabetes of Interrupting Prolonged Sitting With Brief Bouts of Light Walking or Simple Resistance Activities. *Diabetes Care.* DOI: 10.2337/dc15-2336

PubMed · PMID 27208318

Randomized crossover study in adults with type 2 diabetes: breaking up prolonged sitting with 3-minute bouts of light walking or simple resistance activities every 30 minutes lowered post-meal glucose and insulin responses compared with uninterrupted sitting. It did not test a single 15-minute post-meal walk.

10. Centers for Disease Control and Prevention. (2024). National Diabetes Statistics Report, 2024. *U.S. Department of Health and Human Services.*

View Source

General source for the statement that tens of millions of US adults have prediabetes and most do not know it. See the CDC page for current figures: cdc.gov/diabetes/data