





Machine Learning

Linear Regression with multiple variables

Multiple features

Multiple features (variables).

Size (feet ²)	Price (\$1000)
 x	y 
2104	460
1416	232
1534	315
852	178
...	...

$$\underline{h_{\theta}(x) = \theta_0 + \theta_1 x}$$

Multiple features (variables).

Size (feet ²) x_1	Number of bedrooms x_2	Number of floors x_3	Age of home (years) x_4	Price (\$1000) y
2104	5	1	45	460
1416	3	2	40	232
1534	3	2	30	315
852	2	1	36	178
...

Notation:

- n = number of features
- $x^{(i)}$ = input (features) of i^{th} training example.
- $x_j^{(i)}$ = value of feature j in i^{th} training example.

$n = 4$

$m = 47$

$$\underline{x^{(2)}} = \begin{bmatrix} 1416 \\ 3 \\ 2 \\ 40 \end{bmatrix}$$

$$x_3^{(2)} = 2$$

Hypothesis:

Previously: $h_{\theta}(x) = \theta_0 + \theta_1 x$

$$h_{\theta}(x) = \theta_0 + \theta_1 x_1 + \theta_2 x_2 + \theta_3 x_3 + \theta_4 x_4$$

e.g. $\underline{h_0(x)} = \underline{80} + \underline{0.1x_1} + \underline{0.01x_2} + 3x_3 - 2x_4$
↑ ↑ ↑
age

$$\rightarrow h_{\theta}(x) = \theta_0 + \theta_1 x_1 + \theta_2 x_2 + \dots + \theta_n x_n$$

For convenience of notation, define $x_0 = 1$. ($x_0^{(i)} = 1$)

$$x = \begin{bmatrix} x_0 \\ x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \in \mathbb{R}^{n+1}$$

$$\theta = \begin{bmatrix} \theta_0 \\ \theta_1 \\ \theta_2 \\ \vdots \\ \theta_n \end{bmatrix} \in \mathbb{R}^{n+1}$$

$$h_{\theta}(x) = \theta_0 x_0 + \theta_1 x_1 + \dots + \theta_n x_n$$

$$= \theta^T x$$

Multivariate refers to
multiple
features/variables
working on

$$\underbrace{[\theta_0 \ \theta_1 \ \dots \ \theta_n]}_{\theta^T} \begin{bmatrix} x_0 \\ x_1 \\ \vdots \\ x_n \end{bmatrix}_x$$

(n+1) x 1
matrix

theta transpose ضرب ال
x vector في ال
h(x) هيديني ال

Multivariate linear regression. ←



Machine Learning

Linear Regression with multiple variables

Gradient descent for multiple variables

Hypothesis: $h_{\theta}(x) = \theta^T x = \theta_0 x_0 + \theta_1 x_1 + \theta_2 x_2 + \dots + \theta_n x_n$

Handwritten notes: $x_0 = 1$ (with arrow pointing to x_0), θ (underlined), $n+1$ -dimensional vector

Parameters: $\theta_0, \theta_1, \dots, \theta_n$

Handwritten notes: θ (underlined), $n+1$ -dimensional vector

Cost function:

$$J(\theta_0, \theta_1, \dots, \theta_n) = \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$$

Handwritten notes: $J(\theta)$ (underlined), $J(\theta)$ (underlined)

Gradient descent:

Repeat {

$\rightarrow \theta_j := \theta_j - \alpha \left[\frac{\partial}{\partial \theta_j} J(\theta_0, \dots, \theta_n) \right]$

Handwritten notes: $J(\theta)$ (underlined), $J(\theta)$ (underlined)

}

(simultaneously update for every $j = 0, \dots, n$)

Gradient Descent

Previously ($n=1$):

Repeat {

→ $\theta_0 := \theta_0 - \alpha \underbrace{\frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})}_{\frac{\partial}{\partial \theta_0} J(\theta)}$

→ $\theta_1 := \theta_1 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_1^{(i)}$
(simultaneously update θ_0, θ_1)

}

New algorithm ($n \geq 1$):

Repeat {

→ $\theta_j := \theta_j - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_j^{(i)}$

(simultaneously update θ_j for $j = 0, \dots, n$)

}

→ $\theta_0 := \theta_0 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_0^{(i)}$

→ $\theta_1 := \theta_1 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_1^{(i)}$

→ $\theta_2 := \theta_2 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_2^{(i)}$

...



Machine Learning

Linear Regression with multiple variables

Gradient descent in
practice I: Feature Scaling


Feature Scaling

Idea: Make sure features are on a similar scale.

Dividing by max value
that the feature could
take

E.g. $x_1 = \text{size (0-2000 feet}^2\text{)}$ ←


$x_2 = \text{number of bedrooms (1-5)}$ ←



$$\rightarrow x_1 = \frac{\text{size (feet}^2\text{)}}{2000}$$

$$\rightarrow x_2 = \frac{\text{number of bedrooms}}{5}$$

$$0 \leq x_1 \leq 1 \quad 0 \leq x_2 \leq 1$$



Feature Scaling

Get every feature into approximately a $-1 \leq x_i \leq 1$ range.

$$x_0 = 1$$

$$0 \leq x_1 \leq 3 \quad \checkmark$$

$$-2 \leq x_2 \leq 0.5 \quad \checkmark$$

$$-100 \leq x_3 \leq 100 \quad \times$$

$$-0.0001 \leq x_4 \leq 0.0001 \quad \times$$

$$-3 \text{ to } 3 \quad \checkmark$$

$$-\frac{1}{3} \text{ to } \frac{1}{3} \quad \checkmark$$

Mean normalization

Replace x_i with $x_i - \mu_i$ to make features have approximately zero mean
(Do not apply to $x_0 = 1$).

E.g. $\rightarrow x_1 = \frac{\text{size} - 1000}{2000}$

Average size = 1000

$$x_2 = \frac{\# \text{bedrooms} - 5}{4}$$

1-5 bedrooms

$$\rightarrow [-0.5 \leq x_1 \leq 0.5, -0.5 \leq x_2 \leq 0.5]$$

$$x_1 \leftarrow \frac{x_1 - \mu_1}{S_1}$$

← avg value of x_1 in training set

range (max-min)
(or standard deviation)

$$x_2 \leftarrow \frac{x_2 - \mu_2}{S_2}$$



Machine Learning

Linear Regression with multiple variables


Gradient descent in practice II: Learning rate

Gradient descent

$$\rightarrow \theta_j := \theta_j - \alpha \frac{\partial}{\partial \theta_j} J(\theta)$$

- “Debugging”: How to make sure gradient descent is working correctly.
- How to choose learning rate α .

Making sure gradient descent is working correctly.




→ Example automatic convergence test:

→ Declare convergence if $J(\theta)$ decreases by less than 10^{-3} in one iteration.

No. of iterations differ according to applications


Making sure gradient descent is working correctly.



- For sufficiently small α , $J(\theta)$ should decrease on every iteration.
- But if α is too small, gradient descent can be slow to converge.

Summary:

- If α is too small: slow convergence.
- If α is too large: $J(\theta)$ may not decrease on every iteration; may not converge. (Slow converge also possible.)



To choose α , try

$\dots, \underline{0.001}, \underline{0.003}, \underline{0.01}, \underline{0.03}, \underline{0.1}, \underline{0.3}, \underline{1}, \dots$

Arrows indicate the sequence of values, with labels $\approx 3\times$ showing the progression from 0.001 to 0.003, 0.003 to 0.01, 0.01 to 0.03, 0.03 to 0.1, and 0.1 to 0.3.



Machine Learning

Linear Regression with multiple variables

Features and
polynomial regression

Housing prices prediction

$$h_{\theta}(x) = \theta_0 + \theta_1 \times \underbrace{\text{frontage}}_{x_1} + \theta_2 \times \underbrace{\text{depth}}_{x_2}$$

Area

$$x = \underline{\text{frontage} \times \text{depth}}$$

$$h_{\theta}(x) = \theta_0 + \theta_1 x$$

↖ land area



Polynomial regression



$$\theta_0 + \theta_1 x + \theta_2 x^2$$

$$\theta_0 + \theta_1 x + \theta_2 x^2 + \theta_3 x^3$$

Feature scaling

Size: 1-1000

Size²: 1-1,000,000

Size³: 1-10⁹

$$h_{\theta}(x) = \theta_0 + \theta_1 x_1 + \theta_2 x_2 + \theta_3 x_3$$

$$= \theta_0 + \theta_1 (\text{size}) + \theta_2 (\text{size})^2 + \theta_3 (\text{size})^3$$

$$\rightarrow x_1 = (\text{size})$$

$$\rightarrow x_2 = (\text{size})^2$$

$$\rightarrow x_3 = (\text{size})^3$$

Choice of features



$$\rightarrow h_{\theta}(x) = \theta_0 + \theta_1(\text{size}) + \theta_2(\text{size})^2$$

$$\rightarrow h_{\theta}(x) = \theta_0 + \theta_1(\text{size}) + \theta_2\sqrt{(\text{size})}$$





Machine Learning

Linear Regression with multiple variables

Normal equation

Gradient Descent



Normal equation: Method to solve for θ
analytically.

Intuition: If 1D ($\theta \in \mathbb{R}$)

$\rightarrow J(\theta) = a\theta^2 + b\theta + c$

$\frac{\partial}{\partial \theta} J(\theta) = \dots \stackrel{\text{set}}{=} 0$

Solve for θ



$\theta \in \mathbb{R}^{n+1}$ $J(\theta_0, \theta_1, \dots, \theta_m) = \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$

$\frac{\partial}{\partial \theta_j} J(\theta) = \dots \stackrel{\text{set}}{=} 0$ (for every j)

Solve for $\theta_0, \theta_1, \dots, \theta_n$

Examples: $m = 4$.

	Size (feet ²)	Number of bedrooms	Number of floors	Age of home (years)	Price (\$1000)
x_0	x_1	x_2	x_3	x_4	y
1	2104	5	1	45	460
1	1416	3	2	40	232
1	1534	3	2	30	315
1	852	2	1	36	178

$X = \begin{bmatrix} 1 & 2104 & 5 & 1 & 45 \\ 1 & 1416 & 3 & 2 & 40 \\ 1 & 1534 & 3 & 2 & 30 \\ 1 & 852 & 2 & 1 & 36 \end{bmatrix}$
 $m \times (n+1)$

$y = \begin{bmatrix} 460 \\ 232 \\ 315 \\ 178 \end{bmatrix}$
 m -dimensional vector

$\theta = (X^T X)^{-1} X^T y$

m examples $(x^{(1)}, y^{(1)}), \dots, (x^{(m)}, y^{(m)})$; n features.

$$\underline{x^{(i)}} = \begin{bmatrix} x_0^{(i)} \\ x_1^{(i)} \\ x_2^{(i)} \\ \vdots \\ x_n^{(i)} \end{bmatrix} \in \mathbb{R}^{n+1}$$

X
(design matrix)

$$= \begin{bmatrix} \text{---} (x^{(1)})^T \text{---} \\ \text{---} (x^{(2)})^T \text{---} \\ \vdots \\ \text{---} (x^{(m)})^T \text{---} \end{bmatrix}$$

$m \times (n+1)$

E.g. If $\underline{x^{(i)}} = \begin{bmatrix} 1 \\ x_1^{(i)} \end{bmatrix}$

$\Theta = (X^T X)^{-1} X^T y$

$$\begin{bmatrix} 1 & x_1^{(1)} \\ 1 & x_1^{(2)} \\ \vdots & \vdots \\ 1 & x_1^{(m)} \end{bmatrix} \quad \begin{bmatrix} y^{(1)} \\ y^{(2)} \\ \vdots \\ y^{(m)} \end{bmatrix}$$

$m \times 2$

$$\theta = (X^T X)^{-1} X^T y \leftarrow$$

$(X^T X)^{-1}$ is inverse of matrix $X^T X$.

Set $A = X^T X$

$$(X^T X)^{-1} = A^{-1}$$

Octave: `pinv(X' * X) * X' * y`

$$\text{pinv}(X^T * X) * X^T * y$$

$$\theta = (X^T X)^{-1} X^T y$$

$$\min_{\theta} J(\theta)$$

If using normal equation so feature scaling isn't important

X'	X^T
Feature Scaling	
	$0 \leq x_1 \leq 1$
	$0 \leq x_2 \leq 1000$
	$0 \leq x_3 \leq 10^{-5}$ ✓

With the normal equation, computing the inversion has complexity $O(n^3)$. So if we have a very large number of features, the normal equation will be slow. In practice, when n exceeds 10,000 it might be a good time to go from a normal solution to an iterative process.

m training examples, n features.

Cost of inversion

Gradient Descent

- • Need to choose α .
- • Needs many iterations.
- Works well even when n is large.

$n = 10^6$

Normal Equation

- • No need to choose α .
- • Don't need to iterate.
- Need to compute
- • $(X^T X)^{-1}$ ~~$n \times n$~~ $O(n^3)$
- Slow if n is very large.

$n = 100$

$n = 1000$

$n = 10000$



Machine Learning

Linear Regression with multiple variables

Normal equation
and non-invertibility
(optional)

Normal equation

$$\theta = \underline{(X^T X)^{-1} X^T y}$$

$$\underline{X^T X}$$

- What if $\boxed{X^T X}$ is non-invertible? (singular/degenerate)

Works properly even if $X^T X$ is singular

- Octave: `pinv(X' * X) * X' * y`

θ

Pseudo inverse

$\boxed{\text{pinv}}$
inv

What if $X^T X$ is non-invertible?

Causes of singular $X^T X$

- Redundant features (linearly dependent).

E.g. $x_1 = \text{size in feet}^2$

~~$x_2 = \text{size in m}^2$~~

$$x_1 = (3.28)^2 x_2$$

$$1\text{m} = 3.28\text{ feet}$$

$$\rightarrow m = 10 \leftarrow$$

$$\rightarrow n = 100 \leftarrow$$

$$\Theta \in \mathbb{R}^{101}$$

- Too many features (e.g. $m \leq n$).

Sometimes work but not always a good idea

- Delete some features, or use regularization.

↓ later

Solutions to the above problems include deleting a feature that is linearly dependent with another or deleting one or more features when there are too many features.