How does intermittent fasting affect memory and attention in adults?

ResearchGenie
AI-generated research draft

AI-generated research draft for human review. Not a verified or published research paper. Every claim must be checked against its cited source before any external use.

I. Introduction

Intermittent fasting (IF) effects on adult memory and attention are not uniform; they depend on population health status, activity level, protocol timing, and study duration. A meta-analysis of adults with obesity reported significant improvements in attention when IF was combined with hypoxia training relative to controls [1], whereas a pilot study in older adults (mean age ~70 years) found that an 8-week nightly fasting intervention increased cognitive function scores [2]. In contrast, a small study of young adults (mean age 25.8 years) observed decreases in Montreal Cognitive Assessment scores after both long-term and intermittent fasting, with the latter showing a numerically larger decline [3]. These divergent findings suggest that IF may benefit specific clinical cohorts—such as those with obesity or memory decline—but may not improve, and could transiently impair, cognition in younger, healthy populations.

The central tension driving this review is whether short-term cognitive gains reflect a generalizable benefit of IF or are context-dependent artifacts of population, protocol, and measurement. Evidence indicates that activity level modulates outcomes: sedentary elderly adults experienced declines in associative learning during Ramadan diurnal intermittent fasting [4], while physically active peers showed improvements in attention and executive function [4]. Similarly, the timing of the eating window appears critical; Sharifi et al. suggest that omitting dinner may confer neuroprotective effects linked to metabolic markers, whereas other temporal placements might not [5][3].

However, these interpretations are constrained by methodological limitations. Small sample sizes, short intervention durations (typically 4–12 weeks), and reliance on self-reported adherence compromise statistical power and generalizability [5][6][2]. The absence of control groups in several trials prevents causal inference, while heterogeneous outcome measures and lack of standardized IF definitions hinder cross-study comparisons [1][5][4]. Furthermore, most studies focus on English-language publications and North American or Middle Eastern cohorts, limiting applicability to diverse populations and cultural contexts [6][4].

This review synthesizes available evidence to map the conditions under which IF influences memory and attention in adults. It does not preclude future findings but emphasizes that current data support only narrow, context-specific conclusions. The analytical scope includes evaluating population differences, protocol variables, measurement validity, and the need for larger, longer, diverse trials to resolve unresolved questions about long-term safety and optimal implementation.

II. Background

Intermittent fasting (IF) and related time-restricted feeding protocols have been investigated for their impact on memory and attention in adults, yet findings vary by population, protocol, and duration. A meta-analysis of 28 studies involving 2,134 adults with obesity reported significant improvements in memory performance when intermittent hypoxia training was combined with fasting interventions compared to control conditions [1]. In contrast, a study of young female handball players found that baseline intraday differences in attention and simple reaction time were larger in the afternoon than in the morning before Ramadan but were reduced during the first ten days of Ramadan and blunted or reversed by the final ten days, suggesting protocol timing modulates cognitive effects [7]. In older adults (mean age 69.7 years) with self-reported memory decline, an 8-week nightly fasting intervention (14 h fasting/night) was associated with a statistically significant increase in cognitive function as measured by the Memory and Attention Phone Screener [2]. These divergent results highlight that population characteristics—such as age, baseline fitness, and metabolic status—likely shape cognitive outcomes."

Evidence also indicates domain-specific and heterogeneous effects within populations. In a sample of 16 young adults (mean age 25.8 years), both long-term fasting and intermittent fasting led to decreases in total Montreal Cognitive Assessment scores, with the decline numerically larger in the intermittent fasting group, though this difference was not statistically significant; short-term memory showed a large effect size reduction in the intermittent fasting condition [3]. A systematic review of eight studies in older adults (aged 60+) suggested that time-restricted eating and intermittent fasting may positively impact cognitive function but emphasized that outcomes appear contingent on the temporal placement

of the eating window rather than fasting duration alone [5]. These findings imply that not only the length of the fast but also when it occurs relative to circadian rhythms and daily activity patterns matters for cognition.

Methodological constraints limit confidence in generalizing these results. Reviews note that many studies rely on self-reported fasting practices, which may introduce bias, and that outcome measures vary widely across trials, hindering synthesis [5]. Small sample sizes (often 100 participants) and short intervention durations (typically 4–12 weeks) reduce power to detect subtle cognitive changes and impede assessment of long-term efficacy or safety [6]. Additionally, most included studies focused on overweight adults in the USA and were published in English, limiting generalizability to other age groups, populations, and food cultures [6]. Blinding is often not feasible in dietary interventions, further affecting validity [6]."

Population-level differences appear critical. In sedentary elderly adults, Ramadan diurnal intermittent fasting was associated with a significant decline in associative learning performance compared to baseline, whereas other cognitive domains remained unchanged; both groups experienced sleep deterioration and increased daytime sleepiness, with worse effects in the sedentary group [4]. By contrast, physically active elderly adults showed significant improvements in attention, executive function, inhibition, recognition memory, and associative memory under the same fasting regimen [4]. These divergent trajectories suggest that physical activity level may buffer or exacerbate cognitive responses to fasting, potentially by modulating sleep quality, metabolic stress, or neuroinflammatory markers. The authors hypothesized that moderate-intensity physical activity during Ramadan could mitigate negative effects on sleep and cognition, though this remains speculative [4]."

Mechanistic explanations remain largely inferential. Authors propose that IF may promote synaptic plasticity, neurogenesis, and neuroprotection, which are relevant to cognitive function, but these mechanisms require direct investigation in human trials [5]. Similarly, combined metabolic and cognitive benefits have been interpreted as evidence that fasting mitigates obesity-associated cognitive decline via reduced oxidative stress and neuroinflammation, yet organokine regulation pathways (e.g., BDNF) remain unconfirmed [1]. The lack of standardized definitions for time-restricted eating and intermittent fasting further complicates mechanistic comparisons across studies [5]. Consequently, while short-term metabolic and cognitive benefits are supported in specific contexts, the evidence is insufficient to establish long-term safety or optimal clinical protocols without further rigorous trials that address heterogeneity, duration, and population diversity [1][6].

III. Literature Review

Evidence regarding intermittent fasting (IF) and cognitive outcomes in adults reveals a stark divergence based on population health status and activity level. In adults with obesity, a meta-analysis of 28 studies involving over 2,000 participants reported significant improvements in attention performance when IF was combined with hypoxia training relative to control conditions [1]. Similarly, a pilot study in older adults (mean age ~70 years) with self-reported memory decline found that an 8-week prolonged nightly fasting intervention was associated with a statistically significant increase in cognitive function as measured by the Memory and Attention Phone Screener [2]. These findings suggest potential benefits for specific clinical cohorts, yet they contrast sharply with results from younger, healthy adults. Crhová and Kapounková (2020) observed that both long-term fasting and IF led to a decrease in total Montreal Cognitive Assessment (MoCA) scores in a group of young adults, noting that the decline was numerically larger in the IF group, though not statistically significant [3]. The authors interpreted this heterogeneity as evidence that fasting effects are domain-specific, with potential negative influences on short-term memory and generative abilities in younger populations [3].

This discordance is further illuminated by activity-dependent outcomes in elderly populations. Boujelbane et al. (2022) demonstrated that Ramadan diurnal intermittent fasting (RDIF) was associated with a significant decline in associative learning performance specifically in sedentary elderly adults, whereas physically active peers experienced significant improvements across attention, executive function, and memory domains [4]. The study highlighted that both groups suffered sleep deterioration, but the negative cognitive impact appeared contingent on physical activity levels rather than fasting duration alone. Systematic reviews reinforce these inconsistencies, noting that Time-Restricted Eating (TRE) outcomes in older adults may depend on the temporal placement of eating windows rather than fasting duration per se [5].

Methodological constraints complicate the synthesis of these divergent findings. Reviews consistently identify a lack of standardized definitions for IF protocols and diverse outcome measures as primary barriers to generalizability [5][6]. Furthermore, reliance on self-reported adherence introduces bias, while small sample sizes in many trials limit statistical power to detect subtle cognitive shifts [6]. The absence of randomized controlled trials (RCTs) in the existing literature further warrants cautious interpretation of reported associations [5]. Consequently, while short-term metabolic and cognitive benefits are supported for specific subgroups like those with obesity or high physical activity, current evidence is insufficient to establish long-term safety or optimal clinical protocols without addressing these critical methodological gaps.

IV. Discussion

The relationship between intermittent fasting and cognitive function in adults is characterized by significant heterogeneity across populations, intervention protocols, and outcome measures. A meta-analysis of 28 studies involving adults with obesity reported that combined Intermittent Hypoxia Training and Intermittent Fasting yielded significant improvements in memory and attention compared to controls [1]. However, this finding is accompanied by substantial heterogeneity ($I^2 > 50\%$), suggesting that effect sizes vary based on study populations, protocols, or measurement tools [1]. In contrast, a systematic review of eight studies focused specifically on older adults (aged 60+) suggests that Time-Restricted Eating and Intermittent Fasting may positively impact cognitive function, though the authors note the need for additional research to fully comprehend this relationship [5]. This positive association in older adults contrasts with findings from a study of younger adults (mean age 25.8 years), where both long-term fasting and intermittent fasting led to a decrease in total Montreal Cognitive Assessment scores, with the decline being numerically larger in the intermittent fasting group, although this difference was not statistically significant [3]. Furthermore, research on sedentary elderly adults indicates that Ramadan diurnal intermittent fasting was associated with a significant decline in associative learning performance compared to baseline, while other cognitive domains remained unchanged [4]. Conversely, physically active elderly adults experienced significant improvements in attention, executive function, inhibition, and memory under the same fasting regimen [4]. These divergent outcomes highlight that the effect of fasting on cognition is likely contingent on factors such as age, physical activity level, and the temporal placement of eating windows rather than fasting duration alone [5]. The observed improvements in metabolic and cognitive benefits are interpreted by some authors as evidence that these interventions may mitigate obesity-associated cognitive decline through mechanisms such as reduced oxidative stress and neuroinflammation, though they explicitly call for mechanistic investigations into organokine regulation to confirm these explanations [1]. Despite the potential for benefit, the current literature contains critical methodological limitations that constrain the generalizability of findings. A primary limitation is the lack of standardized definitions and protocols for Time-Restricted Eating and Intermittent Fasting, which makes it challenging to compare findings across studies accurately [5]. Additionally, the diverse outcome measures used across studies hinder the synthesis of results and limit the generalizability of findings regarding cognitive outcomes [5]. Variations in participant characteristics, such as age, gender, and health status, further complicate the interpretation of results regarding cognitive function [5]. The reliance on self-reported Intermittent Fasting practices across several studies introduces the possibility of bias, warranting cautious interpretation of reported associations [5]. Moreover, all reviewed studies were published in English, representing a possible bias as relevant materials on intermittent fasting exist in other languages such as Chinese and Hindi [6]. The short duration of most studies also impedes the detection of significant changes in some parameters while lacking data on long-term efficacy [6]. Finally, underreporting of adverse events across the included literature constitutes a significant gap, necessitating the development and validation of standardized safety monitoring frameworks for future trials [1]. Consequently, while current evidence supports the efficacy of these interventions for short-term metabolic and cognitive benefits in specific populations, the findings are not yet sufficient to establish long-term safety or optimal clinical implementation protocols without further rigorous trials [1].

V. Research Gap

A. The Activity-Dependent Paradox in Cognitive Outcomes

A fundamental contradiction exists between studies of physically active versus sedentary older adults. Paper [1] reports significant cognitive improvements (executive function, attention) only in the physically active group during Ramadan fasting, while the sedentary group showed declines in associative learning. Conversely, Paper [6] found cognitive improvements in non-active older adults with memory decline using a home-based protocol. However, no study directly compares these populations under identical conditions to isolate the effect of physical activity from the effects of fasting itself.

Evidence: Paper [1] shows active elderly improved on attention ($p=0.005$) while sedentary peers declined in associative learning. Paper [6] reports increased memory and attention in non-active older adults with self-reported decline. The corpus lacks a study controlling for activity levels to determine if fasting benefits are strictly dependent on concurrent exercise.

B. The Obesity-Cognition Interaction Gap

While obesity is linked to cognitive decline, the corpus fails to clarify whether IF improves cognition in obese adults independently of weight loss. Paper [2] reviews metabolic and cognitive health in obesity but notes that included studies often conflate metabolic improvements with cognitive gains. It remains unclear if IF restores attention/memory in obese individuals even without significant BMI reduction.

Evidence: Paper [2] includes a meta-analysis of 2,134 participants with obesity but does not explicitly separate cognitive outcomes by degree of weight loss or metabolic improvement. Paper [7] shows BMI reduction without cognitive differences, suggesting a disconnect between metabolic and cognitive metrics in non-obese samples.

C. The Timing-Window Uncertainty

There is no consensus on the optimal timing of fasting windows for cognitive performance. Paper [3] demonstrates that intraday differences in attention and reaction time are blunted or reversed during Ramadan, suggesting circadian disruption may impair specific tasks. However, Papers [5] and [7] suggest benefits or neutral effects with 16:8 protocols without specifying if these windows align with individual chronotypes.

Evidence: Paper [3] finds morning-afternoon performance differences are reduced or reversed during fasting periods in athletes. Paper [5] calls for research on 'timing of the eating window' but provides no data on optimal times. Paper [7] uses a standard 16:8 protocol without assessing circadian alignment.

D. The Mechanistic Black Box

The corpus lacks any human study linking specific molecular pathways (e.g., BDNF, insulin sensitivity) to observed cognitive changes. Paper [2] explicitly calls for mechanistic investigations into organokine regulation but provides none. This prevents understanding why some populations improve while others decline.

Evidence: Paper [2] states: 'Priority research areas should include: (1) Mechanistic investigations into organokine regulation (particularly BDNF and leptin pathways);' yet no included study measures these biomarkers alongside cognitive tests. Paper [8] discusses cellular pathways theoretically but offers no empirical human data.

E. The Domain-Specific Vulnerability Gap

Studies measure different cognitive domains inconsistently. Paper [1] measures executive function and associative memory; Paper [4] focuses on short-term memory and verbal production; Paper [7] uses working memory and attentional control tasks. No study compares the relative vulnerability of these domains to fasting stress.

Evidence: Paper [1] reports improvements in executive function but declines in associative learning for sedentary groups. Paper [4] finds short-term memory decreases more than other functions in IF groups. Paper [7] finds no group differences in working memory or attentional control. The lack of standardized, multi-domain batteries across studies prevents meta-analysis of domain-specific effects.

VI. Proposed Novelty and Contribution

Research addressing the activity-dependent paradox (Gap 1) and domain-specific vulnerability (Gap 4) would be novel as no prior study in the corpus directly compares sedentary vs. active older adults under identical fasting conditions or standardizes cognitive domains across populations.

Caveats: Novelty is hedged because the corpus is small and may not represent the entire field. If external literature exists outside these 8 papers, novelty claims may be overstated.

VII. Limitations

The validity of findings regarding intermittent fasting (IF) and cognitive function is constrained by methodological limitations inherent in the current literature. A primary concern across multiple studies is the reliance on small sample sizes, which compromises statistical power and limits generalizability [5][3][6]. This issue is compounded by short intervention durations, often ranging from 4 to 12 weeks, which may fail to capture long-term cognitive impacts or detect late-onset adverse effects [1][6]. Furthermore, the absence of control groups in several pilot trials prevents researchers from distinguishing true intervention efficacy from placebo effects or natural recovery trajectories [2].

Measurement strategies also introduce significant uncertainty. Many studies depend on self-reported adherence to fasting protocols and subjective questionnaires for sleep and cognitive assessment rather than objective biomarkers or actigraphy, raising concerns about reporting bias and measurement error [5][4]. Additionally, the lack of standardized definitions for time-restricted eating (TRE) and IF protocols hinders cross-study comparisons, while variations in participant characteristics—such as age, gender, health status, and physical activity levels—further complicate the interpretation of heterogeneous outcomes [5][4].

Beyond methodological constraints, clinical applicability is limited by safety profiles and adherence challenges. Adherence to IF regimens typically declines after 12 weeks, potentially undermining long-term benefits [6]. Common adverse effects include dehydration, headaches, dizziness, and mood disturbances, which may influence both cognitive performance and regimen sustainability [6]. Consequently, current evidence supports short-term metabolic and cognitive benefits but remains insufficient to establish long-term safety or optimal clinical protocols without further rigorous trials involving diverse populations and longer follow-up periods [1][6].

VIII. Conclusion

The evidence indicates that intermittent fasting (IF) effects on adult memory and attention are not uniform but depend critically on population health status, activity level, protocol timing, and study duration. Robust support exists for context-specific benefits: a meta-analysis of adults with obesity reported significant improvements in attention when IF was combined with hypoxia training relative to controls [1], and a pilot study in older adults (mean age ~70 years) found that an 8-week nightly fasting intervention increased cognitive function scores [2]. These findings suggest potential utility for specific clinical cohorts, such as those with obesity or self-reported memory decline. However, these positive outcomes contrast sharply with results from younger, healthy populations; a study of young adults (mean age 25.8 years) observed decreases

in Montreal Cognitive Assessment scores after both long-term and intermittent fasting, with the latter showing a numerically larger decline [3]. This divergence implies that IF may transiently impair cognition or offer no benefit in younger, healthy groups, whereas it may support cognitive function in older or metabolically compromised cohorts.

Activity level and metabolic context appear to modulate these outcomes significantly. Sedentary elderly adults experienced declines in associative learning during Ramadan diurnal intermittent fasting [4], while physically active peers showed improvements in attention and executive function [4]. Similarly, the timing of the eating window influences results; omitting dinner may confer neuroprotective effects linked to metabolic markers, whereas other temporal placements might not [5][3]. Yet, these interpretations are constrained by substantial methodological heterogeneity. The validity of findings is compromised by small sample sizes, short intervention durations (typically 4–12 weeks), and reliance on self-reported adherence rather than objective biomarkers or actigraphy [5][6][2]. The absence of control groups in several trials prevents causal inference, while heterogeneous outcome measures and lack of standardized IF definitions hinder cross-study comparisons [1][5][4]. Furthermore, most studies focus on English-language publications and North American or Middle Eastern cohorts, limiting applicability to diverse populations.

Consequently, what remains conditional is the generalizability of short-term cognitive gains. Current data support only narrow, context-specific conclusions rather than a universal benefit. What cannot yet be concluded includes the long-term safety profile of IF protocols, the optimal duration required for sustained cognitive effects, and whether observed benefits in specific cohorts reflect a true biological mechanism or are artifacts of population selection and measurement bias. The main boundary conditions are population age and health status, physical activity level, fasting window placement, and intervention length. Sources of uncertainty stem from small samples, short durations, subjective reporting, lack of control groups, and non-standardized protocols.

Future research priorities must address these bottlenecks to resolve unresolved questions. First, larger, longer, diverse trials are needed to distinguish transient effects from sustained benefits and to assess long-term safety [1][6]. Second, studies must standardize definitions of time-restricted eating (TRE) and IF protocols to enable valid cross-study comparisons [5][4]. Third, trials should incorporate objective measures such as actigraphy and biomarkers to reduce reporting bias [5][4]. Fourth, research must explicitly test the moderating roles of activity level and metabolic status across diverse populations, including non-Western cohorts, to determine whether current findings apply broadly or are culturally and demographically specific [6][4]. Finally, causal mechanisms linking fasting windows to cognitive changes require investigation beyond correlation, particularly regarding neuroprotective pathways versus potential acute stress responses in younger adults.

References

[1] [1] J. Guo, N. Zhang, and J. Chen, "Comprehensive impact of Intermittent Hypoxia Training and Intermittent Fasting on metabolic and cognitive health in adults with obesity: an umbrella systematic review and meta-analysis," Frontiers in Nutrition, 2025, doi: 10.3389/fnut.2025.1664600.

[2] [2] D. L. James, L. Larkey, and M. Maxfield, "Prolonged nightly fasting in older adults with memory decline: A single-group pilot study exploring changes in cognitive function and cardiometabolic risk factors," Journal of Clinical and Translational Science, 2024, doi: 10.1017/cts.2024.676.

[3] [3] M. Crhová and K. Kapounková, "Influence of long-term fasting and intermittent fasting on the cognitive abilities," Studia sportiva, 2020, doi: 10.5817/sts2020-1-2.

[4] [4] M. A. Boujelbane, K. Trabelsi, and H. Jahrami, "Time-restricted feeding and cognitive function in sedentary and physically active elderly individuals: Ramadan diurnal intermittent fasting as a model," Frontiers in Nutrition, 2022, doi: 10.3389/fnut.2022.1041216.

[5] [5] S. Sharifi, F. Rostami, and K. B. Khorzoughi, "Effect of time-restricted eating and intermittent fasting on cognitive function and mental health in older adults: A systematic review," Preventive Medicine Reports, 2024, doi: 10.1016/j.pmedr.2024.102757.

[6] [6] S. Brogi, R. Tabanelli, and S. Puca, "Intermittent Fasting: Myths, Fakes and Truth on This Dietary Regimen Approach," Foods, 2024, doi: 10.3390/foods13131960.

[7] [7] H. Bougrine, M. Cherif, and H. Chtourou, "Does Ramadan intermittent fasting affect the intraday variations of cognitive and high-intensity short-term maximal performances in young female handball players?," Biological Rhythm Research, 2023, doi: 10.1080/09291016.2023.2198794.