

Main page Contents Featured content Current events Random article Donate to Wikipedia Wikipedia store

Interaction

Help About Wikipedia Community portal Recent changes Contact page

Tools

What links here Related changes Upload file Special pages Permanent link Page information Wikidata item Cite this page

Print/export

Article Talk

Read

Edit | View history

Search Wikipedia

Pick's theorem

From Wikipedia, the free encyclopedia

For the theorem in complex analysis, see Schwarz lemma § Schwarz-Pick theorem.

Given a simple polygon constructed on a grid of equal-distanced points (i.e., points with integer coordinates) such that all the polygon's vertices are grid points, Pick's theorem provides a simple formula for calculating the area A of this polygon in terms of the number i of lattice points in the interior located in the polygon and the number b of lattice points on the boundary placed on the polygon's perimeter:[1]


$$A=i+rac{b}{2}-1.$$


In the example shown, we have i = 7 interior points and b = 8 boundary points, so the area is $A = 7 + \frac{8}{2} - 1$ = 7 + 4 - 1 = 10 square units.

Note that the theorem as stated above is only valid for *simple* polygons, i.e., ones that consist of a single piece and do not contain holes. For a polygon that has h holes, with a boundary in the form of h+1 simple closed curves, the slightly more complicated formula $i + \frac{b}{2} + h - 1$ gives the area.

The result was first described by Georg Alexander Pick in 1899.^[2] The Reeve tetrahedron shows that there is no analogue of Pick's theorem in three dimensions that expresses the volume of a polytope by counting its interior and boundary points. However, there is a generalization in higher dimensions via Ehrhart polynomials. The formula also generalizes to surfaces of polyhedra.







The triangle with vertices at the lower left, lower right, and upper right points has i = 12 and b = 14, giving by Pick's theorem $A = i + \frac{b}{2} - 1 = 18$; this is Create a book

Download as PDF

Printable version

Languages

العربية

Deutsch

Español

Français

한국어

Italiano

Português

Русский

中文

XA 12 more



O

- 2 See also
- 3 References
- 4 External links

confirmed by the triangle area formula $\frac{1}{2}$ × base × height = $\frac{1}{2}$ × 9 × 4 = 18.

Proof [edit]

Consider a polygon P and a triangle T, with one edge in common with P. Assume Pick's theorem is true for both P and T separately; we want to show that it is also true for the polygon PT obtained by adding T to P. Since P and T share an edge, all the boundary points along the edge in common are merged to interior points, except for the two endpoints of the edge, which are merged to boundary points. So, calling the number of boundary points in common c, we have [3]

$$i_{PT}=i_P+i_T+(c-2)$$

and

$$b_{PT} = b_P + b_T - 2(c-2) - 2.$$

From the above follows

$$i_P+i_T=i_{PT}-(c-2)$$

and

$$b_P + b_T = b_{PT} + 2(c-2) + 2.$$

Since we are assuming the theorem for P and for T separately,

$$egin{aligned} A_{PT} &= A_P + A_T \ &= \left(i_P + rac{b_P}{2} - 1
ight) + \left(i_T + rac{b_T}{2} - 1
ight) \ &= i_P + i_T + rac{b_P + b_T}{2} - 2 \ &= i_{PT} - (c - 2) + rac{b_{PT} + 2(c - 2) + 2}{2} - 2 \ &= i_{PT} + rac{b_{PT}}{2} - 1. \end{aligned}$$

Therefore, if the theorem is true for polygons constructed from n triangles, the theorem is also true for polygons constructed from n+1 triangles. For general polytopes, it is well known that they can always be triangulated. That this is true in dimension 2 is an easy fact. To finish the proof by mathematical induction, it remains to show that the theorem is true for triangles. The verification for this case can be done in these short steps:

- observe that the formula holds for any unit square (with vertices having integer coordinates);
- deduce from this that the formula is correct for any rectangle with sides parallel to the axes;
- deduce it, now, for right-angled triangles obtained by cutting such rectangles along a diagonal;
- now any triangle can be turned into a rectangle by attaching such right triangles; since the formula is correct for the right triangles and for the rectangle, it also follows for the original triangle.

The last step uses the fact that if the theorem is true for the polygon PT and for the triangle T, then it's also true for P; this can be seen by a calculation very much similar to the one shown above.

See also [edit]

Integer points in convex polyhedra

References [edit]

- 1. A Trainin, J. (November 2007). "An elementary proof of Pick's theorem". Mathematical Gazette. 91 (522): 536-540.
- 3. A Beck, Matthias; Robins, Sinai (2007). Computing the Continuous Discretely, Integer-point enumeration in polyhedra. Undergraduate Texts in Mathematics. New York: Springer-Verlag. ch. 2. ISBN 978-0-387-29139-0. MR 2271992 2.

External links [edit]

- Pick's Theorem (Java)
 at cut-the-knot
- Pick's Theorem proof by Tom Davis
- Pick's Theorem by Ed Pegg, Jr., the Wolfram Demonstrations Project.
- Weisstein, Eric W. "Pick's Theorem" №. *MathWorld*.

Categories: Digital geometry | Lattice points | Euclidean plane geometry | Theorems in geometry | Area | Polygons | Analytic geometry

This page was last edited on 31 March 2018, at 13:52 (UTC).

Text is available under the Creative Commons Attribution-ShareAlike License; additional terms may apply. By using this site, you agree to the Terms of Use and Privacy Policy. Wikipedia® is a registered trademark of the Wikimedia Foundation, Inc., a non-profit organization.

Privacy policy About Wikipedia Disclaimers Contact Wikipedia Developers Cookie statement Mobile view



