


ECODOME Research Submission

Rooted Resilence

Mangroves and Sediment dynamics during coastal storm

Nethra Shree. S

Environmental Research Paper

May 23, 2026

i) Table of Contents:

Section	Page No.
i. Acknowledgement	2
1. Abstract	3
2. Introduction	3
3. Research Objective	4
4. Literature Overview	4
5. Theoretical Framework	6
6. Methodology	6
6.1 Methodology Note	7
7. Innovation Aspect	7
8. Research Significance	7
9. Results / Findings	8
9.1 Impact of Mangrove Density on Sediment Stability	8

9.2 Influence of Storm Intensity on Coastal Change	9
9.3 Case Study: Sundarbans Mangrove Ecosystem	10
9.4 Case Study: Kerala Coastal Region	11
9.5 Comparison Between Kerala and Sundarbans	12
10. Discussion	12
11. Limitations of the Study	13
12. Policy Implications	13
13. Conclusion	14
14. Future Scope	15
ii) References	15
iii) Appendix A: Glossary of Key Terms	19
iv) Appendix B: Figures Overview	20

ii) Acknowledgement: I would like to express my gratitude to the ECODOME research initiative for providing the opportunity to explore environmental resilience and coastal sustainability through this study. I also acknowledge the use of secondary research sources, satellite-based datasets and scientific literature that supported the completion of this paper.

1. Abstract:

Coastal regions are increasingly exposed to climate change, rising sea levels, and stronger storm events, all of which are altering shoreline dynamics and increasing sediment movement. Mangrove ecosystems play a significant role in reducing these impacts by attenuating wave energy, slowing water flow, and trapping sediments through their dense root structures. This study investigates the relationship between mangrove coverage and sediment behaviour during storm events, with particular emphasis on how vegetation density influences shoreline stability. The analysis is based on secondary environmental studies and documented case examples from cyclone-prone coastal regions. Findings indicate that areas with higher mangrove density consistently show reduced sediment displacement and improved shoreline stability during extreme weather conditions. This study reinforces the importance of mangroves as nature-based coastal protection systems and highlights their relevance in climate adaptation and sustainable coastal management.

Keywords: Mangroves; coastal erosion; sediment stability; storm surge; cyclone impact; NDVI analysis; ecosystem resilience; coastal protection; climate change adaptation; nature-based solutions.

2. Introduction:

Coastal regions are increasingly facing pressure from climate change, rising sea levels, and more frequent storms. In many tropical areas, these forces not only reshape shorelines gradually but also cause sudden erosion and sediment movement during extreme weather events, making both ecosystems and coastal settlements more vulnerable.

Mangrove ecosystems play a key role in reducing this impact. Their dense root systems reduce wave energy, slow water movement, and trap sediments, helping to stabilise coastlines during storm conditions. They also act as a natural buffer between land and sea, supporting both ecological balance and coastal protection.

However, mangrove cover has declined in several regions due to urban expansion, land conversion, and unplanned coastal development. This reduction has weakened natural protection systems and increased exposure to erosion and sediment loss during storms. While the ecological importance of mangroves is well

established, their specific influence on sediment behaviour during storm events still requires closer examination.

Understanding this relationship is important for improving coastal management strategies. As climate-related risks continue to intensify, conserving and restoring mangrove ecosystems may play a crucial role in strengthening long-term coastal resilience.

The paper is organised into several sections. The literature review examines existing research on mangrove ecosystems and coastal sediment dynamics, followed by the methodology used for the study. The subsequent sections present the findings and case-based observations, discuss their broader implications, and conclude by highlighting the significance of mangrove conservation in sustainable coastal management.

3. Research Objective:

To analyse the relationship between mangrove density and coastal stability under varying storm conditions using secondary environmental data and satellite-based observations, and to assess the role of mangroves in reducing sediment movement and coastal erosion.

4. Literature Overview:

Research on coastal ecosystems has consistently highlighted the protective role of mangrove forests in stabilizing shoreline environments. Kathiresan and Bingham (2001) found that mangrove root systems significantly enhance sediment deposition by reducing nearshore water velocity, which contributes to long-term shoreline formation and stability in tropical coastal regions. Their work also emphasized the ability of mangroves to act as natural buffers against physical coastal disturbances.

Similarly, Alongi (2008) reported that mangrove forests provide effective protection against storm surges, particularly in cyclone-prone regions of Southeast Asia. His study observed that coastal zones with intact mangrove cover experienced reduced bstructural damage and lower levels of sediment disturbance during extreme weather events compared to degraded areas.

Environmental assessments by organisations such as the United Nations Environment Programme (UNEP, 2014) further support these findings, noting that mangrove density is directly linked to improved sediment retention and reduced coastal erosion. However, while existing literature strongly establishes the general protective function of mangroves, fewer studies have examined in detail how variations in mangrove density specifically influence sediment behaviour during high-intensity storm conditions. This gap is important in the context of increasing storm frequency and intensity driven by climate change.

In addition to these findings, recent coastal studies highlight that mangroves are most effective when considered as part of a wider coastal system. They often interact with nearby ecosystems such as tidal marshes and seagrass beds, which together strengthen sediment retention and reduce wave energy along shorelines. This suggests that mangrove performance is influenced not only by their structure and density, but also by their ecological surroundings. As a result, there is a growing shift in research toward understanding mangroves as interconnected components of coastal systems rather than isolated protective zones.

Recent studies have also looked at how mangroves respond to environmental stress and human activity over time. Friess et al (2019) found that their effectiveness in protecting coastlines depends on local conditions such as sediment supply, tidal range, and coastal development. In areas where natural sediment flow is disrupted or mangrove forests are fragmented, their ability to stabilise shorelines tends to reduce. This shows that mangrove performance is shaped not only by their natural structure but also by external environmental and human factors. These findings highlight the need to consider both ecological and human influences when studying long-term coastal protection.




Figure 1: Mangrove Root System and Sediment Trapping Mechanism

This figure illustrates how mangrove prop roots reduce wave energy and trap sediments, contributing to coastal stability during storm events.

Source: Adapted from ResearchGate (for educational purposes only)

5. Theoretical Framework:

This study is guided by the concept of ecosystem-based coastal resilience, which explains how natural systems such as mangroves contribute to shoreline stability and protection against environmental disturbances. It also draws on the idea of nature-based solutions, where ecological systems are used to reduce the impact of climate-related hazards. In this context, mangrove ecosystems are understood as dynamic protective structures that interact with hydrological and sedimentary processes to influence coastal stability.

6. Methodology:

Building upon the existing literature on mangrove ecosystems and coastal dynamics, this study adopts a qualitative research design based entirely on secondary data analysis. Relevant information was gathered from peer-reviewed journal articles, published environmental studies, and reports from recognised organisations such as the United Nations Environment Programme (UNEP), focusing on mangrove ecosystems and coastal sediment dynamics.

A comparative analytical approach was employed to examine existing findings across multiple cyclone-prone coastal regions. This involved evaluating documented case studies that highlight differences in sediment behaviour and shoreline stability between areas with varying levels of mangrove coverage.

Particular attention was given to studies discussing hydrodynamic processes, wave attenuation, and sediment retention mechanisms associated with mangrove root systems.

The collected data was systematically reviewed to identify recurring patterns and relationships between mangrove density and sediment displacement during storm events. No primary data collection or fieldwork was conducted; instead, the study relies exclusively on validated secondary sources to derive conclusions regarding coastal resilience and ecosystem performance under extreme weather conditions.

6.1 Methodology Note:

This study adopts a secondary-data-based analytical approach to ensure scalability and consistency in assessing coastal processes across multiple regions. The focus is on synthesising validated environmental findings to support comparative interpretation of mangrove ecosystem performance under storm conditions.

7. Innovation Aspect:

This study uses satellite-based remote sensing data (Landsat) along with comparative analysis to understand how mangrove density affects coastal stability during storm events. Instead of only describing mangroves as protective barriers, the research focuses on how different levels of mangrove density influence erosion and sediment movement, providing a clearer and more practical way to assess coastal resilience.

8. Research Significance:

This study contributes to the growing understanding of nature-based coastal protection by highlighting the importance of mangrove density in reducing storm-induced shoreline instability. The findings support the use of ecosystem-based approaches in climate adaptation and sustainable coastal management.

9. Results / Findings:

9.1 Impact of Mangrove Density on Sediment Stability:

The analysis of secondary environmental studies and coastal case examples shows a clear pattern between mangrove cover and sediment stability during storm events. Areas with healthier and denser mangrove forests generally experience less sediment movement compared to regions where mangroves are degraded or sparse.

In cyclone-affected coastal regions, mangrove-rich zones tend to remain relatively more stable during storms, with reduced erosion and slower changes in shoreline shape. This pattern is consistently observed in regions where mangrove cover remains relatively dense.

On the other hand, locations with limited mangrove cover show more noticeable shoreline changes after storm events, including higher sediment loss and greater physical disturbance of the coast. Overall, the findings suggest that mangrove density plays an important role in how stable a coastline remains during extreme weather conditions.



Figure 2: Coastal Erosion Comparison in Mangrove and Non-Mangrove Regions

This figure illustrates differences in shoreline stability between areas with dense mangrove cover and regions with limited or no mangrove vegetation, highlighting variations in erosion patterns during storm conditions.

Source: Elsevier (ScienceDirect), academic research article (peer-reviewed environmental study)

9.2 Influence of Storm Intensity on Coastal Change:

Further comparison of case examples also suggests that the impact of mangrove coverage varies depending on the intensity of storm events. During stronger cyclonic conditions, regions with dense mangrove forests still experience some level of shoreline change, but the extent of erosion and sediment loss remains comparatively lower than in areas with limited vegetation. This indicates that while mangroves do not completely prevent coastal alteration, they significantly reduce the severity of storm-induced changes across different levels of storm intensity.




Figure 3: Mangrove Density Distribution Based on Vegetation Index (NDVI Analysis)

This figure represents spatial variation in mangrove density across coastal regions using vegetation index analysis, where higher values indicate healthier and denser mangrove cover.

Source: *European Journal of Remote Sensing*, Taylor & Francis Group (remote sensing dataset / publication)

9.3 Case study: Sundarbans Mangrove Ecosystem

The Sundarbans, located across India and Bangladesh, represent one of the largest mangrove ecosystems in the world and serve as an important example of natural coastal protection. This region is frequently exposed to cyclones and tidal disturbances, making it highly vulnerable to shoreline erosion and sediment displacement. Studies conducted after major cyclonic events, including Cyclone Amphan, observed that areas with dense mangrove cover experienced comparatively lower physical damage and reduced coastal erosion than more exposed coastal zones. The dense root networks of mangroves in the Sundarbans help slow wave movement and retain sediments, contributing to greater shoreline stability during extreme weather conditions. The case of the Sundarbans therefore demonstrates the practical significance of mangrove ecosystems in reducing the environmental impact of storms and supporting long-term coastal resilience.




Figure 4: Sundarbans Mangrove Extent

Shows the geographical distribution of mangrove forests across the Sundarbans region.

Source: USGS/NASA Landsat via Google Earth Engine

In addition to its ecological significance, the Sundarbans also provides a clear example of how mangrove ecosystems interact with human settlements in high-risk coastal zones. A large population living in and around this region depends on agriculture, fishing, and small-scale livelihoods that are frequently affected by cyclonic activity. During major storm events, areas closer to intact mangrove belts have been observed to experience relatively reduced inland water intrusion compared to more exposed villages located near degraded forest zones. This difference highlights the protective role of mangroves not only in controlling physical shoreline changes but also in reducing the extent of storm-related disruption to human activity.

Furthermore, long-term observations in the region suggest that mangrove loss due to illegal logging and land conversion has increased vulnerability in certain coastal stretches. In contrast, areas where conservation efforts and reforestation initiatives have been implemented show signs of improved shoreline stability over time. This indicates that restoration efforts can play a meaningful role in gradually rebuilding natural coastal resilience. The Sundarbans therefore serves not only as a natural laboratory for studying mangrove dynamics, but also as an example of how ecological conservation directly connects to human safety and sustainability.

9.4 Case Study: Kerala Coastal Region (India)

The coastal region of Kerala, India, provides another important example of how mangrove ecosystems influence shoreline stability during extreme weather events. This region is frequently affected by monsoon-driven flooding and periodic cyclonic disturbances, making it a relevant site for studying coastal change. In several coastal stretches of Kerala, areas with remaining mangrove patches have shown comparatively better resistance to shoreline erosion during storm surges than nearby regions where mangroves have been heavily cleared for development and aquaculture.

Field-based observations from cyclone-impacted zones suggest that mangrove-fringed coastlines experience reduced wave penetration and slower inland water movement during extreme rainfall events. This contributes to comparatively lower levels of sediment displacement and coastal damage. However, in areas where mangroves have been replaced by human settlements or aquaculture farms, the absence of natural buffering systems has led to more pronounced erosion and greater vulnerability to flooding.

This case highlights that even fragmented mangrove systems can still provide measurable protective benefits, but their effectiveness is significantly reduced when ecosystem continuity is disrupted. The Kerala coastline therefore reinforces the importance of conserving remaining mangrove patches and restoring degraded coastal vegetation to improve long-term resilience against climate-related coastal hazards.

9.5 Comparison between Kerala and Sunderbans:

A comparative analysis between the Kerala coastline and the Sundarbans highlights variations in mangrove effectiveness under different coastal conditions. The Sundarbans, with its extensive and continuous mangrove cover, demonstrates significantly higher resilience to cyclonic storms, with reduced shoreline erosion and better sediment retention. In contrast, the Kerala coastline, where mangrove patches are more fragmented and unevenly distributed, shows comparatively higher vulnerability to coastal erosion and storm-induced shoreline changes. This difference suggests that mangrove continuity and density play a critical role in determining the level of coastal protection, rather than the presence of mangroves alone.

10. Discussion:

The findings of this study support the idea that mangrove ecosystems play an important role in protecting coastal areas during storm events. In regions with dense mangrove cover, there is clearly less sediment movement, which can be linked to how mangrove roots slow down waves and reduce their impact on the shoreline.

On the other hand, areas with fewer or damaged mangroves tend to experience more noticeable changes during storms, including stronger erosion and quicker shifts in shoreline shape. This suggests that when mangroves are missing or reduced, coastal areas become much more exposed to the effects of extreme weather.

One interesting point from the findings is that mangrove protection does not work in a simple “present or absent” way. Instead, the level of mangrove cover seems to matter significantly. Areas with thicker mangrove growth appear to cope better with storm conditions, while partially degraded areas still face

significant damage. This shows that even minor changes in mangrove density can influence the stability of the coastline.

Overall, the results highlight that mangroves are not just helpful as general protection, but their density plays a key role in the ability of coastal areas to withstand storms. This makes their conservation and restoration especially important in regions facing increasing climate-related risks.

Beyond the immediate physical effects observed, these findings also highlight broader implications for coastal resilience under changing climate conditions. As storm intensity and sea-level rise continue to increase, the protective role of mangroves becomes more significant not only in reducing erosion but also in maintaining the long-term stability of coastal ecosystems. This indicates that mangrove conservation should be viewed as a key strategy in climate adaptation planning, rather than only as an environmental preservation effort. Strengthening and restoring mangrove belts may therefore offer a sustainable and nature-based solution to reducing future coastal vulnerability.

11. Limitations of the Study:

This study is based on secondary data sources, including published research and documented case studies, which limits direct field-level validation of mangrove density and sediment movement. Variations in data collection methods across different studies may also influence the consistency of comparisons. In addition, local geographical differences, such as tidal range, coastal shape, and human activity, can affect sediment behaviour independently of mangrove coverage. Therefore, while the findings show a clear overall pattern, they should be interpreted with consideration of these contextual variations.

12. Policy Implications:

The findings of this study highlight the need to recognise mangrove ecosystems as a central component of coastal management strategies. Given their role in reducing sediment displacement and stabilising shorelines during storm events, the protection of existing mangrove forests should be prioritised within national and regional environmental policies.

Coastal development planning should incorporate strict regulations to prevent further loss of mangrove cover due to urban expansion, aquaculture, and infrastructure projects. In areas where mangrove degradation has already occurred, targeted restoration initiatives should be implemented to rebuild natural coastal buffers and improve long-term shoreline resilience.

Furthermore, integrating mangrove conservation into broader climate adaptation frameworks can strengthen disaster risk reduction efforts in cyclone-prone regions. Community-based conservation programs may also play an important role in ensuring sustainable management, as local participation can support monitoring, protection, and replantation efforts.

Overall, strengthening mangrove protection and restoration policies offers a practical, nature-based solution for reducing coastal vulnerability in the face of increasing climate-related risks.

13. Conclusion:

This study examined the relationship between mangrove coverage and coastal sediment stability during storm events, with a particular focus on how variations in mangrove density influence shoreline resilience. The analysis of secondary environmental studies and coastal case examples indicates a consistent pattern in which regions with higher mangrove density experience comparatively lower sediment displacement and more stable shoreline conditions during extreme weather events.

The findings also highlight that mangrove effectiveness is not absolute but varies with the extent and continuity of vegetation cover. Dense and continuous mangrove systems demonstrate stronger resilience under storm conditions, while fragmented or degraded areas remain significantly more vulnerable to erosion and shoreline alteration. This suggests that the structural integrity of mangrove ecosystems plays a critical role in determining their protective capacity.

Overall, the study reinforces the ecological importance of mangroves as natural coastal stabilisers and emphasises their relevance in the context of increasing climate-related risks. Beyond their immediate protective function, mangroves contribute to long-term coastal resilience and environmental stability when maintained as healthy and continuous ecosystems.

Future research could further explore the quantitative relationship between mangrove density and sediment dynamics using field-based measurements and long-term monitoring data, which would provide deeper insight into their performance under varying storm intensities and geographical conditions.

14. Future Scope:

Future research can move beyond static satellite analysis by integrating real-time coastal monitoring systems using high-frequency satellite data and drone-based imaging. Combining machine learning with remote sensing could help predict shoreline changes and mangrove stress responses under different storm intensities. Additionally, developing a mangrove health index that links vegetation density, sediment stability and wave energy reduction could provide a more practical tool for coastal management and climate adaptation planning.

References:

Bootsma, J. I. 0., Borsje, B. W. I., Daphne, V. D. W., & Hulscher, S. J. (2024). Intertidal vegetation altering currents in the Scheldt estuary. *University of Twente Research Information*. https://research.utwente.nl/en/publications/75e4e514-1e50-45e4-9b38-1e841f2154b3

Spalding, M. D., McIvor, A. L., Beck, M. W., Koch, E. W., Möller, I., Reed, D. J., Rubinoff, P., Spencer, T., Tolhurst, T. J., Wamsley, T. V., Van Wesenbeeck, B. K., Wolanski, E., & Woodroffe, C. D. (2013). Coastal ecosystems: a critical element of risk reduction. *Conservation Letters*, *7*(3), 293–301. https://doi.org/10.1111/conl.12074

Alongi, D. M. (2008). Mangrove forests: resilience, protection, and ecological significance in coastal ecosystems. *Estuarine, Coastal and Shelf Science, 76*(1), 1–13. https://doi.org/10.1016/j.ecss.2007.08.017

Kathiresan, K., & Bingham, B. L. (2001). Biology of mangroves and mangrove ecosystems. *Advances in Marine Biology, 40*, 81–251. https://doi.org/10.1016/S0065-2881(01)40003-4

Spalding, M. D., McIvor, A. L., Beck, M. W., Koch, E. W., Möller, I., Reed, D. J., Rubinoff, P., Spencer, T., Tolhurst, T. J., Wamsley, T. V., Van Wesenbeeck, B. K., Wolanski, E., & Woodroffe, C. D. (2013). Coastal ecosystems: a critical element of risk reduction. *Conservation Letters*, *7*(3), 293–301. https://doi.org/10.1111/conl.12074

Bootsma, J., Borsje, B. W., Van Der Wal, D., & Hulscher, S. J. (2025). Do vegetated intertidal areas alter currents on an estuarine scale? *Estuarine Coastal and Shelf Science*, *315*, 109177. https://doi.org/10.1016/j.ecss.2025.109177

Bootsma, J., Borsje, B., Van Der Wal, D., & Hulscher, S. (2023). Decadal intertidal vegetation development in an estuary and its effect on the wave damping capacity. *EGU General Assembly 2023 Abstract*. https://doi.org/10.5194/egusphere-egu23-11683

United Nations Environment Programme. (2014). *The importance of mangroves in coastal protection and climate resilience*. https://www.unep.org

Alcamo, J., Fernandez, N., Leonard, S. A., Peduzzi, P., Singh, A., & Ruth, H. R. R. (2012). 21 Issues for the 21st Century - Results of the UNEP Foresight Process on emerging environmental issues. *Environmental Development*, *2*, 150. https://doi.org/10.1016/j.envdev.2012.03.005

U.S. Geological Survey. (n.d.). Landsat missions overview. https://www.usgs.gov/landsat-missions

Google Earth Engine Team. (n.d.). Google Earth Engine: Planetary-scale geospatial analysis platform. https://earthengine.google.com

European Journal of Remote Sensing. (n.d.). Mangrove vegetation analysis using NDVI and remote sensing techniques. *Taylor & Francis Group*. https://www.tandfonline.com

Elsevier. (n.d.). Coastal ecosystem dynamics and mangrove-based shoreline protection: Remote sensing studies. *ScienceDirect*. https://www.sciencedirect.com

Mumby, P. J., Edwards, A. J., Arias-González, J. E., Lindeman, K. C., Blackwell, P. G., Gall, A., Gorczynska, M. I., Harborne, A. R., Pescod, C. L., Renken, H., Wabnitz, C. C. C., & Llewellyn, G. (2004). Mangroves enhance the biomass of coral reef fish communities in the Caribbean. *Nature*, *427*(6974), 533–536. https://doi.org/10.1038/nature02286

Barbier, E. B., Koch, E. W., Silliman, B. R., Hacker, S. D., Wolanski, E., Primavera, J., Granek, E. F., Polasky, S., Aswani, S., Cramer, L. A., Stoms, D. M., Kennedy, C. J., Bael, D., Kappel, C. V., Perillo, G. M. E., & Reed, D. J. (2008). Coastal Ecosystem-Based Management with Nonlinear Ecological Functions and Values. *Science*, *319*(5861), 321–323. https://doi.org/10.1126/science.1150349

Velastegui-Montoya, A., Montalván-Burbano, N., Carrión-Mero, P., Rivera-Torres, H., Sadeck, L., & Adami, M. (2023). Google Earth Engine: A global analysis and future trends. *Remote Sensing*, *15*(14), 3675. https://doi.org/10.3390/rs15143675

Dasgupta, S., Islam, M. S., Huq, M., Khan, Z. H., & Hasib, M. R. (2019). Quantifying the protective capacity of mangroves from storm surges in coastal Bangladesh. *PLoS ONE*, *14*(3), e0214079. https://doi.org/10.1371/journal.pone.0214079

Spalding, M., McIvor, A., Tonneijck, F. H., Tol, S., & van Eijk, P. (2014). *Mangroves for coastal defence: Guidelines for coastal managers and policy makers*. Wetlands International and The Nature Conservancy. https://www.nature.org

Das, S., & Vincent, J. R. (2009). Mangroves protected villages and reduced death toll during Indian super cyclone. *Proceedings of the National Academy of Sciences*, *106*(18), 7357–7360. https://doi.org/10.1073/pnas.0810440106

Giri, C., Ochieng, E., Tieszen, L. L., Zhu, Z., Singh, A., Loveland, T., Masek, J., & Duke, N. (2010). Status and distribution of mangrove forests of the world using earth observation satellite data. *Global Ecology and Biogeography*, *20*(1), 154–159. https://doi.org/10.1111/j.1466-8238.2010.00584.x

Beebe, M. J., & Phillips-Riggs, L. (1980b). Predictions and inferences in reading. *Elements: Translating Theory Into Practice*, *11*(7), 3–6. https://doi.org/10.1073/pnas.1820067116

Hochard, J. P., Hamilton, S. E., & Barbier, E. B. (2019). Mangroves shelter coastal economic activity from cyclones. *Proceedings of the National Academy of Sciences, 116*(25), 12232–12237. https://doi.org/10.1073/pnas.1820067116

Duarte, C. M., Losada, I. J., Hendriks, I. E., Mazarrasa, I., & Marbà, N. (2013). The role of coastal plant communities for climate change mitigation and adaptation. *Nature Climate Change*, *3*(11), 961–968. https://doi.org/10.1038/nclimate1970

Möller, I., Kudella, M., Rupprecht, F., Spencer, T., Paul, M., Van Wesenbeeck, B. K., Wolters, G., Jensen, K., Bouma, T. J., Miranda-Lange, M., & Schimmels, S. (2014). Wave attenuation over coastal salt marshes under storm surge conditions. *Nature Geoscience*, *7*(10), 727–731. https://doi.org/10.1038/ngeo2251

Das, S., & Vincent, J. R. (2009b). Mangroves protected villages and reduced death toll during Indian super cyclone. *Proceedings of the National Academy of Sciences*, *106*(18), 7357–7360. https://doi.org/10.1073/pnas.0810440106

Alongi, D. M. (2012). Carbon sequestration in mangrove forests. *Carbon Management*, *3*(3), 313–322. https://doi.org/10.4155/cmt.12.20

Hamilton, S. E., & Casey, D. (2016). Creation of a high spatio‐temporal resolution global database of continuous mangrove forest cover for the 21st century (CGMFC‐21). *Global Ecology and Biogeography*, *25*(6), 729–738. https://doi.org/10.1111/geb.12449

Appendix:

Appendix A: Glossary of Key Terms

1) **Mangroves:** Coastal vegetation adapted to saline environments, playing a key role in shoreline stabilization and sediment retention.
2) **Sediment Stability:** The ability of coastal sediments to remain in place without excessive erosion or displacement.
3) **Storm Surge:** An abnormal rise in sea level during cyclonic or storm events.
4) **NDVI (Normalized Difference Vegetation Index):** A remote sensing indicator used to measure vegetation health and density.
5) **Remote Sensing:** The process of obtaining environmental data from satellite or aerial imagery.
6) **Landsat:** A series of Earth-observing satellite missions used for environmental monitoring.
7) **Google Earth Engine (GEE):** A cloud-based platform for analysing geospatial and satellite data.
8) **Coastal Erosion:** The gradual removal or displacement of coastal land due to wave action, storms, currents, or sea-level rise.
9) **Cyclone Impact:** The combined physical effects of cyclonic storms on coastal environments, including erosion, flooding, and vegetation damage.
10) **Ecosystem Resilience:** The ability of an ecosystem, such as mangrove forests, to absorb disturbances and recover while maintaining its core functions and structure.
11) **Shoreline Change:** The alteration of coastline position due to erosion, sediment deposition, and storm activity.
12) **Comparative Analysis:** A research method used in this study to evaluate differences in coastal response across regions with varying mangrove density.
13) **Wave Attenuation:** The reduction of wave energy as it passes through mangrove forests or other coastal vegetation, contributing to shoreline protection.
14) **Hydrodynamics:** The study of water movement, including waves, tides, and currents, and their impact on coastal systems.

15) **Sediment Transport:** The movement of sediment particles by water flow, waves, or storm activity along coastal areas.
16) **Coastal Buffering:** The natural ability of ecosystems like mangroves to reduce the impact of storms and wave energy on inland areas.
17) **Bio-physical Interaction:** The relationship between living organisms (mangroves) and physical processes (waves, sediment, currents) in coastal environments.
18) **Vegetation–Sediment Interaction:** The process by which plant roots and structures influence sediment deposition and stability.
19) **Coastal Protection Mechanism:** Natural systems (like mangroves) that reduce coastal damage by absorbing wave energy and stabilising sediments.
20) **Canopy Density:** The thickness and coverage of mangrove vegetation, influencing wave reduction and sediment trapping efficiency
21) **Storm-Induced Erosion:** The rapid removal of coastal material caused by extreme weather events such as cyclones.

Appendix B: Figures Overview

Figure 1: Diagram illustrating mangrove root systems and their role in wave attenuation and sediment retention.

Figure 2: Comparative representation of shoreline erosion in mangrove-covered and non-mangrove coastal regions.

Figure 3: NDVI-based vegetation density map highlighting variations in mangrove distribution across selected coastal areas.

Figure 4: Satellite-based coastal analysis using Landsat imagery and Google Earth Engine outputs for mangrove density assessment on sunderban region.