SEDIMENT DYNAMICS AND THEIR ROLE IN MANGROVE ECOSYSTEM STABILITY

Research Paper

Prepared for: ECODOME Youth Network

Prepared by: Rounak Gupta

Institution: ECODOME Youth Network

Date: July 28th, 2026

Abstract

Mangrove ecosystems occupy the dynamic boundary between land and sea, where their survival depends strongly on the movement, deposition, erosion, and characteristics of sediment. This paper examines the role of sediment dynamics in maintaining mangrove ecosystem stability, with particular attention to sediment supply, tidal transport, surface-elevation change, root–sediment interactions, shoreline erosion, and resilience to relative sea-level rise. Using a literature-based analytical approach, the study synthesizes established scientific understanding of the feedbacks between mangrove vegetation and coastal geomorphology. Mangrove roots can reduce near-bed water velocity, promote sediment trapping, stabilize substrates, and contribute organic material to soils. At the same time, the stability of mangrove forests depends on sufficient mineral and organic sediment accumulation to maintain suitable surface elevations relative to changing sea levels. Sediment deficits caused by altered river flows, dams, dredging, coastal development, or erosion may weaken this capacity. Conversely, excessive or abrupt sedimentation can bury roots and seedlings or modify hydrological conditions. The paper concludes that sediment dynamics should be treated as a central component

of mangrove conservation and restoration, particularly in coastal areas exposed to sea-level rise and increasing human pressure.

Keywords

Mangroves; sediment dynamics; sediment deposition; coastal erosion; accretion; sea-level rise; ecosystem stability; coastal resilience.

1. Introduction

Mangroves are intertidal forests found along tropical and subtropical coastlines, estuaries, deltas, and sheltered shores. They provide habitat for diverse species, support fisheries, store carbon, improve water quality, and reduce the exposure of coastal communities to waves and erosion. Their ecological importance, however, is closely linked to the physical environment in which they grow.

Sediment is one of the most important components of that environment. Rivers, tides, waves, currents, storms, and biological processes continually transport and redistribute mineral and organic material across mangrove landscapes. The balance between sediment deposition and erosion influences soil elevation, shoreline

position, nutrient availability, seedling establishment, and the ability of mangrove forests to remain within a suitable tidal range.

Mangroves also modify sediment movement. Their trunks, aerial roots, pneumatophores, and below-ground root systems increase hydraulic resistance, slow water locally, trap suspended particles, and stabilize deposited material. This creates a two-way relationship: sediment dynamics influence mangrove development, while mangrove vegetation influences sediment dynamics.

Understanding this feedback is increasingly important as sea-level rise, land subsidence, river regulation, coastal engineering, aquaculture, and urban development alter sediment budgets. This paper therefore examines how sediment supply, deposition, erosion, and sediment characteristics contribute to mangrove ecosystem stability and how disruptions to these processes can increase ecosystem vulnerability.

2. Objectives and Research Question

The principal objective of this study is to evaluate the relationship between sediment dynamics and mangrove ecosystem stability. Specifically, the paper aims to explain the mechanisms of sediment transport and deposition in mangrove environments; examine how mangrove vegetation traps and stabilizes sediment; assess the

consequences of erosion and sediment deficits; explore the relationship between sediment accretion and resilience to relative sea-level rise; and identify implications for mangrove conservation and restoration.

Research Question: How do sediment deposition, erosion, sediment supply, and sediment characteristics influence the long-term stability and resilience of mangrove ecosystems under changing environmental conditions?

3. Methodology

This paper uses a qualitative literature-review approach based on peer-reviewed research and authoritative scientific assessments concerning mangrove geomorphology, sediment transport, coastal erosion, surface-elevation change, and sea-level rise. The analysis focuses on mechanisms and relationships that recur across mangrove environments rather than claiming new field measurements. Evidence is interpreted comparatively to identify conditions under which sediment processes strengthen or weaken ecosystem stability. This approach is appropriate for developing an integrated understanding of a system in which biological and physical processes operate together across multiple spatial and temporal scales.

4. Sediment Dynamics in Mangrove Ecosystems

Sediment dynamics describe the erosion, transport, deposition, resuspension, and consolidation of particles within an environment. In mangrove systems, sediment may originate from river catchments, coastal waters, eroding shorelines, marine sources, and biological production. Its movement is controlled by tidal range, currents, waves, storm events, channel morphology, vegetation density, particle size, and the availability of suspended material.

During tidal flooding, water can carry suspended sediment into a mangrove forest. As flow encounters dense vegetation and complex root structures, local water velocity may decline, encouraging particles to settle. Fine sediments can accumulate on the forest floor, while organic matter from roots, leaves, and other biological material contributes to soil formation. During energetic conditions, however, previously deposited sediment may be resuspended or eroded.

The sediment budget of a mangrove therefore reflects a dynamic balance rather than continuous accumulation. Stability depends not simply on the presence of sediment, but on whether deposition, erosion, compaction, organic production, and subsidence

collectively maintain land elevations and substrate conditions suitable for mangrove growth.

5. Mangrove–Sediment Feedbacks

Mangrove vegetation and sediment processes are strongly interconnected. Above-ground roots and stems create drag that changes water movement, while below-ground roots reinforce soil and can reduce its susceptibility to erosion. Vegetation may promote deposition by increasing the residence time of water and providing physical structures around which sediment accumulates.

As the substrate becomes more stable, conditions may become favorable for further vegetation establishment, creating a positive feedback between plant growth and sediment retention. Yet this relationship is not unlimited. The effect varies according to species, forest structure, tidal setting, wave energy, sediment concentration, and geomorphology. Mangroves should therefore not be treated as universal sediment traps under every environmental condition.

Organic contributions are also important. Root growth and the accumulation of organic matter can contribute to soil volume and elevation. Consequently, both mineral sediment deposition and biological soil-building processes influence the vertical development of mangrove surfaces.

6. Sediment Dynamics and Ecosystem Stability

Ecosystem stability can be understood as the capacity of a mangrove system to maintain its structure and ecological functions while responding to environmental disturbance. Sediment dynamics support this stability in several ways.

First, sediment deposition can help maintain surface elevation within the tidal zone. Mangroves occupy a limited elevation range in relation to tides; if the soil surface becomes too low relative to sea level, flooding duration and frequency may increase beyond tolerable conditions for some species. Second, stable substrates support root anchorage and seedling establishment. Third, sediment can carry nutrients and influence soil chemistry, affecting productivity and vegetation patterns. Finally, sediment accumulation and root reinforcement can contribute to shoreline stability.

Erosion produces the opposite pressure when sediment losses exceed gains. Persistent erosion can expose roots, undermine trees, remove seedlings, shift creek banks, and cause shoreline retreat. Ecosystem stability therefore depends on the balance between processes that build and retain the substrate and processes that remove or lower it.

7. Sea-Level Rise and Sediment Accretion

Relative sea-level rise is a major challenge for mangrove ecosystems because it changes the relationship between the forest surface and tidal water levels. Mangroves may persist where vertical surface-elevation gain can keep pace with relative sea-level rise or where forests can migrate landward into suitable areas.

Sediment availability is central to this response. Mineral deposition during tidal inundation can contribute to vertical accretion, while root production and organic matter can further influence elevation. Where sediment supply is limited, land is subsiding, or coastal development blocks landward migration, vulnerability may increase.

The relationship is complex: a high sediment supply can enhance elevation gain, but sedimentation that is too rapid or hydrologically disruptive may stress vegetation. Effective resilience therefore depends on the rate, timing, composition, and distribution of sediment, together with biological productivity and local geomorphology.

8. Human Pressures on Sediment Budgets

Human activities can alter the amount and movement of sediment reaching mangrove coastlines. Dams and reservoirs may retain river-borne sediment upstream. Dredging, navigation channels, embankments, reclamation, and coastal infrastructure can redirect

flows or modify erosion and deposition patterns. Conversion of mangroves for aquaculture or development removes vegetation that previously influenced hydrodynamics and substrate stability.

These interventions may produce effects far beyond the immediate project site because coastal sediment systems are connected. Management decisions should therefore consider sediment pathways at the scale of estuaries, deltas, and catchments rather than focusing only on individual forest patches.

9. The Sundarbans as a Relevant Example

The Sundarbans, spanning the Ganges–Brahmaputra–Meghna delta in India and Bangladesh, illustrates the close relationship between mangrove ecology and deltaic sediment processes. Tidal channels, river discharge, sediment transport, erosion, accretion, cyclones, salinity, subsidence, and human interventions interact across a highly dynamic landscape.

Different parts of the delta may experience different patterns of erosion and deposition, making broad generalizations inappropriate. The example nevertheless demonstrates an important principle: mangrove conservation cannot be separated from the geomorphological and hydrological processes that create and reshape the land on which the forest exists. Maintaining sediment

connectivity and understanding local elevation change are therefore important elements of long-term management.

10. Conservation and Restoration Implications

Mangrove restoration is more likely to succeed when physical site conditions are addressed before planting. Projects should evaluate tidal connectivity, elevation, wave exposure, sediment supply, erosion risk, and historical changes in shoreline position. Planting seedlings in a site that lacks appropriate hydrology or sediment stability may produce poor survival even when suitable species are selected.

Sediment-aware management can include protecting natural river and tidal connections, avoiding unnecessary disruption of sediment pathways, restoring hydrology where feasible, monitoring erosion and accretion, and allowing space for landward mangrove migration. Repeated measurements of surface elevation and sediment deposition can help managers determine whether restored or natural forests are maintaining their position relative to sea level.

Conservation should also recognize that mangroves are part of larger catchment-to-coast systems. Upstream water and sediment management can influence downstream coastal resilience, making

coordination across sectors and administrative boundaries important.

11. Discussion

The evidence indicates that sediment dynamics are not merely a background physical process; they are a fundamental part of mangrove ecosystem functioning. Mangrove stability emerges from feedbacks among hydrodynamics, sediment availability, vegetation structure, root growth, organic accumulation, and elevation change.

A useful management principle follows from this relationship: increasing mangrove area alone does not guarantee resilience. A forest may remain vulnerable if sediment supply is insufficient, erosion is severe, subsidence is rapid, or tidal connectivity has been disrupted. Similarly, sediment accumulation should not automatically be interpreted as beneficial because excessive burial or altered drainage can negatively affect vegetation.

Future research and monitoring should therefore integrate ecological indicators with measurements of shoreline change, sediment deposition, surface elevation, subsidence, and hydrodynamic conditions. Such integration is particularly important for evaluating mangrove responses to climate change.

12. Conclusion

Sediment dynamics play a central role in the stability, development, and resilience of mangrove ecosystems. Sediment transport and deposition contribute to substrate formation and surface-elevation maintenance, while mangrove roots and vegetation can modify flow, trap particles, reinforce soils, and contribute organic material. Together, these processes create feedbacks that influence shoreline stability, forest development, and responses to environmental change.

When sediment supply is disrupted or erosion exceeds deposition, mangroves may experience root exposure, shoreline retreat, altered inundation, and reduced resilience. Under relative sea-level rise, the capacity of a mangrove surface to gain elevation—and the availability of space for landward migration—becomes especially important.

For this reason, successful mangrove conservation should combine biological protection with sediment and hydrological management. Understanding how water, sediment, vegetation, and elevation interact provides a stronger basis for protecting mangrove ecosystems in a changing climate.

References

Alongi, D. M. (2008). Mangrove forests: Resilience, protection from tsunamis, and responses to global climate change. Estuarine, Coastal and Shelf Science, 76(1), 1–13.

Friess, D. A., Rogers, K., Lovelock, C. E., Krauss, K. W., Hamilton, S. E., Lee, S. Y., Lucas, R., Primavera, J., Rajkaran, A., & Shi, S. (2019). The state of the world's mangrove forests: Past, present, and future. Annual Review of Environment and Resources, 44, 89–115.

Krauss, K. W., McKee, K. L., Lovelock, C. E., Cahoon, D. R., Saintilan, N., Reef, R., & Chen, L. (2014). How mangrove forests adjust to rising sea level. New Phytologist, 202(1), 19–34.

Lovelock, C. E., Cahoon, D. R., Friess, D. A., Guntenspergen, G. R., Krauss, K. W., Reef, R., Rogers, K., Saunders, M. L., Sidik, F., Swales, A., Saintilan, N., Thuyen, L. X., & Triet, T. (2015). The vulnerability of Indo-Pacific mangrove forests to sea-level rise. Nature, 526, 559–563.

McIvor, A., Spencer, T., Möller, I., & Spalding, M. (2012). Storm surge reduction by mangroves. Natural Coastal Protection Series: Report 2. The Nature Conservancy and Wetlands International.

Woodroffe, C. D., Rogers, K., McKee, K. L., Lovelock, C. E., Mendelssohn, I. A., & Saintilan, N. (2016). Mangrove sedimentation

and response to relative sea-level rise. Annual Review of Marine Science, 8, 243–266.