

4.7.63

AI25BTECH11002 - Ayush Sunil Labhade

October 5, 2025

Question:

Prove that in any ($\triangle ABC$),

$$\cos A = \frac{b^2 + c^2 - a^2}{2bc}, \quad (0.1)$$

where (a,b,c) are the magnitudes of the sides opposite to the vertices (A,B,C) respectively.

Solution:

By definition of the side lengths,

$$c = |\mathbf{B} - \mathbf{A}|, \quad (0.2)$$

$$b = |\mathbf{C} - \mathbf{A}|, \quad (0.3)$$

$$a = |\mathbf{C} - \mathbf{B}|. \quad (0.4)$$

The cosine of angle A is given by

$$\cos A = \frac{(\mathbf{B} - \mathbf{A})^T (\mathbf{C} - \mathbf{A})}{|\mathbf{B} - \mathbf{A}| \cdot |\mathbf{C} - \mathbf{A}|} \quad (0.5)$$

$$= \frac{(\mathbf{B} - \mathbf{A})^T (\mathbf{C} - \mathbf{A})}{bc}. \quad (0.6)$$

Now, express a^2 in terms of $\mathbf{B} - \mathbf{A}$ and $\mathbf{C} - \mathbf{A}$. Observe that

$$\mathbf{C} - \mathbf{B} = (\mathbf{C} - \mathbf{A}) - (\mathbf{B} - \mathbf{A}). \quad (0.7)$$

Let $\mathbf{u} = \mathbf{C} - \mathbf{A}$ and $\mathbf{v} = \mathbf{B} - \mathbf{A}$ for brevity. Then,

$$\mathbf{C} - \mathbf{B} = \mathbf{u} - \mathbf{v}. \quad (0.8)$$

Hence,

$$a^2 = |\mathbf{C} - \mathbf{B}|^2 \quad (0.9)$$

$$= (\mathbf{u} - \mathbf{v})^T (\mathbf{u} - \mathbf{v}) \quad (0.10)$$

$$= \mathbf{u}^T \mathbf{u} + \mathbf{v}^T \mathbf{v} - 2\mathbf{u}^T \mathbf{v} \quad (0.11)$$

$$= b^2 + c^2 - 2(\mathbf{C} - \mathbf{A})^T (\mathbf{B} - \mathbf{A}). \quad (0.12)$$

Rearranging,

$$(\mathbf{C} - \mathbf{A})^T (\mathbf{B} - \mathbf{A}) = \frac{b^2 + c^2 - a^2}{2}. \quad (0.13)$$

Substitute into the expression for $\cos A$:


$$\cos A = \frac{(\mathbf{B} - \mathbf{A})^T(\mathbf{C} - \mathbf{A})}{bc} \quad (0.14)$$

$$= \frac{1}{bc} \cdot \frac{b^2 + c^2 - a^2}{2} \quad (0.15)$$

$$= \frac{b^2 + c^2 - a^2}{2bc}. \quad (0.16)$$

Thus, the required identity is proved.

Graph:



Figure