

2.9.18

AI25BTECH11002 - Ayush Sunil Labhade

Question:

Find the volume of a cuboid whose edges are given by $-3\hat{i} + 7\hat{j} + 5\hat{k}$, $-5\hat{i} + 7\hat{j} - 3\hat{k}$ and $7\hat{i} - 5\hat{j} - 3\hat{k}$.

Solution: Given:

Point	Vector
a	$\begin{pmatrix} 3 \\ 7 \\ 5 \end{pmatrix}$
b	$\begin{pmatrix} -5 \\ 7 \\ -3 \end{pmatrix}$
c	$\begin{pmatrix} 7 \\ -5 \\ -3 \end{pmatrix}$

TABLE I: Given data

To find volume we need to compute $[a \ b \ c]$ We will compute by finding the determinant of $[a \ b \ c]$:

$$\mathbf{D} = (a \ b \ c) \quad (1)$$

$$\mathbf{D} = \begin{pmatrix} 3 & -5 & -7 \\ 7 & 7 & -5 \\ 5 & -3 & -3 \end{pmatrix} \quad (2)$$

On computing,

$$\det(\mathbf{D}) = 264 \quad (3)$$

$$\therefore [a \ b \ c] = 264 \quad (4)$$

Thus, the volume is 264.

Graph:

