1

ASSIGNMENT-2

Ojaswa Pandey

Download all python codes from

https://github.com/behappy0604/Summer— Internship—IITH/tree/main/Assignment—2

and latex-tikz codes from

https://github.com/behappy0604/Summer— Internship—IITH/tree/main/Assignment—2

1 Question No. 2.39

Construct a quadrilateral MORE where MO = 6, OR = 4.5, $\angle M = 60^{\circ}$, $\angle O = 105^{\circ}$ and $\angle R = 105^{\circ}$.

2 SOLUTION

1) Let us generalize the given data:

$$\angle M = 60^{\circ} = \theta \tag{2.0.1}$$

$$\angle O = 105^{\circ} = \alpha \tag{2.0.2}$$

$$\angle R = 105^\circ = \gamma \tag{2.0.3}$$

$$\|\mathbf{O} - \mathbf{M}\| = 6 = a,$$
 (2.0.4)

$$\|\mathbf{R} - \mathbf{O}\| = 4.5 = b,$$
 (2.0.5)

$$\mathbf{M} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \mathbf{O} = \begin{pmatrix} 6 \\ 0 \end{pmatrix} \tag{2.0.6}$$

2) Also, Let us assume the other two sides as

$$\|\mathbf{R} - \mathbf{E}\| = c \tag{2.0.7}$$

$$||\mathbf{M} - \mathbf{E}|| = d \tag{2.0.8}$$

$$\theta = \theta_1 + \theta_2 \tag{2.0.9}$$

$$\delta = 180^{\circ} - \alpha = 75^{\circ} \tag{2.0.10}$$

3) Now on calculating, we get

$$\implies \angle E + 270^{\circ} = 360^{\circ}, \qquad (2.0.11)$$

$$\implies \angle E = 90^{\circ} \tag{2.0.12}$$

4) Now taking sum of all the angles given and (2.0.12) we get

$$\angle M + \angle O + \angle R + \angle E = 360^{\circ}$$
 (2.0.13)

So construction of given quadrilateral is possible as sum of all the angles is equal to

360°.

5) Now, using cosine formula in $\triangle MOR$ we can find RM:

$$\implies \|\mathbf{R} - \mathbf{M}\|^2 =$$
$$\|\mathbf{M} - \mathbf{O}\|^2 + \|\mathbf{O} - \mathbf{R}\|^2 - 2 \times \|\mathbf{M} - \mathbf{O}\| \times \|\mathbf{O} - \mathbf{R}\| \cos O$$
(2.0.14)

$$\implies RM = 8.38$$
 (2.0.15)

$$\implies \theta = \arcsin 31.24^{\circ}$$
 (2.0.16)

6) Now in $\triangle MER$, we know

$$\angle M = 28.76^{\circ}, \angle E = 90^{\circ}.$$
 (2.0.17)

We know that sum of the angles of a triangle is 180°

$$\implies \angle M + \angle E + \angle R = 180^{\circ}$$
 (2.0.18)

$$\implies 28.76^{\circ} + 90^{\circ} + \angle R = 180^{\circ}$$
 (2.0.19)

$$\implies \angle R = 61.24^{\circ} \tag{2.0.20}$$

7) Now applying sine law of triangle, we get EM= 7.34=d

Lemma 2.1. Exact co-ordinates of the given R and E can be expresssed as

$$\mathbf{R} = \mathbf{O} + |b| \times \begin{pmatrix} \cos \delta \\ \sin \delta \end{pmatrix} \tag{2.0.21}$$

$$\implies |b| = \frac{\|\mathbf{R} - \mathbf{O}\|}{\|\begin{pmatrix} \cos \delta \\ \sin \delta \end{pmatrix}\|}$$
 (2.0.22)

$$\mathbf{E} = \mathbf{M} + |d| \times \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} \tag{2.0.23}$$

$$\implies |d| = \frac{||\mathbf{E} - \mathbf{M}||}{\|\begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix}\|}$$
 (2.0.24)

Proof. • Calculating the co-ordinates of R:

The vector equation of a line can be given by as,

$$\implies$$
 R = **O** + |b| × $\begin{pmatrix} \cos \delta \\ \sin \delta \end{pmatrix}$ (2.0.25)

$$\implies |b| = \frac{\|\mathbf{R} - \mathbf{O}\|}{\|\begin{pmatrix} \cos 75^{\circ} \\ \sin 75^{\circ} \end{pmatrix}\|}$$
 (2.0.26)

$$\implies |b| = \frac{4.5}{1} \tag{2.0.27}$$

$$\implies b = 4.5 \tag{2.0.28}$$

Putting the values in the above equation we get,

$$\implies$$
 R = $\begin{pmatrix} 6 \\ 0 \end{pmatrix}$ + 4.5 × $\begin{pmatrix} 0.259 \\ 0.965 \end{pmatrix}$ (2.0.29)

$$\implies \mathbf{R} = \begin{pmatrix} 6 \\ 0 \end{pmatrix} + \begin{pmatrix} 1.16 \\ 4.35 \end{pmatrix} \tag{2.0.30}$$

$$\mathbf{R} = \begin{pmatrix} 7.16 \\ 4.35 \end{pmatrix} \tag{2.0.31}$$

• Calculating the co-ordinates of E:

The vector equation of a line can be given by as,

$$\implies \mathbf{E} - \mathbf{M} = d \times \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} \tag{2.0.32}$$

$$\implies |d| = \frac{||\mathbf{E} - \mathbf{M}||}{\|\begin{pmatrix} \cos 60^{\circ} \\ \sin 60^{\circ} \end{pmatrix}\|}$$
 (2.0.33)

$$\implies |d| = \frac{7.34}{1} \tag{2.0.34}$$

$$\implies d = 7.34$$
 (2.0.35)

Putting the values in the above equation we

get,

$$\implies$$
 E = $\begin{pmatrix} 0 \\ 0 \end{pmatrix} + 7.34 \times \begin{pmatrix} 0.50 \\ 0.87 \end{pmatrix}$ (2.0.36)

$$\implies \mathbf{E} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} + \begin{pmatrix} 3.67 \\ 6.36 \end{pmatrix} \tag{2.0.37}$$

$$\mathbf{E} = \begin{pmatrix} 3.67 \\ 6.36 \end{pmatrix} \tag{2.0.38}$$

8) Now, we have the coordinate of vertices M,O,R,E as,

$$\mathbf{M} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \mathbf{O} = \begin{pmatrix} 6 \\ 0 \end{pmatrix}, \mathbf{R} = \begin{pmatrix} 7.16 \\ 4.35 \end{pmatrix}, \mathbf{E} = \begin{pmatrix} 3.67 \\ 6.36 \end{pmatrix}.$$
(2.0.39)

9) On constructing the given quadilateral on python and marking angle we get:




Fig. 2.1: Quadrilateral MORE