

## NCERT Exemplar Solutions

## Class 12 – Mathematics

## Chapter 1 – Relations and Functions

## Objective Type Questions

## Question 1.28:

Let  $T$  be the set of all triangles in the Euclidean plane and let a relation  $R$  on  $T$  be defined as  $aRb$ , if  $a$  is congruent to  $b$ ,  $\forall a, b \in T$ . Then,  $R$  is

- (a) reflexive but not transitive
- (b) transitive but not symmetric
- (c) equivalence
- (d) None of these

## Solution 1.28: (c)

Consider that  $aRb$ , if  $a$  is congruent to  $b$ ,  $\forall a, b \in T$ .

Since, every triangle is congruent to itself.

Hence,  $R$  is reflexive.

...(1)

Let  $aRb \Rightarrow a \cong b$

$\Rightarrow b \cong a \Rightarrow bRa$

$\Rightarrow bRa$

If first triangle is congruent to second triangle, then the second triangle is also congruent to first triangle.

Hence,  $R$  is symmetric.

...(2)

Let  $aRb$  and  $bRc$

$\Rightarrow a \cong b$  and  $b \cong c$

$\Rightarrow a \cong c \Rightarrow aRc$

Hence,  $R$  is transitive.

...(3)

Therefore,  $R$  is equivalence relation.

**Question 1.29:**

Consider the non-empty set consisting of children in a family and a relation  $R$  defined as  $aRb$ , if  $a$  is brother of  $b$ . Then,  $R$  is

- (a) symmetric but not transitive
- (b) transitive but not symmetric
- (c) neither symmetric nor transitive
- (d) both symmetric and transitive

**Solution 1.29: (b)**

We are given that a relation  $R$  defined  $aRb \Rightarrow a$  is brother of  $b$   
 $aRa \Rightarrow a$  is brother of  $a$ , which is not true.

Hence,  $R$  is not reflexive.

$aRb \Rightarrow a$  is brother of  $b$ .

This does not mean  $b$  is also a brother of  $a$  and  $b$  can be a sister of  $a$ .

Hence, it is not symmetric.

$aRb \Rightarrow a$  is brother of  $b$

and  $bRc \Rightarrow b$  is a brother of  $c$ .

So,  $a$  is brother of  $c$ .

Hence,  $R$  is transitive.

**Question 1.30:**

The maximum number of equivalence relations on the set  $A = \{1, 2, 3\}$  is

- (a) 1
- (b) 2
- (c) 3
- (d) 5

**Solution 1.30: (d)**

We are given that,  $A = \{1, 2, 3\}$

The maximum number of equivalence relations on the set  $A = \{1, 2, 3\}$  is 5, which are given below:

$$R_1 = \{(1, 1), (2, 2), (3, 3)\}$$

$$R_2 = \{(1, 1), (2, 2), (3, 3), (1, 2), (2, 1)\}$$

$$R_3 \{(1,1), (2,2), (3,3), (1,3), (3,1)\}$$

$$R_4 = \{(1,1), (2,2), (3,3), (2,3), (3,2)\}$$

$\therefore$  Maximum number of equivalence relation on the set  $a = \{1, 2, 3\} = 5$

**Question 1.31:**

If a relation  $R$  on the set  $\{1, 2, 3\}$  be defined by  $R = \{(1, 2)\}$ , then  $R$  is

- (a) reflexive
- (b) transitive
- (c) symmetric
- (d) None of these

**Solution 1.31: (b)**

We are given a relation  $R$  on the set  $\{1, 2, 3\}$  be defined by  $R = \{(1, 2)\}$

It is clear that  $R$  is transitive.

**Question 1.32:**

Let us define a relation  $R$  in  $R$  as  $aRb$  if  $a \geq b$ . Then,  $R$  is

- (a) an equivalence relation
- (b) reflexive, transitive but not symmetric
- (c) symmetric, transitive but not reflexive
- (d) neither transitive true nor reflexive but symmetric

**Solution 1.32: (b)**

We are given that,  $aRb$  if  $a \geq b$

$$\Rightarrow aRa \Rightarrow a \geq a \text{ which is true.}$$

For relation  $aRb$  to be symmetric, we must have  $a \geq b$  and  $b \geq a$  which can't be possible.

Hence,  $R$  is not symmetric.

For relation  $aRb$  to be transitive, we must have  $aRb$  and  $bRc$

$$\Rightarrow a \geq b \text{ and } b \geq c$$

$$\Rightarrow a \geq c$$

Hence,  $R$  is transitive

**Question 1.33:**

If  $A = \{1, 2, 3\}$  and consider the relation  $R = \{(1,1), (2,2), (3,3), (1,2), (2,3), (1,3)\}$

Then,  $R$  is

- (a) reflexive but not symmetric
- (b) reflexive but not transitive
- (c) symmetric and transitive
- (d) neither symmetric nor transitive

**Solution 1.33: (a)**

We are given that,  $A = \{1, 2, 3\}$  and  $R = \{(1, 1), (2, 2), (3, 3), (1, 2), (2, 3), (1, 3)\}$

Since,  $(1, 1), (2, 2), (3, 3) \in R$

Hence,  $R$  is reflexive

Now,  $(1, 2) \in R$  but  $(2, 1) \notin R$

Hence,  $R$  is not symmetric

Since,  $(1, 2) \in R$  and  $(2, 3) \in R$

Also,  $(1, 3) \in R$

Hence,  $R$  is transitive.

**Question 1.34:**

The identity element for the binary operation  $*$  defined on  $Q - \{0\}$  as  $a * b = \frac{ab}{2} \forall a, b \in Q - \{0\}$  is

- (a) 1
- (b) 0
- (c) 2
- (d) None of these

**Solution 1.34: (c)**

We are given that,  $a * b = \frac{ab}{2}, \forall a, b \in Q - \{0\}$

Let us suppose  $e$  be the identity element for  $*$ .

$$\therefore a * e = \frac{ae}{2}$$

$$\Rightarrow a = \frac{ae}{2}$$

$$\Rightarrow e = 2$$

**Question 1.35:**

If the set  $a$  contains 5 elements and the set  $b$  contains 6 elements, then the number of one-one and onto mappings from  $a$  to  $b$  is

- (a) 720
- (b) 120
- (c) 0
- (d) None of these

**Solution 1.35: (c)**

Since, the number of elements in  $B$  is more than  $A$ .

Hence, there cannot be any one-one and onto mapping from  $A$  to  $B$ .

**Question 1.36:**

If  $a = \{1, 2, 3, \dots, n\}$  and  $b = \{a, b\}$ . Then, the number of surjections from  $a$  into  $b$  is

- (a)  ${}^n P_2$
- (b)  $2^n - 2$
- (c)  $2^n - 1$
- (d) None of these

**Solution 1.36: (d)**

We are given that,  $a = \{1, 2, 3, \dots, n\}$  and  $b = \{a, b\}$ .

If  $a$  and  $b$  are two non-empty finite sets containing  $m$  and  $n$  elements respectively, then the number of surjection from  $a$  into  $b$  is given by

$${}^n C_m \times m!, \text{ if } n \geq m$$

$$0, \text{ if } n < m$$

$$\text{Here, } m = 2$$

$$\text{Hence, the number of surjection from } a \text{ into } b \text{ is } {}^n C_2 \times 2! = \frac{n!}{2!(n-2)!} \times 2!$$

**Question 1.37:**

If  $f: R \rightarrow R$  be defined by  $f(x) = \frac{1}{x}$ ,  $\forall x \in R$  Then,  $f$  is

- (a) one-one
- (b) onto
- (c) bijective

(d)  $f$  is not defined

**Solution 1.37: (d)**

We are given that,  $f(x) = \frac{1}{x}, \forall x \in R$

If  $x = 0$ , then  $f(x) = \infty$  or not defined.

Hence,  $f(x)$  is not defined function.

**Question 1.38:**

If  $f: R \rightarrow R$  be defined by  $f(x) = 3x^2 - 5$  and  $g: R \rightarrow R$  by  $g(x) = \frac{x}{x^2 + 1}$ . Then,  $g \circ f$  is

(a)  $\frac{3x^2 - 5}{9x^4 - 30x^2 + 26}$

(b)  $\frac{3x^2 - 5}{9x^4 - 6x^2 + 26}$

(c)  $\frac{3x^2}{x^4 + 2x^2 - 4}$

(d)  $\frac{3x^2}{9x^4 + 30x^2 - 2}$

**Solution 1.37: (a)**

We are given that,  $f(x) = 3x^2 - 5$  and  $g(x) = \frac{x}{x^2 + 1}$

$$\begin{aligned}
 \text{Now, } g \circ f &= g\{f(x)\} \\
 &= g(3x^2 - 5) \quad [\because f(x) = 3x^2 - 5] \\
 &= \frac{3x^2 - 5}{(3x^2 - 5)^2 + 1} \\
 &= \frac{3x^2 - 5}{9x^4 - 30x^2 + 25 + 1} \\
 &= \frac{3x^2 - 5}{9x^4 - 30x^2 + 26}
 \end{aligned}$$

**Question 1.39:**

Which of the following functions from  $\mathbb{Z}$  into  $\mathbb{Z}$  are bijections?

(a)  $f(x) = x^3$

(b)  $f(x) = x + 2$

(c)  $f(x) = 2x + 1$

(d)  $f(x) = x^2 + 1$

**Solution 1.39: (b)**

Consider, the second option i.e.,  $f(x) = x + 2$

$$\text{Now, } f(x_1) = f(x_2)$$

$$\Rightarrow x_1 + 2 = x_2 + 2$$

$$\Rightarrow x_1 = x_2$$

Hence,  $f(x) = x + 2$  is one-one function.

Now, let us suppose,  $y = x + 2$

$$x = y - 2 \in \mathbb{Z}, \forall y \in \mathbb{Z}$$

Hence,  $f(x)$  is one-one and onto.

**Question 1.40:**

If  $f: \mathbb{R} \rightarrow \mathbb{R}$  be the functions defined by  $f(x) = x^3 + 5$ , then  $f^{-1}(x)$  is

(a)  $(x + 5)^{\frac{1}{3}}$

(b)  $(x - 5)^{\frac{1}{3}}$

(c)  $(5 - x)^{\frac{1}{3}}$

(d)  $5 - x$

**Solution 1.40: (b)**

We are given that,  $f(x) = x^3 + 5$

Let us suppose,  $y = x^3 + 5$

$$\Rightarrow x^3 = y - 5$$

$$\Rightarrow x = (y - 5)^{\frac{1}{3}}$$

$$\Rightarrow f^{-1}(y) = (y - 5)^{\frac{1}{3}} \quad \left[ \begin{array}{l} \because f(x) = y \\ \Rightarrow x = f^{-1}(y) \end{array} \right]$$

$$\Rightarrow f^{-1}(x) = (x - 5)^{\frac{1}{3}}$$

**Question 1.41:**

If  $f: A \rightarrow B$  and  $g: B \rightarrow C$  be the bijective functions, then  $(gof)^{-1}$  is

(a)  $f^{-1}og^{-1}$

(b)  $fog$

(c)  $g^{-1}of^{-1}$

(d)  $gof$

**Solution 1.41: (a)**

We are given that,  $f: A \rightarrow B$  and  $g: B \rightarrow C$  are the bijective functions

$$\therefore (gof)^{-1} = f^{-1}og^{-1}$$

**Question 1.42:**

If  $f: R - \left\{ \frac{3}{5} \right\} \rightarrow R$  be defined by  $f(x) = \frac{3x+2}{5x-3}$ , the

(a)  $f^{-1}(x) = f(x)$

(b)  $(f \circ f)x = -x$

(c)  $g^{-1}of^{-1}$

(d)  $gof$

**Solution 1.42: (d)**

We are given that,  $f(x) = \frac{3x+2}{5x-3}$

Let us suppose,  $y = \frac{3x+2}{5x-3}$



$$\Rightarrow y(5x-3) = 3x+2$$

$$\Rightarrow 5xy - 3y = 3x+2$$

$$\Rightarrow x(3-5y) = -3y-2$$

$$\Rightarrow x = \frac{3y+2}{5y-3}$$

$$\Rightarrow f^{-1}(y) = \frac{3y+2}{5y-3} \quad \left[ \begin{array}{l} \because f(x) = y \\ \Rightarrow x = f^{-1}(y) \end{array} \right]$$

$$\Rightarrow f^{-1}(x) = \frac{3x+2}{5x-3}$$

$$\therefore f^{-1}(x) = f(x)$$

**Question 1.43:**

If:  $[0,1] \rightarrow [0,1]$  be defined by  $\begin{cases} x, & \text{if } x \text{ is rational} \\ 1-x, & \text{if } x \text{ is irrational} \end{cases}$  then  $(f \circ f)x$  is

If  $x$  is irrational

(a) constant

(b)  $1+x$

(c)  $x$

(d) None of these

**Solution 1.43: (c)**

We are given that,  $f : [0,1] \rightarrow [0,1]$  be defined by  $f(x) = \begin{cases} x, & \text{if } x \text{ is rational} \\ 1-x, & \text{if } x \text{ is irrational} \end{cases}$

Now,  $(f \circ f)x = f(f(x))$

$= x$

**Question 1.44:**

If  $f : [2, \infty) \rightarrow \mathbb{R}$  be the function defined by  $f(x) = x^2 - 4x + 5$ , then the range of  $f$  is

(a)  $\mathbb{R}$

(b)  $[1, \infty)$

(c)  $[4, \infty)$

(d)  $[5, \infty)$

**Solution 1.44: (b)**

We are given that,  $f(x) = x^2 - 4x + 5$

Let us suppose,  $y = x^2 - 4x + 5$

$$\Rightarrow y = x^2 - 4x + 4 + 1 = (x-2)^2 + 1$$

$$\Rightarrow (x-2)^2 = y-1$$

$$\Rightarrow x-2 = \sqrt{y-1}$$

$$\Rightarrow x = 2 + \sqrt{y-1}$$

For the function to be real, the term inside the square root must be greater than or equal to zero.

$$\therefore y-1 \geq 0$$

$$\Rightarrow y \geq 1$$

Therefore, the range of  $f$  is  $(1, \infty)$ .

**Question 1.45:**

If  $f : N \rightarrow R$  be the function defined by  $f(x) = \frac{2x-1}{2}$  and  $g : Q \rightarrow R$  be another function defined

by  $g(x) = x + 2$ . Then,  $(g \circ f)\left(\frac{3}{2}\right)$  is

(a) 1

(b) 1

(c)  $\frac{7}{2}$

(d) None of these

**Solution 1.45: (d)**

We are given that,  $f(x) = \frac{2x-1}{2}$  and  $g(x) = x + 2$

Now,  $(g \circ f)(x) = g\{f(x)\}$

$$\begin{aligned}\therefore (g \circ f) \frac{3}{2} &= g \left[ f \left( \frac{3}{2} \right) \right] \\ &= g \left( \frac{2 \times \frac{3}{2} - 1}{2} \right) \\ &= g(1) \\ &= 1 + 2 \\ &= 3\end{aligned}$$

**Question 1.46:**

If  $f : \mathbb{R} \rightarrow \mathbb{R}$  be defined by  $f(x) = \begin{cases} 2x : x > 3 \\ x^2 : 1 < x \leq 3 \\ 3x : x \leq 1 \end{cases}$  Then,  $f(-1) + f(2) + f(4)$  is

- (a) 9
- (b) 14
- (c) 5
- (d) None of these

**Solution 1.46: (a)**

We are given that,  $f(x) = \begin{cases} 2x : x > 3 \\ x^2 : 1 < x \leq 3 \\ 3x : x \leq 1 \end{cases}$

$$\begin{aligned}\text{Now, } f(-1) + f(2) + f(4) &= 3(-1) + (2)^2 + 2 \times 4 \\ &= -3 + 4 + 8 \\ &= 9\end{aligned}$$

**Question 1.47:**

If  $f : \mathbb{R} \rightarrow \mathbb{R}$  be given by  $f(x) = \tan x$ , then  $f^{-1}(1)$  is

- (a)  $\frac{\pi}{4}$

$$(b) \left\{ n\pi + \frac{\pi}{4} : n \in \mathbb{Z} \right\}$$

(c) Does not exist

(d) None of these

### Solution 1.47: (a)

We are given that,  $f(x) = \tan x$

Let us suppose,  $y = \tan x$

$$\Rightarrow x = \tan^{-1} y$$

$$\Rightarrow f^{-1}(y) = \tan^{-1} y \quad \left[ \begin{array}{l} \because f(x) = y \\ \Rightarrow x = f^{-1}(y) \end{array} \right]$$

$$\Rightarrow f^{-1}(x) = \tan^{-1} x$$

$$\Rightarrow f^{-1}(1) = \tan^{-1} 1$$

$$= \tan^{-1} \tan \frac{\pi}{4} = \frac{\pi}{4} \quad \left[ \because \tan \frac{\pi}{4} = 1 \right]$$

Copyright © Jagranjosh.com

All rights reserved. No part or the whole of this eBook may be copied, reproduced, stored in retrieval system or transmitted and/or cited anywhere in any form or by any means (electronic, mechanical, photocopying, recording or otherwise), without the written permission of the copyright owner. If any misconduct comes in knowledge or brought in notice, strict action will be taken.