



ELSEVIER

Contents lists available at ScienceDirect

Best Practice & Research Clinical Rheumatology

journal homepage: www.elsevierhealth.com/berh



1

Current trends in sport injury prevention

Carolyn A. Emery^{a,*}, Kati Pasanen^b

^a Chair Sport Injury Prevention Research Centre, Faculty of Kinesiology, Pediatrics and Community Health Sciences, Cumming School of Medicine, University of Calgary, 2500 University Dr. NW, Calgary, Alberta, T2N1N4, Canada

^b Sport Injury Prevention Research Centre, Faculty of Kinesiology, University of Calgary, 2500 University Dr. NW, Calgary, Alberta, T2N1N4, Canada



A B S T R A C T

Keywords:

Musculoskeletal injury
Athletic injury
Primary prevention
Neuromuscular training
Sporting rules
Sport equipment
Consequences of injury
Osteoarthritis

Participation in sport and recreation has important positive implications for public health across the lifespan; however, the burden of sport-related musculoskeletal injury is significant, with the greatest risk being in youth and young adults. Moving upstream to primary prevention of injury is a public health priority that will have significant implications for reducing the long-term consequences of musculoskeletal injury including early post-traumatic osteoarthritis. The primary targets for the prevention of musculoskeletal injury in sport include neuromuscular training (NMT), rule modification, and equipment recommendations. Currently, there is significant high-quality evidence to support the widespread use of NMT warm up programs in team and youth sport, with an expected significant impact of reducing the risk of musculoskeletal injury by over 35%. Policy disallowing body checking in youth ice hockey has led to a >50% reduction in injuries, and rules limiting contact practice in youth American football has significant potential for injury prevention. There is evidence to support the use of bracing and taping in elite sport to reduce the risk of recurrent ankle sprain injury but not for use to prevent the primary injury, and wrist guards are protective of sprain injuries in snowboarding. Future research examining the maintenance of NMT programs across real-world sport and school settings, optimization of adherence, additional benefit of workload modification, and evaluation of rule changes in other sports is needed.

© 2019 Elsevier Ltd. All rights reserved.

* Corresponding author.

E-mail addresses: caemery@ucalgary.ca (C.A. Emery), kati.pasanen@ucalgary.ca (K. Pasanen).

Why is sport injury prevention important?

Sport and recreation are encouraged as part of a healthy lifestyle across the lifespan and in all populations; however, the sport-related injury burden is significant, and there is a relative paucity of research evaluating injury prevention strategies in all sports and across all ages [1]. Participation and injury rates in youth and young adult sport are the highest; sport is the leading cause of injury in youth [2–5]. An estimated 20% of schoolchildren are absent from school at least one day a year due to sports injuries, and 1 in 3 youth seek medical attention for a sport-related injury annually [1–3]. It is also estimated that 1 in 3 working adults lost at least one day a year from work due to a sport-related injury [1]. Sport accounts for the greatest proportion of all injuries in youth and also has a significant impact in the adult population [1,6]. An Australian study estimates the direct cost of sport-related injury over seven years to be \$265 million Australian dollars [7]. This injury burden highlights the need for evidence-informed injury prevention interventions to reduce risk of injury in youth and across the lifespan. Lower extremity injuries account for over 60% of the overall injury burden in sport, and 60% of these are ankle and knee joint injuries [1,2].

The burden of overweight and obesity in youth has increased over the past 25 years with over 30% of children and adolescents classified as overweight or obese in Canada, and similar trends are found worldwide [8,9]. Sport injury may contribute to this burden with an estimated 8% of youth dropping out of sport annually because of injury or fear of injury [10]. A reduction in future participation in physical activity, which will adversely affect future health including obesity and post-traumatic osteoarthritis. Reducing the public health burden of sport injury in society would have a major impact on quality of life through the promotion of physical activity. There is evidence that low levels of physical activity participation are associated with multiple-cause morbidity and mortality (e.g., cancer, cardiovascular disease, diabetes, and mental illness) [11–13]. Joint injury is also a leading cause of early post-traumatic osteoarthritis (OA), with an estimated 4-fold increased risk of developing OA following knee joint injury [14]. Further, recent evidence indicates that as early as 3–10 years following a sport-related knee joint injury, youth will have more clinical symptoms, lower knee-related quality of life, higher adiposity, weaker knee muscles, poorer dynamic balance, and more structural changes (10-fold greater risk of MRI-defined osteoarthritis) consistent with future OA than uninjured controls [15–17].

Sport and recreation participation is extremely important in youth and for all age groups to maintain a healthy lifestyle, promote healthy growth and development, prevent chronic disease, and reduce stress, among other benefits. An active population is optimal; however, participation in any physical activity must be balanced with the injury risk. As such, the primary focus of this chapter will be lower extremity injury prevention. Participation in sport and recreation has important implications for public health benefits across the lifespan, and mitigating injury risk is paramount.

What is the optimal approach to sport injury prevention?

The four-step model for injury prevention proposed by van Mechelen has been the foundation for developing and evaluating sport injury prevention programs over the last 25 years (Fig. 1) [18]. This model indicates that the prevention of sport injuries begins with establishing the extent of injury in a given population through surveillance systems, followed by identifying risk factors for injury in that population. This leads to the development and validation of injury prevention strategies followed by their evaluation through measuring the impact of the prevention strategy on the incidence of injury using appropriate surveillance systems. Randomized controlled trials (RCT) are the optimal methods to evaluate the efficacy of a prevention strategy but not always feasible or ethical. As such, less rigorous study designs (e.g., quasi-experimental, cohort, and case-control) are also used to evaluate injury prevention intervention efficacy and effectiveness with their inherent methodological limitations that introduce study biases and limit the interpretation of study findings to some extent [19].

The “best” injury prevention program is one that can and will be adopted and sustained by athletes, coaches, and sporting bodies. Therefore, the implementation of injury prevention research into real-world settings has received a great deal of attention [20–22]. Wide-scale implementation of cost-effective intervention measures in real-life circumstances is an ongoing challenge that goes beyond evaluation through an RCT [21]. The Translating Research into the Injury Prevention Practice (TRIPP)




Fig. 1. The four-step model for sport injury prevention as described by van Mechelen [18] (with permission).

Framework has been widely cited to address these challenges [20]. This framework is an extension of the van Mechelen model and describes two additional steps that are required to translate effectiveness of prevention strategies into practice (Fig. 2) [23]. These additional steps involve understanding the real-world context for which an intervention is being developed and evaluating the intervention in a real-world setting. For example, if an injury prevention program is to be transferable and sustainable in the real world, it will be important to consider the sport participants' age group, level of play, sport type, and organizational structure in which the original intervention was evaluated. The RE-AIM framework (Reach Efficacy Adoption Implementation Maintenance Framework) was originally developed to evaluate the public health impact of health promotion interventions and describes five cross-cutting dimensions that identify the translatability and feasibility of a program [22]. The five steps to translate research into action described include 1) reach the target population, 2) effectiveness or efficacy, 3) adoption by target staff, settings, or institutions, 4) implementation consistency and costs and adaptations made during delivery, and 5) maintenance of intervention effects in individuals and settings over time [24]. Increasing attention has been paid to this framework in the context of sport injury prevention strategy evaluation, and this will continue to be an area of further development in the field [25–27].

The importance of the economic evaluation of injury prevention research has more recently been identified, adding relevant insights into the financial consequences and outcomes of preventive approaches. This information will inform practice and policy related to injury prevention strategies. An efficient use of limited financial resources is imperative, yet only a handful of full economic evaluations in the field of sport medicine have been published [28–31].

When developing an optimal implementation strategy that will maximize the effectiveness of a specific sport injury prevention program, it is critical to consider the multiple factors that may influence adherence to such an approach across many levels of influence (i.e., athlete/child, parent, coach/teacher/trainer, sport organization, and government) [32]. To mitigate the problems of poor adherence, it is important to ensure a balance between evidence and real-world user implementation throughout intervention development and evaluation. Owwoye et al. [33] recommend a focus on the identification




Fig. 2. Translating research into injury prevention practice (TRIPP) as described by Finch [22] (reproduced with permission).

and modification of determinants of adherence for successful implementation and maintenance using a four-step model (Fig. 3).

What is the evidence supporting the use of sport injury prevention programs?

While injuries are arguably predictable and preventable, it is likely impossible to eliminate all injuries in sport; however, injury prevention strategies can reduce the number and severity. On the basis of relative burden, the focus of much of the evidence surrounding musculoskeletal injury prevention in sport has been on reducing the risk of lower extremity injuries. In the past decade, there have been an increasing number of epidemiological studies evaluating the efficacy of prevention strategies in youth and elite sport to prevent lower extremity musculoskeletal injuries. These evidence-informed strategies cross three themes, namely, 1) training strategies, 2) sport rule modifications and policy changes, and 3) equipment recommendations. A diversity of sport-specific (i.e., primarily, team sports) and more general sport populations (e.g., school based) have been targeted for injury prevention strategy evaluation. The greatest number of these strategies is training strategies targeting modifiable intrinsic (athlete related) risk factors (e.g. strength, endurance, and balance) through exercise interventions, primarily neuromuscular in nature. In addition, extrinsic risk (environmental) factors have been addressed through rule modification and equipment strategies in some sports (e.g., body checking in youth ice hockey, noncontact practice in American football, wrist guards for snowboarding, and ankle braces).

Neuromuscular training injury prevention programs

Until the past decade, there were relatively few scientifically rigorous evaluation studies examining the efficacy of injury prevention strategies in sport [34]. Epidemiological research historically focused on the evaluation of prevention strategies in elite adult amateur and professional athlete populations




Fig. 3. The four key steps toward more rigorous approach to promote adherence to interventions as described by Owoeye et al. [33].

where injury surveillance was established with the presence of medical staff, and more recently, there has been a greater focus on youth sport [35]. The majority of prospective studies (e.g., RCT, quasi-experimental, and cohort studies) evaluate neuromuscular training (NMT) strategies in team sport, with increasing attention toward youth sport. NMT programs are typically coach or trainer led, ideally following a comprehensive coach training workshop by a physiotherapist or strength and conditioning coach with expertise in NMT workshop delivery.

Several systematic reviews, some including meta-analyses where data are combined across multiple studies, have been conducted examining studies that have evaluated the effectiveness of NMT strategies in reducing the risk of injury in sport [34–42]. Relevant meta-analyses produce combined estimates of measure of effect using incidence rate ratios (IRR) examining NMT interventions across multiple sport contexts and report consistent findings [35–37]. Examining RCTs only in youth sport (i.e., soccer, European handball, and basketball), seven studies examining overall lower extremity injury outcome were combined to estimate a 35% reduction in lower extremity injury risk [IRR = 0.65 (95% CI; 0.49–0.86)] and five studies examining knee injury outcome specifically estimated a 26% reduction in knee injury risk [IRR = 0.74 (95% CI; 0.51–1.07)] that was clinically relevant but not statistically significant. In another systematic review, meta-analyses of 25 RCTs across a variety of sports (i.e., soccer, European handball, Australian rules football, floorball, and basketball) and age groups examined the protective effect of exercise interventions in reducing the risk of injury [36]. The combined estimate demonstrated a 37% reduction in overall injury risk [IRR = 0.63 (95% CI; 0.53–0.75)], 35% reduction in acute injury risk [IRR = 0.65 (95% CI; 0.5–0.84)], and 47% reduction in overuse injury risk [IRR = 0.53 (95% CI; 0.37–0.75)] [34]. Further, an even greater preventive effect was seen when NMT intervention programs focused on proprioception/balance [IRR = 0.55 (95% CI; 0.35–0.87)] and strength [IRR = 0.32 (95% CI; 0.21–0.48)] while no preventive effect was demonstrated with programs focused on stretching [IRR = 0.96 (95% CI; 0.85–1.1)]. Another study combined RCTs examining the effectiveness of NMT in reducing the risk of ankle sprains specifically and reported a 32% reduction in ankle sprain injury risk in youth basketball and soccer players [IRR = 0.68 (95% CI; 0.46–0.99)] [42]. Further, there is recent evidence that a NMT warm up is effective in reducing injury risk in pediatric soccer players under the age of 13 years by 48% [Hazard Ratio = 0.52 (95% CI; 0.32–0.86)] [43]. More recently, a protective effect of NMT training was found in youth rugby players when exercises were performed at least three times per week [IRR = 0.28 (95% CI; 0.14–0.51)] [44]. Further, to a sport-specific context, there is building evidence regarding the effectiveness of such NMT warm up programs in reducing the overall risk of sport and recreational injury in a school physical education context across the spectrum of age groups of children and adolescents [45–48].

NMT training programs, typically consist of exercises aimed to improve balance, strength, and agility (e.g., coordination, cutting and landing techniques) [41]. NMT programs are often introduced as part of an extended warm up routine, and as such, NMT components include aerobic, balance, strength, and agility (see Fig. 4). Ideally, levels of progression are built into each NMT exercise component such that participants can progress difficulty as they see fit. Examples of NMT components are provided in Fig. 4. To successfully reduce the burden of sport-related injury, it is critical to develop and implement feasible NMT programs that fit in a real-world sporting context (e.g., part of the warm up routine, no required equipment). Further research is required to fully understand the mechanism by which

AEROBIC

Players run from one side of the gym to the other (15–20m), using arms and a relaxed running technique. Focus on good posture.

1. Forward run

1–2 laps

**2. Forward run with backward zig-zag variations**

1–2 laps

**3. Skipping**

1–2 laps



Level 1
Forward-backward skipping



Level 2
Sideways skipping

Level 1
Backward zig-zag shuffles

Level 2
Backward zig-zag carioca

4. Forward run progressing speed

1–2 laps

**AGILITY**

Ensure soft landing on toes. Focus on good posture, and knee-over-toe position.

5. One-leg jumps over a line

10–15 reps per leg




Level 1
Forward-Backward



Level 2
Side-to-side

6. Jumps in place

8–12 reps



Level 1
Squat jumps



Level 2
Skate jumps

STRENGTH

Maintain body in a straight line from head to toes.

7. Plank

20–30 seconds



Level 1
On elbows



Level 2
With leg lifts

8. Side Plank

20–30 seconds per side



Level 1
On elbows



Level 2
With leg or arm lifts

9. Hamstrings

Maintain a straight line from head to knees, slowly lower the body toward the mat.

Level 1 3–6 reps

Level 2 7–10 reps

**10. Walking Lunges**

5–10 metres



Level 1
With torso rotation



Level 2
With knee lift, forward-backward

11. Side Lunges

Level 1
3-D
2–4 reps in each direction



Level 2
With arm circles
8–12 reps

**BALANCE**

Players stand facing each other while balancing on one leg with hip and knee slightly bent and knee aligned over the planted foot. The other knee is slightly bent. Focus on good posture and knee-over-toe position.

12. Single-leg balance

4–6 reps per leg



Level 1
With torso rotation



Level 2
With ball roll

13. Single-leg balance

4–6 reps per leg



Level 1
With ball toss



Level 2
With jump catch

different components of NMT reduce injury risk and optimization of adherence (i.e., utilization frequency, utilization fidelity, duration fidelity, exercise fidelity, and cumulative utilization) to NMT programs across sport contexts [33]. There is ample evidence to support a dose–response relationship of greater exposure to NMT leading to a greater reduction in injury risk across sport contexts [44,49–53]. To assess the effectiveness of differing dosages of the NMT programs, adherence needs to be measured and quantified. Assessing exercise fidelity, or the extent to which a program is followed as prescribed (i.e., correct execution of NMT warm up program), is an area that has been under-researched and not often considered when measuring adherence [54]. As such, it is not known how movement quality during the NMT warm up influences injury rates and the importance of movement quality in performing NMT components may be underestimated. As one can imagine, there are challenges in analyzing movement quality in a real-world sporting context setting where qualified personnel may observe and evaluate movement patterns, which could be influenced by observer bias and is a time-consuming process [54]. The use of traditional biomechanical motion capture equipment to quantify movement patterns is expensive and requires trained individuals to operate the equipment, and this type of analysis is often confined to a laboratory or clinic [55]. Recent advancements in wearable technologies have provided the means to measure movement quantity and quality outside of a biomechanics laboratory [56]. The future development of wearable technologies to determine movement quality during an NMT warm up program to provide information about how exercise fidelity during an NMT warm up program influences injury rates is promising.

Further to NMT programs, Gabbett 2016 highlights the potential for load modification in reducing the risk of injuries in sport [57]. He proposes the ‘Training-Injury Prevention Paradox’ whereby athletes accustomed to high training loads may have a lower risk of injury than athletes training at lower workloads [57]. He argues that excessive and rapid increases in training load may increase the risk of noncontact musculoskeletal injuries (e.g., tendinopathies) and that the acute-to-chronic training workload ratio may predict training-related injuries [56]. With greater availability of wearable technologies to monitor workload (e.g., jump load in volleyball and basketball), there exists a greater opportunity to better understand the influence of workload on injury risk [58,59]. This research provides a foundation for the development and evaluation of future workload modification interventions to reduce the risk of such sport-related injuries.

Research evaluating training strategies aimed at reducing the risk of upper extremity injuries is far more limited. A Shoulder Injury Prevention Programme aimed at throwing athletes to increase glenohumeral internal rotation, external rotation, and scapular muscle strength and improving kinetic chain and thoracic mobility was implemented as part of a European handball warm up program in elite players and evaluated in a cluster-RCT study, reporting a 28% reduction in shoulder problems (Odds Ratio = 0.72, 95% CI; 0.52–0.98) [60]. A systematic review examining risk factors and shoulder injury prevention programs for throwing athletes summarizes the limitations in evaluating injury prevention training programs in overhead athletes due to the limited number of athletes uninjured at baseline and limited statistical power as a result of examining new injury outcomes [61].

In summary, there is evidence from adult elite, adolescent, and pediatric athletes and sport participants to inform best practice in NMT warm up program recommendations to prevent injuries across multiple team sport contexts including soccer, European handball, Australian rules football, basketball, floorball, and rugby. Translation of these findings and adaptation of sport-specific NMT programs to a more generalized population such as physical education students across child and adolescent age groups has also proven to be effective in reducing the risk of injuries in sport and recreation. It stands to reason that the development and implementation of sport-specific NMT warm up programs in sports where evaluation has not been previously done will also be effective; however, future research is necessary. Implementation and evaluation of such programs in a real-world context are necessary to gain a further understanding of adherence and maintenance of such programs across sport contexts. Research examining exercise fidelity, quality of movement, and workload also requires more attention,

Fig. 4. Examples of neuromuscular training warm up program components; 4a Examples of strength and balance components and 4b Neuromuscular training warm up: Examples of aerobic and agility components [acknowledgment Sport Injury Prevention Research Centre, University of Calgary, K Pasanen, with permission].

with promising opportunities using wearable technologies. Further, future research evaluating upper extremity injuries should target additional throwing sport populations.

Sporting rule modification and policy changes

Sporting rules and policy are important for the regulation of the sport, but some have been implemented specifically to reduce the risk of injury. An example of evidence-informed policy change related to the age of introduction of body checking in youth ice hockey. Body checking is a tactic used to gain competitive advantage by changing direction or leaving the established skating lane to make contact with the body of the opponent, or using hips, shoulders, or arms to push off and separate the opponent from the puck [62]. The optimal age of introduction to body checking has been a topic of debate in North America for over a decade and several studies have contributed to evidence-informed body checking policy change. In a meta-analysis, policy-allowing body checking in youth increased injury risk more than 2-fold (combined Risk Ratio = 2.45; 95% CI 1.7 to 3.6) [63]. Supporting this, evaluation of a national evidence-informed policy change in Canada disallowing body checking at all levels of play in 11- and 12-year-old leagues led to a 50% reduction in injury risk [IRR = 0.5 (95% CI; 0.33–0.75)] [64]. This evidence subsequently informed policy disallowing body checking in older age groups (ages 13–14 years) in non-elite leagues (lowest 60–70% by division of play), which also reduced the risk of injury 54% [IRR = 0.46 (95% CI; 0.27–0.77)] [65].

The lessons learned over a decade of research examining body checking policy change in youth ice hockey may have implications for other team sports such as American football, rugby, lacrosse, and soccer. Other studies evaluating rule changes in youth team sport report an association of a high school football policy restricting teams to no more than two collision practices a week with fewer head contacts in games and practices and that, combined with coach safety training, is associated with 63% lower rates of injury in practices than those with coach training alone [IRR = 0.37 (95% CI, 0.26–0.53)] [66]. Further research is required to determine whether such a strategy will have implications for the long-term reduction of injury risk in games. While there are limited studies examining the impact of rule and contact policy changes in sport, these findings have arguably exceeded the public health impact of other injury prevention approaches such as training strategies in some sports such as ice hockey.

Equipment recommendations

Using protective equipment may help prevent musculoskeletal injury and/or severity in sport. Such equipment may include bracing/taping and wrist guards. In a systematic review conducted in adult and youth athletes, Dizon et al. [67] report in a systematic review a protective effect of ankle bracing and taping among previously injured adult and youth athletes in reducing the risk of ankle sprain re-injury 69% and 71%, respectively. Janssen et al. [68] demonstrate a superior effect of ankle bracing over NMT training in reducing the incidence of recurrent ankle sprain 47% compared to the NMT alone, with similar reported full compliance in the bracing (48%) and NMT (45%) groups. Interestingly, when the interventions were combined, full compliance reduced to 28%, with no significant benefit to the combined approach over NMT alone [66]. The evidence does not support the preventative effect of ankle bracing/taping or knee bracing as a primary strategy to prevent ankle and knee injuries in youth sport, except in the case of recurrent injury [69]. Further, recommendations for bracing compared to taping may be based on personal preference, comfort, availability of medical personnel for taping, and potential cost implications of each [70].

Other equipment strategies examined include wrist guard use in snowboarding [71]. A systematic review examining wrist guards in snowboarding reveals a significant protective effect in reducing the risk of wrist injury (Odds Ratio = 0.23), wrist fracture [Risk Ratio (RR) = 0.29], and wrist sprains (RR = 0.17) [71]. It is estimated that for every 50 snowboarders who wear wrist guards, one wrist injury will be prevented [71]. This research has potential implications for other sports including skateboarding and roller blading, and further research is required to translate these findings to other contexts. Breakaway bases have been examined in baseball and softball and have consistently demonstrated a protective effect on sliding injuries in adult and youth population [72–74]. While there

is some evidence to support the use of protective equipment in youth sport, there is also evidence to support less than optimal uptake of equipment strategies (e.g., wrist guard use in snowboarding) [71]. Combining educational approaches (e.g., social media) and policy (legislation and facility/sport association requirements) may be the best way to encourage use of protective equipment.

The role of sport injury prevention in reducing the global burden of MSK disorders

Moving upstream to primary prevention of injury in a healthy and active population may ultimately have the greatest public health impact in reducing the burden of injuries and their consequences, including early post-traumatic OA. The significant consequences of musculoskeletal injury in contributing to reduced levels of physical activity, overweight/obesity, OA, and health care costs inform the need for rigorous evaluation of primary injury prevention strategies in sport and recreation to reduce this burden. There is significant high-quality evidence to support the widespread use of NMT warm up programs in team sports and youth sport widely, with an expected significant impact in reducing the risk of musculoskeletal injury by over 35%. Future research examining the maintenance of NMT programs across real-world sport and school settings, optimization of adherence and exercise fidelity, and additional benefit of workload modification is critical. Sport-specific rule modifications and contact policy changes in youth collision sport can have significant public health impact in the reduction of injuries in youth sport. Policy disallowing body checking in some age groups and levels of play in youth ice hockey has led to a >50% reduction in injuries, and rules limiting contact practice in youth American football has significant potential for best practice in injury prevention. Future research evaluating such rule changes in other collision sports is needed. There is evidence to support the use of bracing and taping in elite sport to reduce the risk of recurrent ankle sprain injury but not for use as a primary prevention practice. Further, wrist guards are protective of wrist sprain injuries in snowboarding. Participation in sport and recreation has important implications for public health benefits across the lifespan, and mitigating injury risk is paramount.

The need for future research in the primary prevention should not, however, be underestimated, including the need for methodologically rigorous evaluation studies including RCTs in sports where research has not been undertaken. The Osteoarthritis Research Society International (OARSI) RCT recommendations were developed to inform the design, conduct, and analytical approaches to RCTs evaluating the preventative effect of primary musculoskeletal injury prevention strategies and strategies aimed at the secondary prevention of early post-traumatic OA to inform the highest level of evidence in the field [75]. Further, implementation research in the real-world context including mixed methods studies to inform barriers and facilitators to injury prevention best practice and optimization of positive behavior change in sport including implementation of NMT warm up programs broadly to have a greater public health impact in reducing the burden of musculoskeletal injuries [27]. RCTs may not always be feasible or ethical to evaluate potential training strategies, rule changes, and equipment recommendations in sport, and therefore, case–control studies (e.g., equipment), cohort studies (e.g., rule changes), quasi-experimental designs, and comparative effectiveness studies (e.g., training strategies) may be considered. Economic evaluation of injury prevention strategies is critical to inform cost-savings associated with injury prevention programs. Results from such economic evaluations will support the injury prevention community in advocating for best practice and policy with government and sporting association stakeholders who have the potential to influence a significant reduction in the burden of musculoskeletal injury in sport and recreation [28–31].

Primary prevention of injury may be the “low hanging fruit” that will have the greatest impact in reducing the burden of musculoskeletal injury in sport and recreation. Once injury has occurred, it is imperative that those individuals who are at high risk of recurrent injury and/or risk of significant consequences of injury such as early post-traumatic OA are identified, and secondary prevention strategies also considered. These recommendations are beyond the scope of this chapter and will be addressed in chapter 10. An improved evidence-base in the primary and secondary prevention of musculoskeletal injury in sport and recreation will inform health care practitioners, sport and health administrators, legislators, athletes, coaches, parents, and the public to make the best possible evidence-informed decisions regarding the prevention of injury in the broad and diverse communities in which physical activity is promoted through sport and recreation.

Practice points

- Neuromuscular training warm up programs including strength, balance, aerobic, and agility components can reduce the risk of musculoskeletal injuries at least 35% in team sport and other youth sport settings
- Sport rule modifications and contact policy changes in youth collision sport may have significant public health implications in reducing the risk of injury, and their potential should not be underestimated
- Ankle bracing will reduce the risk of recurrent ankle sprain injuries in elite sport but not considered as a strategy for primary prevention in healthy athletic populations with no previous history of ankle injury
- Wrist guards are protective of injury in snowboarding and hence should be promoted
- Workload modification interventions require evaluation to further inform the prevention of acute and chronic injuries in elite sport

Research agenda

- Research examining the maintenance of NMT programs across real-world sport and school settings, optimization of adherence, and additional benefit of workload modification is needed.
- Novel outcome measurements using wearable technologies may inform research examining exercise fidelity and workload modification.
- Research evaluating rule modifications and contact policy in youth collision sports is recommended
- The effectiveness of wrist guards in sports beyond snowboarding should be evaluated.
- In the upscaling and broad-scale implementation and evaluation of injury prevention implementation strategies in real-world settings; attention to adherence, exercise fidelity, behavior change, barriers, and facilitators, and long-term maintenance is recommended

Conflicts of interest

Carolyn Emery and Kati Pasanen have no conflicts of interest to declare.

Funding statement

Carolyn Emery is the Chair of Sport Injury Prevention Research Centre, Faculty of Kinesiology, University of Calgary, an International Research Centre for Injury Prevention in Sport and Protection of Athlete Health supported by the International Olympic Committee. Carolyn holds a Chair in Pediatric Rehabilitation supported by the Alberta Children's Hospital Foundation.

We obtained written consent for the use of [Fig. 4](#) in our chapter.

References

- [1] Conn JM, Annett JL, Gilchrist J. Sports and recreation related injury episodes in the US population. *Inj Prev* 2003;9:117–23.
- [2] Emery CA, Tyreman H. Sport participation, sport injury, risk factors and sport safety practices in Calgary and area junior high Schools. *J Paediatr Child Health* 2009;14:439–44.
- [3] Emery CA, Meeuwisse WH, McAllister JR. A survey of sport participation, sport injury and sport safety practices in adolescents. *Clin J Sport Med* 2006;16:20–6.
- [4] Emery CA. Injury prevention and future research. In: Caine DJ, Maffulli N, editors. *Epidemiology of pediatric sports injuries: individual sports*. Basel, Switzerland: Karger; 2005. p. 179–200.
- [5] King MA, Pickett W, King AJC. Injury in Canadian youth: a secondary analysis of the 1993–94 health behaviour in school-aged children survey. *Can J Public Health* 1998;89:397–401.
- [6] Emery C. Risk factors for injury in child and adolescent sport: a systematic review. *Clin J Sport Med* 2003;13:256–68.

- [7] Finch CF, Kemp JL, Clapperton AJ. The incidence and burden of hospital-treated sports related injury in people aged 15+ years in Victoria, Australia, 2004–2010: a future epidemic of osteoarthritis? *Osteoarthritis Cartilage* 2015;23:1138–43.
- [8] Roberts KC, Shields M, Groh MD, Aziz A, Gilbert J. Overweight and obesity in children and adolescents: results from the 2009 to 2011 Canadian health measures survey. *Health Rep* 2012;23:1–5.
- [9] de Onis M, Onyango AW, Borghi E, Siyam A, Nishida C, Siekmann J. Development of a WHO growth reference for school-aged children and adolescents. *Bull World Health Organ* 2007;85:660–7.
- [10] Grimmer KA, Jones D, Williams J. Prevalence of adolescent injury from recreational exercise: an Australian perspective. *J Adolesc Health* 2000;27:1–6.
- [11] Blair SN, Kohl HW, Barlow CE. Changes in physical fitness and all-cause mortality: a prospective study of healthy and unhealthy men. *JAMA* 1995;273:1093–8.
- [12] Paffenbarger RS, Kamput JB, Lee IM. Changes in physical fitness and other lifestyle patterns influence longevity. *Med Sci Sports Exerc* 1994;26:857–65.
- [13] Jebb S, Moore M. Contribution of a sedentary lifestyle and inactivity to the aetiology of overweight and obesity: current evidence and research issues. *Med Sci Sports Exerc* 1999;31:S534–41.
- [14] Richmond S, Fukuchi R, Ezzat A, Schneider KJ, Schneider GM, Emery CA. Are joint injury, sport activity, physical activity, obesity, or occupational activities predictors for osteoarthritis? A systematic review. *J Orthop Sports Phys Ther* 2013;43(8):515–24.
- [15] Whittaker J, Toomey CM, Nettel-Aguirre A, Jaremko JL, Doyle-Baker PH, Woodhouse LJ, et al. Health-related outcomes following a youth sport-related knee injury. *Med Sci Sports Exerc* 2018. <https://doi.org/10.1249/MSS.0000000000001787>. epub ahead of print.
- [16] Toomey CM, Whittaker JL, Nettel-Aguirre A, Reimer RA, Woodhouse LJ, Ghali B, et al. Higher fat mass is associated with a history of knee injury in youth sport. *J Orthop Sports Phys Ther* 2017;47(2):80–7.
- [17] Whittaker JL, Toomey C, Nettel-Aguirre A, Jaremko JL, Doyle-Baker PH, Woodhouse LJ, et al. Health-related outcomes following a youth sport-related knee injury. *Med Sci Sports Exerc* 2018. <https://doi.org/10.1249/MSS.0000000000001787>.
- [18] van Mechelen W, Hlobil H, Kemper HC. Incidence, severity, aetiology and prevention of sports injuries. A review of concepts. *Sports Med* 1992;14:82–99.
- [19] Emery CA. In: Verhagen E, van Mechelen W, editors. Research designs for evaluation studies in research methods in sport medicine. Oxford University Press; 2010. p. 169–80.
- [20] Verhagen E, van Stralen M, Van Mechelen W. Behaviour, the key factor for sports injury prevention. *Sports Med* 2010;40:899–906.
- [21] Timpka T, Ekstrand J, Svanström L. From sports injury prevention to safety promotion in sports. *Sports Med* 2006;36:733–45.
- [22] Finch C. A new framework for research leading to sports injury prevention. *J Sci Med Sport* 2006;9:3–91.
- [23] Verhagen E, Finch CF, Voogt N, Bruinsma A. A knowledge transfer scheme to bridge the gap between science and practice: an integration of existing research frameworks into a tool for practice. *Br J Sports Med* 2013;48:698–701.
- [24] Glasgow RE, Vogt TM, Boles SM. Evaluating the public health impact of health promotion interventions: the RE-AIM framework. *Am J Public Health* 1999;89:1322–7.
- [25] O'Brien J, Finch C. The implementation of injury prevention exercise programmes in team ball sport: a systematic review employing the RE-AIM framework. *Sports Med* 2014;44:1305–18.
- [26] O'Brien J, Finch C. A systematic review of core implementation components in team ball sport injury prevention trials. *Inj Prev* 2014;20:357–62.
- [27] Finch C, Donaldson A. A sports setting matrix for understanding the implementation context for community sport. *Br J Sports Med* 2010;44:973–8.
- [28] Hupperets M, Verhagen E, Heymans M, Bosmans JE, van Tulder MW, van Mechelen W. Potential savings of a program to prevent ankle sprain recurrence: economic evaluation of a randomized controlled trial. *Am J Sports Med* 2010;38:2194–200.
- [29] Krist MR, van Beijsterveldt AMC, de Wit GA, Backx FJG. Preventive exercises reduced injury-related costs among adult male amateur soccer players: a cluster-randomised trial. *J Physiother* 2013;59:15–23.
- [30] Lacny SL, Marshall DA, Currie G, Kulin NA, Meeuwisse WH, Kang J, et al. Reality check: the cost-effectiveness of removing body checking from youth ice hockey. *Br J Sports Med* 2014;48:1299–305.
- [31] Marshall D, Lopatina E, Lacny S, Emery CA. Economic impact of a neuromuscular training prevention strategy to reduce injuries in youth soccer. *Br J Sports Med* 2016;50:1388–93.
- [32] Keats M, Emery CA, Finch C. Are we having fun yet? Fostering adherence to injury preventive exercise recommendations in young athletes. *Sports Med* 2012;42:175–84.
- [33] Owoeye OBA, McKay CD, Verhagen EALM, Emery CA. Advancing adherence research in sport injury prevention. *Br J Sports Med* 2017. <https://doi.org/10.1136/bjsports-2017-098272>. epub ahead of print.
- [34] Emery CA. Injury prevention and future research. In: Caine DJ, Maffulli N, editors. *Epidemiology of pediatric sports injuries: team sports*. Basel, Switzerland: Karger; 2005. p. 170–91.
- [35] Emery CA, Roy TO, Whittaker JL, Nettel-Aguirre A, Mechelen W. Neuromuscular training injury prevention strategies in youth sport: a systematic review and meta-analysis. *Br J Sports Med* 2015;49:865–70.
- [36] Laursen JB, Bertelsen DM, Andersen LB. The effectiveness of exercise interventions to prevent sports injuries: a systematic review and meta-analysis of randomised controlled trials. *Br J Sports Med* 2014;48:871–7.
- [37] Herman K, Barton C, Malliaras P, Morrissey D. The effectiveness of neuromuscular warm-up strategies that require no additional equipment, for preventing lower limb injuries during sport participation: a systematic review. *BMC Med* 2012;10:1–12.
- [38] Abernethy L, Bleakley C. Strategies to prevent injury in adolescent sport: a systematic review. *Br J Sports Med* 2007;41:627–38.
- [39] Schiff MA, Caine DJ, O'Halloran R. Injury prevention in sports. *Am J Lifestyle Manag* 2010;4:42–64.
- [40] Caine D, Maffulli N, Caine C. Epidemiology of injury in child and adolescent sports: injury rates, risk factors and prevention. *Clin Sports Med* 2008;27:19–56.

- [41] Hübscher M, Zech A, Pfeifer K, Hänsel F, Vogt L, Banzer W. Neuromuscular training for sports injury prevention: a systematic review. *Med Sci Sports Exerc* 2010;42:413–21.
- [42] Owoseye OBA, Palacios-Derflingher LM, Emery CA. Prevention of ankle sprain injuries in youth soccer and basketball: effectiveness of a neuromuscular training program and examining risk factors. *Clin J Sport Med* 2018;28:325–31.
- [43] Rössler R, Junge A, Bizzini M, Verhagen E, Chomiak J, aus der Fünften K, et al. A multinational cluster randomised controlled trial to assess the efficacy of '11+ Kids': a warm-up programme to prevent injuries in children's football. *Sports Med* 2018; 48:1493–504.
- [44] Hislop MD, Stokes KA, Williams S, McKay CD, England ME, Kemp SPT, et al. Reducing musculoskeletal injury and concussion risk in schoolboy rugby players with a pre-activity movement control exercise programme: a cluster randomised controlled trial. *Br J Sports Med* 2017. <https://doi.org/10.1136/bjsports-2016-097434>. epub ahead of print.
- [45] Collard DC, Verhagen EA, Chinapaw MJ, Bekker I, vanMechelen W. Effectiveness of a school-based physical activity injury prevention program: a cluster randomized controlled trial. *Arch Pediatr Adolesc Med* 2010;164:145–50.
- [46] Emery CA, Cassidy JD, Klassen TP, Rosychuk R, Rowe B. Effectiveness of a home-based balance-training program in reducing sports-related injuries among healthy adolescents: a cluster randomized controlled trial. *CMAJ* 2005;172: 749–54.
- [47] Richmond SA, Kang J, Doyle-Baker PK, Nettel-Aguirre A, Emery CA. A school-based injury prevention program to reduce sport injury risk and improve healthy outcomes in youth: a pilot, clustered randomized controlled trial. *Clin J Sport Med* 2016;26:291–8.
- [48] Emery CA, van den Berg C, Richmond SA, Palacios-Derflingher, Nettel-Aguirre A, McKinlay M, et al. A school-based program to reduce injuries through neuromuscular training: iSPRINT a cluster-randomized controlled trial. *Clin J Sport Med* 2018;28(3):e49.
- [49] Sugimoto D, Myer GD, Barber-Foss KD, Hewett TE. Dosage effects of neuromuscular training intervention to reduce anterior cruciate ligament injuries in female athletes: meta- and sub-group analyses. *Sports Med* 2014;44:551–62.
- [50] Häggglund M, Atroshi I, Wagner P, Walden M. Superior compliance with a neuromuscular training programme is associated with fewer ACL injuries and fewer acute knee injuries in female adolescent football players: secondary analysis of an RCT. *Br J Sports Med* 2013;47:986–91.
- [51] Steffen K, Meeuwisse WH, Romiti M, Kang J, McKay C, Bizzini M, et al. Evaluation of how different implementation strategies of an injury prevention programme (FIFA 11+) impact team adherence and injury risk in Canadian female youth football players: a cluster-randomised trial. *Br J Sports Med* 2013;47:480–7.
- [52] Soligard T, Myklebust G, Steffen K, Holme I, Silvers H, Bissini M, et al. Comprehensive warm-up programme to prevent injuries in young female footballers: cluster randomised controlled trial. *BMJ* 2008;337:a2469.
- [53] Pasanen K, Parkkari J, Pasanen M, Hiilloskorpi H, Mäkinen T, Järvinen M, et al. Neuromuscular training and the risk of leg injuries in female floorball players: cluster randomised controlled study. *BMJ* 2008 1;337:96–102.
- [54] Fortington LV, Donaldson A, Lathlean T, Young W, Gabbe BK, Lloyd D. When "just doing it" is not enough: assessing the fidelity of player performance of an injury prevention exercise program. *J Sci Med Sports* 2015;18:272–7.
- [55] Simon SR. Quantification of human motion: gait analysis—benefits and limitations to its application to clinical problems. *J Biomech* 2004;37(12):1869–80.
- [56] Benson LC, Clermont CA, Bosnjak E, Ferber R. The use of wearable devices for walking and running gait analysis outside of the lab: a systematic review. *Gait Posture* 2018;63:124–38.
- [57] Gabbett TJ. The training-injury prevention paradox: should athletes be training smarter and harder? *Br J Sports Med* 2016; 50:273–80.
- [58] MacDonald K, Bahr R, Baltich J, Whittaker J. Validation of an inertial measurement unit for the measurement of jump count and height. *Phys Ther Sport* 2017;25:15–9.
- [59] Ghali B, Owoseye OBA, Stilling C, Palacios-Derflingher L, Jordan M, Pasanen K, et al. Workload and patellar tendinopathy in youth basketball. *Clin J Sport Med* 2018;28:e60.
- [60] Andersson SH, Bahr R, Clarsen B, Myklebust G. Preventing overuse shoulder injuries among throwing athletes: a cluster-randomised controlled trial in 660 elite handball players. *Br J Sports Med* 2017;51:1073–80.
- [61] Asker M, Brooke HL, Waldén M, Tranaeus U, Johansson F, Skillgate E, et al. Risk factors for, and prevention of, shoulder injuries in overhead sports: a systematic review with best-evidence synthesis. *Br J Sports Med* 2018;52:1312–9.
- [62] Emery CA, Kang J, Shrier I, Goulet C, Hagel B, Benson B, et al. Risk of injury associated with body checking among youth ice hockey players. *JAMA* 2010;303:2265–72.
- [63] Emery CA, Hagel B, Decloe M, McKay CD. Risk factors for injury and severe injury in youth ice hockey: a systematic review of the literature. *Inj Prev* 2010;16:113–8.
- [64] Black AM, Hagel BE, Palacios-Derflingher L, Schneider KJ, Emery CA. The risk of injury associated with body checking among Pee Wee ice hockey players: an evaluation of hockey Canada's national body checking policy change. *Br J Sports Med* 2017;51:1767–72.
- [65] Emery CA, Palacios-Derflingher L, Eliason P, Black AM, Krolikowski M, Spencer N, et al. Evaluation of body checking policy in Bantam ice hockey players: a multivariable analysis. *Clin J Sport Med* 2018;28:e48.
- [66] Kerr ZY, Yeargin S, McLeod TCV, Nitto VC, Mensh J, Dodge T, et al. Comprehensive coach education and practice contact restriction guidelines result in lower injury rates in youth American football. *Orthop J Sports Med* 2015;3. 2325967115594578.
- [67] Dizon JMR, Reyes JJB. A systematic review on the effectiveness of external ankle supports in the prevention of inversion ankle sprains among elite and recreational players. *J Sci Med Sports Exerc* 2010;13:309–17.
- [68] Janssen KW, vanMechelen W, Verhagen E. Bracing superior to neuromuscular training for the prevention of self-reported recurrent ankle sprains: a three-arm randomised controlled trial. *Br J Sports Med* 2014;48:1235–9.
- [69] Yang J, Marshall SW, Bowling JM, Runyan CW, Mueller FO, Lewis MA. Use of discretionary protective equipment and rate of lower extremity injury in high school athletes. *Am J Epidemiol* 2005;161:511–9.
- [70] Mickel TJ, Bottoni CR, Tsuji G, Chang K, Baum L, Tokushige KAS. Prophylactic bracing versus taping for the prevention of ankle sprains in high school athletes: a prospective, randomized trial. *J Foot Ankle Surg* 2006;45:360–5.

- [71] Russell K, Hagel B, Francescutti LH. The effect of wrist guards on wrist and arm injuries among snowboarders: a systematic review. *Clin J Sport Med* 2011;17:145–50.
- [72] Janda DH, Wojtyś EM, Hankin FM, Milbry E, Benedict MA. Softball sliding injuries: a prospective study comparing standard and modified bases. *JAMA* 1988;259:1848–50.
- [73] Sendre RA, Keating TM, Hornak JE, Newitt PA. Use of the hollywood impact base and standard stationary base to reduce sliding and base-running injuries in baseball and softball. *Am J Sports Med* 1994;22:450–3.
- [74] Dick R, Sauers EL, Agel J, Keuter G, Marshall SW, McCarty K, et al. Descriptive epidemiology of collegiate men's baseball injuries: national Collegiate Athletic Association injury surveillance system 1988-1989 through 2003-2004. *J Athl Train* 2007;42:183–93.
- [75] Emery CA, Roos EM, Verhagen E, Finch CF, Bennell K, Story B, et al. OARSI clinical trials recommendations: design and conduct of clinical trials for primary prevention of osteoarthritis by joint injury prevention in sport and recreation. *Osteoarthritis Cartilage* 2015;23:815–25.