



# Will Landscape Fire Increase in the Future? A Systems Approach to Climate, Fire, Fuel, and Human Drivers

Karin L. Riley<sup>1</sup> · A. Park Williams<sup>2</sup> · Shawn P. Urbanski<sup>3</sup> · David E. Calkin<sup>1</sup> · Karen C. Short<sup>1</sup> · Christopher D. O'Connor<sup>1</sup>

© This is a U.S. Government work and not under copyright protection in the US; foreign copyright protection may apply 2019

## Abstract

The extent of the Earth's surface burned annually by fires is affected by a number of drivers, including but not limited to climate. Other important drivers include the amount and type of vegetation (fuel) available and human impacts, including fire suppression, ignition, and conversion of burnable land to crops. Prior to the evolution of hominids, area burned was dictated by climate via direct influences on vegetation, aridity, and lightning. In the future, warming will be accompanied by changes in distribution, frequency, intensity, and timing of precipitation that may promote or suppress fire activity depending on location. Where area burned increases, fire may become self-regulating by reducing fuel availability. The effects of climate change on fire regimes will be strongly modulated by humans in many areas. Here, we use a systems approach to outline major drivers of changes in area burned. Due to the array of interacting drivers working in concert with climate's influence on burned area, and uncertainty in the direction and magnitude of changes in these drivers, there is very high uncertainty for much of the globe regarding how fire activity and accompanying smoke emissions will change in the coming decades.

**Keywords** Area burned · Climate change · Fire activity · Emissions · Systems approach

## Introduction

Fire has long been a widespread form of disturbance within the global environment, with patterns of fire activity varying across space and time [1, 2]. Fire has reciprocal influences on biotic assemblages, including human communities, and changes in fire regimes due to environmental or human drivers—or both—can have significant socioecological impacts (e.g., [3–6]).

This article is part of the Topical Collection on *Air Pollution*

Karin L. Riley  
kriley@fs.fed.us

<sup>1</sup> Forestry Sciences Laboratory, Rocky Mountain Research Station, US Forest Service, 800 E. Beckwith Ave., Missoula, MT 59801, USA

<sup>2</sup> Lamont-Doherty Earth Observatory, Columbia University, 61 Route 9W, Palisades, NY 10964, USA

<sup>3</sup> Fire Sciences Laboratory, Rocky Mountain Research Station, US Forest Service, 5775 W. Broadway, Missoula, MT 59808, USA

Increasingly destructive wildfires<sup>1</sup> in some regions have directed much attention to the influence of warming temperatures on future fire potential. Fire seasons are lengthening globally, via an increase in severe fire weather potential on the six continents where fire occurs naturally [7]. Within this changing fire environment, wildfires may be becoming more resistant to control actions [8, 9]—a finding consistent with public perception of and concern over an increase in fire frequency and size in North America, Mediterranean Europe, and Australia (e.g., [10]). However, satellite observations of burned area (including all landscape fire, not just wildfire) show that over the past two decades, the global annual burned area has actually declined, driven primarily by land use and socioeconomic change in Africa [10–12]. Both may be true: while global area burned has recently declined, an increasingly warm environment is causing wildfires to burn under more extreme conditions at a time when the amount of homes and

<sup>1</sup> Wildfire is one class of landscape fires, distinct from agricultural burning and other types of controlled burning or “prescribed” fire. Wildfires can be human-caused or naturally ignited, but because they are “unplanned” from a fire-management perspective, they often provoke a suppression response.

infrastructure located in the fire-vulnerable wildland-urban interface continues to grow [10, 13]. These inconsistencies demonstrate the complex interactions and forcings resulting from human manipulation of the environment including the use of fire, how climate establishes the conditions for wildfire potential, and the interplay between human activity, fire, and the changing climate [1, 14].

The complex interactions between humans and natural systems are being increasingly explored using systems-based approaches often termed social ecological systems (SES) or coupled human and natural systems (CHANS) [15, 16]. These systems approaches explore how humans influence and are in turn influenced by natural disturbance processes over space and time. A simplified example is the wildfire paradox: attempts by humans to remove wildfire from fire-dependent ecosystems have led to fuel buildup that makes future attempts to eliminate wildfire more difficult, leading to more frequent management effort, and increased losses when these efforts fail [9]. In recent years, there have been a number of efforts to map how additional social and ecological factors and feedbacks including climate change influence wildfire management and outcomes (see for example [17–20]).

In this manuscript, we take a systems approach to examining wildfire and climate change globally, acknowledging that humans and fuel are also important parts of the system and that reciprocal effects exist between many components of the system (Fig. 1). We review the contemporary literature pertaining to key drivers of landscape fire activity globally (including wildfire as well as agricultural burning and other types of controlled burning or “prescribed” fire) and accompanying atmospheric pollutants from smoke. We present a stylized systems framework that focusses on how climate, ecosystem condition (specifically burnable fuel), wildfire, and human actions interact. We use the term “drivers” to indicate these influences (e.g., climate, fuel, and human actions) which push the system toward increasing (positive forcing), decreasing (negative forcing), or relatively stable area burned. Over time, these influences are dynamic and may switch in direction or magnitude; for example, systems experiencing

positive fire feedbacks (increasing area burned) eventually become fuel-limited, introducing a negative feedback that regulates subsequent fire activity. While our main objective is to examine the question of whether burned area will increase in the future, it is not possible to address this question without understanding how climate, fire, ecosystems, and humans have interacted in the past. Therefore, we begin by examining the system of interactions before the advent of hominids and move forward in time, examining how hominids have altered these interactions and what this might mean for the future. (Note that we use the term “hominid” to refer to humans and extinct relatives, i.e., genus *Homo*, but not to great apes.)


## Fire Before Hominids

Earth is an intrinsically fire-prone planet, with the presence of an oxygenated atmosphere, fuel in the form of biomass (i.e., vegetation), and a tilted axis that drives seasonal dryness [21]. The first evidence of fire in the Earth system appeared in the fossil record approximately contemporaneously with terrestrial plant life, about 420 million years ago [22]. Continuous evidence of fossil trees and charcoal appeared since the late Devonian, approximately 390 million years ago [13]. As oxygen levels rose during the Late Permian due to the proliferation of vegetation, fires began to occur in an increasing number of ecosystems, as elevated oxygen rendered vegetation flammable at higher fuel moistures by facilitating ignition and continued combustion [22, 23].


In general, fire activity is promoted by elevated temperature, as long as vegetation is available to burn. For example, ice cores collected in Greenland and Antarctica demonstrate that periods of increased temperature coincided with elevated fire activity on millennial timescales in North and South America, between 110,000 and 10,000 years ago [24, 25]. (During this time period, inferences are necessarily at millennial to centennial scales and across broad spatial domains due to the nature of ice core records.)

Prior to the advent of hominids, the amount of fire on the landscape was a function of the abundance of fuel and the prevalence of weather conditions conducive to fire, both of which were driven by climate (Fig. 2). Climate acted as a top-down control on vegetation/fuel type and abundance. Climate also set constraints on the prevalence of conducive fire weather, with warming periods being linked to an increase in conducive weather episodes. Climate fluctuations can also influence the abundance of lightning strikes and affect whether lightning co-occurs with precipitation (which affects ignition probability). Apart from other natural ignition sources, including volcanic activity, lightning was the primary source of landscape fire before hominids.

Vegetation and fire also act in a series of forcings and feedbacks that can operate over annual to millennial timescales



**Fig. 1** The landscape fire-climate-fuel-human system. The directionality and strength of the influences are place- and time-specific



**Fig. 2** The pre-hominid landscape fire system. Climate exerts a top-down control on fuel and fire weather, which in turn together influence the levels of landscape fire activity


(Fig. 3). At millennial timescales, the evolution and spread of vegetation may be facilitated or impeded by fire. For example, the spread of C<sub>4</sub> grasses<sup>2</sup> into more mesic environments during the Late Tertiary may have been promoted by increased fire activity, which created favorable environments for grasses by opening woodland canopies [26]. Many ecosystems have evolved in concert with wildland fire and are considered “fire-adapted,” with plants showing various adaptations to fire including vigorous resprouting and serotinous cones that require fire to drop seed [21, 27]. At centennial timescales, shifts in vegetation due to changing climate may facilitate or retard fire spread (e.g., where tundra shifts to boreal forest) [24]. At annual timesteps, in ecosystems with high fire activity, recently burned areas act as a negative forcing on area burned, since it takes some time for vegetation to regrow to the point that it can carry fire [28, 29]. Thus, fire can restrict the availability of burnable fuel at the landscape level [30].

Fire may cause shifts in atmospheric composition drastic enough to cause climate forcing (Fig. 4). Widespread burning of peats and commensurate rise in atmospheric carbon dioxide (CO<sub>2</sub>) has been proposed as one of the possible mechanisms that contributed to the rapid rise in temperature at the Paleocene-Eocene thermal maximum and may have facilitated global change at the end of the Permian [23, 31].

## Fire in the Time of Hominids

Hominids evolved in eastern Africa approximately 2.5 mya. As populations grew and dispersed, they began to act as an additional driver within the system (Fig. 5).

During the Paleolithic and Mesolithic ages, hominids used fire extensively to clear ground for dwellings, to reduce




**Fig. 3** The pre-hominid wildfire system, including climate forcing on fire weather and fuel growth, abundance, and structure. A feedback loop exists between fire and fuel, where fire influences not only the rate of fuel growth but also the vegetation type (e.g., forest versus grassland) and fuel affects the frequency of fire. (When an arrow is added for the first time, in this and in subsequent diagrams, it is shown in gray. The lighter color conveys only that the arrow is new, and nothing about its strength relative to other forcings)

underbrush in order to facilitate movement, to hunt, and to regenerate plant foods for themselves and grazing animals [21]. Impacts on vegetation included the opening of closed-canopy shrublands and woodlands and their replacement by annual species [21].


Between the last glacial maximum (about 21,000 years ago) and the start of widespread agriculture (about 10,000 years ago), global biomass burning increased monotonically, according to charcoal evidence, with this trend being linked to both warming of the climate and expansion of vegetation as the ice sheets retreated [32].

As agriculture became prevalent during the Neolithic, *Homo sapiens* (i.e., humans) started to significantly alter the abundance and distribution of available fuel (Fig. 6). Humans required fire to transform some vegetation types to annual-dominated landscapes [21]. Increasing population in Europe corresponded to a widespread increase in burning between 4000 and 2000 years ago, based on charcoal records [32] and ice core records [33]. This increase may have corresponded with land clearing and conversion to pasture or agricultural land [32, 33].



**Fig. 4** The pre-hominid landscape fire system, including a fire-climate forcing

<sup>2</sup> C<sub>4</sub> grasses utilize a different photosynthesis pathway than other grasses (known as C<sub>3</sub> grasses) which makes them better adapted to areas with warm or hot conditions during the rainy period of the year [26, 176]. Thus, C<sub>4</sub> grasses often dominate today's tropical and subtropical grasslands [26, 177].

**Fig. 5** The landscape fire system, including early-hominid use of fire to influence vegetation

As human-cultivated lands expanded (along with their impacts on fuel continuity and abundance), the upward trend in global fire activity ceased (approximately 10,000 years ago) [21]. Climate was likely also a factor in the global decline in biomass burning from 1 to 900 AD, when a period of global cooling corresponded with continued expansion of human populations and cultivated lands [34]. The influences of these various drivers (i.e., expansion of cultivated land and global cooling) are difficult to apportion.

During the Medieval Climate Anomaly (~900–1300 AD), increased fire activity corresponded with localized warming in boreal Alaska [35, 36], North America's Pacific Northwest [37], and the central Rocky Mountains [38]. Conversely, during the Little Ice Age (~1500–1800 AD), fire activity decreased in many regions globally [32, 35, 39]; however, the decrease in North America also coincided with the dramatic decline in Native American populations and their traditional burning practices [40].

Between 1750 and 1870, global burning increased sharply, with this rise being linked to more pervasive human influences including population growth and accompanying land use change [34]. After 1870, burning declined abruptly, likely due to the widespread expansion of intensive agriculture,

grazing, and the eventual emergence of organized fire suppression [34].



From the systems perspective, hominids represent a new driver that uses fire to exert various controls over vegetation. As hominids migrated across the globe, they began to affect interactions among climate, fire, and ecosystems. Hominids used fire in a localized fashion to make room for dwellings and to encourage or inhibit the growth of certain plants. However, with the advent of agriculture, hominids (i.e., humans) began to use fire as a tool in the conversion of ecosystems to cultivated land—a more long-lasting, widespread, and substantial change. Thus, hominids act as both a positive forcing on fire activity (in using fire to change the landscape) and a negative forcing on fire activity (in converting burnable land to cropland or pasture, which may be burned less frequently or not at all).

## Fire in the Era of Anthropogenic Climate Change

Currently, humans have a far larger influence on the landscape fire system than in the past, due to explosive population growth and technological advances. They influence the extent and composition of available fuel, apply (both intentionally and accidentally) and suppress fire, and impact global climate (Fig. 7). Climate and fuel still exert strong influences on fire activity, however.

During the past few decades, extensive datasets, including global satellite data and weather data assimilated from thousands of recording stations, have allowed for the detailed study of the interplay between fire and weather, as well as smoke dynamics, sometimes at temporal scales as fine as sub-daily and spatial scales smaller than 1 km.

Globally, in recent decades, on the order of 301–464 million hectares (about 2–3% of the planet's land surface) has burned annually [12, 41]. Despite recent increases in

**Fig. 6** The landscape fire system, including direct and indirect influences of humans on vegetation/fuel, e.g., for sustained agricultural uses**Fig. 7** The landscape fire system, including anthropogenic climate change

temperature and fuel aridity, global burned area declined significantly by 24.3% during 1998–2015 [11]. However, this trend varies spatially in direction and strength and is inconclusive for North America, South America, Eurasia, Australia, and Southeast Asia. The global reduction in burned area was driven by a decline of 1.27% per year in Africa [11]. Trends in Africa likely result from human-mediated landscape changes [11], suggesting that, in this region, global climate forcings that would increase fire activity are being overridden by human impacts.

Human population density and prosperity affect area burned [11]. Where density and prosperity are very low (which may represent marginally habitable land), area burned tends to be low. As density and prosperity increase, landscape fire activity also increases (signifying use of fire by rural populations) before dropping precipitously in heavily populated and relatively prosperous areas [11]. The burned area decline in the latter regions is likely a function of perceived threats to highly valued resources such as homes and infrastructure, prompting extensive fire suppression efforts, sometimes at enormous cost [10].

In addition to strong direct human influences, climate exerts substantial controls on fire activity. At the global scale, fire weather seasons have been growing longer, with the global mean fire-season length increasing by approximately 18% between 1979 and 2013 [7]. Fuel aridity has increased due to increases in temperature driven by anthropogenic climate change during the past few decades; fuel aridity and fire danger metrics correlate strongly with area burned [7, 42, 43]. Temperature promotes fuel aridity by increasing the evaporative demand of the atmosphere via the vapor pressure deficit [44]. This effect is implicit via the Clausius-Clapeyron relationship, where the moisture holding capacity of the atmosphere increases exponentially with temperature ( $\sim 7\% \text{ }^{\circ}\text{C}^{-1}$ ) [45]. Although temperature is positively correlated with burned area across the western United States (US) [46, 47], the strongest correlations between area burned and weather/climate variables are found when using metrics related to flammability rather than to temperature alone (e.g., potential evapotranspiration, precipitation anomaly, or fuel moisture) [14, 43, 46, 48, 49]. In fact, one of the strongest drivers of area burned may be the number of wetting rain days (precipitation  $>= 2.54 \text{ mm}$ ) [50].

Climate changes also cause shifts in snow dynamics. Warming reduces the fraction of precipitation falling as snow and speeds the rate of snowpack melt. These changes cause the timing of runoff and infiltration to shift earlier in the year and promote lengthening of the period with dry soils and fuel during the warm fire season [51–55].

## **Smoke and Emissions**

The primary drivers of fire emissions are burned area, fuel loading (amount of material available for combustion in various biomass pools), combustion completeness (fraction of

fuel load burned), and emission factors (mass of a pollutant emitted per unit mass of fuel burned) [56]. Fuel loading and combustion completeness vary significantly across ecosystems and biomass pools, respectively. Combustion completeness of grasses and fine litter pools typically approaches 100%. In contrast, the combustion completeness of live leaves and needles, larger diameter litter (large branches and logs), standing dead trees and shrubs, and organic soil is highly variable, depending on moisture content, ambient meteorology, and terrain. Emission factors vary by fuel component, fuel conditions, and the relative share of flaming or smoldering combustion. Organic soils, peat, and large-diameter dead woody debris favor smoldering combustion, which has much higher emission factors for PM<sub>2.5</sub> than flaming combustion that dominates grasses and fine litter [57, 58].

In recent decades, approximately 2.2 Pg/year of carbon has been emitted by wildland fire globally (including emissions from both natural and human-caused fires) [59]. Carbon from fossil fuel emissions is approximately five times greater at 10 Pg/year (10,000 million metric tonnes) [60]. Following fire, carbon is often reassimilated into vegetation onsite over a period of months or years and thus biomass burning cannot be considered a net carbon flux; however, where forests do not regrow after fire due to type conversions driven by climate changes or maintenance as croplands, net carbon emissions can be significant [61].

Biomass burning is the largest source of carbonaceous aerosols and a leading source of reactive trace gases [59, 62]. Fires in Africa are the leading source of global fire emissions (accounting for 52%), followed by Asia (22%) and South America (15%) [59]. Emissions depend on the area burned, the amount of fuel (biomass available for combustion), and combustion completeness (fraction of fuel consumed) [63].

Wildfire smoke is a substantial health hazard—both acute and long-term exposures are associated with increases in respiratory and cardiovascular morbidity and mortality [64–66]. The smoke constituent presenting the primary health hazard is fine aerosol, i.e., particulate matter with an aerodynamic diameter  $< 2.5 \text{ }\mu\text{m}$  (PM<sub>2.5</sub>) [67]. Globally, it has been estimated that exposure to PM<sub>2.5</sub> from wildfire smoke results in over 300,000 premature deaths annually [68].

## **Fire and Climate in the Future**

Under future climate, changes in temperature, precipitation, humidity, and cloudiness are expected to influence fuel moisture, thereby impacting burned area and combustion completeness, and thus smoke [69–71]. Changes in human habitation and land use patterns will also affect the distribution of fire [72, 73]. The boundaries of biomes may shift, causing changes in fuel structure and type and commensurate fire activity; shifting biome


boundaries may affect climate via large changes in carbon storage as well as changes in surface energy fluxes that affect atmospheric circulation (Fig. 8) [61, 74–76].

Little is known about the relative strengths of the various drivers (e.g., humans, fuels, and climate) on area burned, or how these vary across different biomes and ecosystems [73, 77]. In addition, the magnitude of future climate change is not known due to internal variability in the climate system, uncertainty in modeling of climate response to external forcing, and uncertainty in the magnitude of external forcing due to humans [78, 79]. Thus, predictions of future fire activity are highly uncertain. In the following subsections, we examine predicted changes to key climatic influences on fire activity and underscore the level of uncertainty in these predictions. Then, we discuss likely changes in vegetation and fuels and their implications for fire and climate and end this section with a discussion of likely future fire emission scenarios and implications for air quality and climate.

### **Changes in Climate and Fire Weather: Predicted Effects on Fuel and Fire Activity**

#### **Robust Projections of Climate Change**

Projections of future climate come from General Circulation Models (GCMs), which simulate the global climate system [80]. GCMs project the globe will warm as a result of rising greenhouse gas concentration in the atmosphere, with warming most rapid over continents and at high latitudes [79]. If all else were held constant, widespread increases in wildfire activity would be expected in many regions globally due to warming-induced increases in atmospheric moisture demand and fuel aridity (which is correlated with area burned) [69, 81, 82]. In historically snowy locations, warming-induced shifts to rain and early snowmelt may also reduce soil moisture availability in the subsequent warm months [52, 83].



**Fig. 8** Climate- or human-driven changes in land use may be substantial enough to provide a forcing on global climate

Whether and how these projected increases in temperature actually affect wildfire activity, however, will be heavily dictated by the availability of fuel [82, 84, 85], which will be strongly influenced by precipitation and other climate variables. For example, reductions in precipitation could promote fire activity in very wet regions such as tropical or temperate rainforests, or suppress it by causing reduced vegetation cover in semi-arid deserts. Conversely, increases in precipitation in fuel-limited systems (such as xeric shrublands and semi-arid deserts) could promote fire activity by enhancing surface fuel abundance or suppress it by increasing fuel moisture [77, 86]. CO<sub>2</sub> fertilization, increased precipitation, and longer growing seasons may enhance ecosystem productivity [87], increasing the foliage and litterfall available for burning in some areas. In addition, humans will continue to be strong influences on vegetation change globally, via expansion of croplands, deforestation, fire suppression, etc.

While there is broad agreement on an increase in temperature at both global and local scales, changes to patterns of precipitation are subject to uncertainties associated with seasonal (e.g., monsoonal), interannual (climate mode, e.g., El Niño Southern Oscillation (ENSO)), and decadal (climate oscillations such as Interdecadal Pacific Oscillation (IPO), Atlantic Multidecadal Oscillation (AMO)) climate patterns. Robustly projected changes in precipitation include decreases in annual totals in the subtropics, particularly on the poleward flanks as the overturning Hadley cells expand poleward [88], and increases in annual totals throughout much of the tropics and high latitudes [79]. The seasonality of these changes in precipitation will be important to forecasting changes in fire activity. For example, if decreases in precipitation in the subtropics occur during the winter but not during the summer fire season, fire activity might increase less than if the decreases in precipitation occur during the summer fire season.

Temperature increases will also affect the activity of insects. For example, under climate change, developmental stages of some spruce bark beetles are expected to accelerate, facilitating faster reproduction and increased intensity of outbreaks [89–91], while the reduced frequency of fatal winter temperatures has allowed for expansion of the range of Mountain pine beetle across North America [92, 93]. Tree mortality may increase due to infestation by these insects, with trees likely being more vulnerable due to increased drought stress [94–96]. The expansion of the range of fire-sensitive spruce and fir into surrounding forest types as a result of fire exclusion is also likely to contribute to larger-scale and higher-intensity outbreaks [97]. Increased size and/or frequency of beetle outbreaks could influence subsequent fire activity, though empirical evidence for this is as yet mixed [98, 99].

Models also project increases in precipitation variability and extreme events on both interannual and intra-seasonal bases, mainly because the near-surface specific humidity of the atmosphere increases exponentially as it warms [100,

101]. Climate change may increase tree mortality through more severe droughts [102], altering forest composition and increasing the amount of standing and down woody biomass available for combustion [103]. Over time, drought-driven tree mortality would reduce water demand, potentially creating a negative feedback on subsequent mortality.

Even in the absence of changes in total mean precipitation, changes in precipitation variability would influence landscape fire regimes. More extremes on the sub-seasonal scale may enhance runoff at the expense of soil infiltration, likely reducing live and dead fuel moisture. If precipitation tends to arrive in fewer but larger storms, the number of days with wetting rain will likely decrease, contributing to an increase in fire activity [50]. More variability on interannual scales may result in more fine fuel regrowth between years of extreme fire danger, potentially increasing fire frequency and severity. Importantly, vegetation structure and species composition will be affected by changes in climate and climate-driven changes in fire severity and frequency. For example, increased fire frequency may prevent individuals of some plant species from reaching reproductive age, favoring those more adapted to short fire-return intervals [104, 105].

### Major Uncertainties in Climate Projections

While there are some robustly projected changes in large-scale temperature and precipitation (described above), projections are more uncertain for most individual regions and especially for non-temperature variables. For example, climate variability in many regions is sensitive to nuanced variations in tropical ocean and atmosphere circulation that GCMs still have trouble simulating reliably (e.g., [106]). Among the most notable sources of uncertainties are clouds. Uncertainties in the effect of anthropogenic forcing on clouds over tropical and subtropical oceans cause large uncertainties in both global temperature change and large-scale atmospheric and oceanic circulation that drive changes in regional hydroclimate [107–109] and have implications for future fire activity.

Of course, even considering variables and regions where models are in agreement as to how additional greenhouse gases affect climate, there is great uncertainty in future climate change due to uncertainty in future greenhouse gas emissions [110, 111]. Comparing various potential future emissions scenarios, the multi-model mean projected warming for 1900 to 2100 ranges from approximately 2 °C to 5 °C, depending largely on the trajectory of the twenty-first-century global greenhouse gas emissions [79].

Another uncertainty particularly relevant to extra-tropical precipitation, extreme weather, heatwaves, and fire is the degree to which rapid Arctic warming and ice melt (known as Arctic amplification) affect the jet streams and related features of extra-tropical atmospheric circulation [112–114]. It has been hypothesized that the reduced tropics-to-pole temperature gradient has

caused a slowing of the jet stream and an increase in the latitudinal amplitude of its Rossby waves in the Northern Hemisphere, leading to more persistent and extreme weather events in the Northern Hemisphere extra-tropics [115]. This could lead to exacerbated drought conditions conducive to fire [116]. However, these findings have been shown to be highly sensitive to methodology [117, 118], and there have not yet been clear and consistent trends in jet stream latitude, upper tropospheric wind speeds, or Rossby wave amplitude in observations over the past several decades [113, 119], possibly because observational records are too short for an effect of anthropogenic forcing to have yet clearly emerged [120]. While models do robustly project a poleward migration of the Northern Hemisphere jet streams, this migration is small (1°) and does not correspond to increases in Rossby wave amplitude or frequency of blocking events [121, 122]. Overall, past and future trends in large-scale extra-tropical circulation were and will be driven by far more than Arctic amplification alone and will continue to be difficult to attribute to anthropogenic forcing due to a high degree of interannual variability, and complex dynamics with competing effects, some of which are still poorly understood and modeled [113, 123].

Influences on interannual and seasonal precipitation totals are also the subject of continued research. Projections of the El Niño Southern Oscillation (ENSO), a driver of fire activity in some regions via interannual fluctuations in precipitation [124, 125], are still highly uncertain [126, 127]. In addition, models disagree as to the seasonality of precipitation changes [128], with precipitation during fire season being a stronger driver than non-fire season precipitation in many ecosystems [43, 47].

### Changes in Vegetation and Fuel: Predicted Effects on Climate and Fire

Changes in land cover will also critically affect climate and fire activity across many scales. In recent decades, increases in terrestrial net vegetation productivity have substantially reduced the rate of CO<sub>2</sub> accumulation in the atmosphere resulting from anthropogenic emissions [129]. Further, vegetation influences evapotranspiration amount and timing across all land areas. Evapotranspiration diminishes surface energy that would otherwise be partitioned toward sensible heating [130], and altered surface energy and moisture fluxes affect large-scale atmospheric circulation [131]. Future changes in vegetation cover will therefore have important impacts on soil moisture, runoff, surface temperature, and humidity at local scales, and on large-scale circulation [76, 83], with commensurate effects of fire activity.

Dynamical global vegetation models (DGVMs) used within the Coupled Model Intercomparison Project 5 (CMIP5) experiment have widely varying projections of changes in global terrestrial vegetation productivity and cover, largely due to uncertainties in effects of CO<sub>2</sub> fertilization, nutrient limitation, human land management, and the effects of climate

variability on productivity, disturbances, mortality, and succession [132–140]. Continuing population expansion and conversion of burnable land to cropland or other irrigated space can affect landscape-level burn probabilities [72]. Conversion of rainforests to croplands may liberate large amounts of carbon to the atmosphere [141]. Management interventions such as prescribed burning and fuel treatments have the potential to reduce or limit increases in future burned area, if completed at a broad enough scale [142, 143].

### **Future Landscape Fire Emissions and Implications for Air Quality and Climate**

Given the uncertainty in projections of area burned, recent studies estimate the late twenty-first-century global fire emissions will range between –15 and +62% of present-day emissions, varying with the combination of climate change scenarios and predicted human population changes [69, 144]. Under future climate, fire activity and emissions were predicted by two studies to increase in North America, Europe, and North Asia; decrease significantly in Africa (due to population changes); and remain roughly unchanged in South America [69, 144]. Increased emissions in the mid-latitudes and boreal regions were approximately offset by decreased tropical emissions. However, a third study estimated future global fire emissions would increase by 17% to 62%, with fires in South America accounting for over half the increase [70].

Factors influencing emissions vary across studies. Climate change may increase emissions through increased burned area in some or all regions [69, 70]. These increases may be roughly offset by reductions in the fuel loading available for combustion [69], but in some regions, increased precipitation and fuel availability may result in a combination of increased burned area and fuel load [144]. CO<sub>2</sub> fertilization may affect the burned area and is likely to decrease it via woody encroachment on grasslands, but some of this decrease may be offset by increased emissions due to increased fuel load [69]. Shifts in population, land use change, and timber harvest are likely to decrease emissions [70]. Though human ignitions may increase with population, the net impact of population changes is likely to decrease emissions via fire suppression [70]. The impact of land use change and harvest is expected to decrease global emissions by up to 5% to 35%, depending on the scenario considered [57].


Future emission fluxes and climate conditions are uncertain, resulting in zonal changes in black carbon concentrations (a reasonable proxy for PM<sub>2.5</sub>) varying from –50 to +100% [144]. These zonal means understate the potential human health impacts considering the atmospheric lifetime of black carbon and the magnitude of the black carbon changes over the continents, especially in active fire regions. Under future climate, some regions (e.g., central South America, northeastern Siberia, and northwestern Canada) are likely to experience

degraded air quality due to increased fire emissions; however, future wildfires will be of limited importance for air pollution in most regions [145]. In Southeast Asia, where smoke from fires used for land clearing and agriculture can be a significant health issue in the present day [146], studies project reduced emissions under future climate due to increases in population density and changes in land use [69, 70, 146].

### **Using the Systems Framework: an Example from the Western US**

The systems framework can be leveraged to aid in examining changes in drivers of fire activity under climate change. Here, we use an example from the forests of the western US, a well-studied area. Area burned has increased during the past few decades [147], which is likely an outcome of several factors. Changes in climate (increasing temperatures and decreasing summer precipitation) have acted as a positive forcing on area burned by creating additional windows of weather conducive to fires [7, 42, 50, 52, 54, 147] (Fig. 9a). Changes in fire management now allow for fires to burn and expand in some locations where they are expected to deliver benefits to highly valued resources, also acting as a positive forcing on area burned (e.g., [148]). However, fire suppression is still largely the default reaction to ignitions, causing area burned to remain below the early-twentieth-century levels [149, 150]. Thus, fire suppression acts as a negative forcing on fire activity over the short term, but contributes to increases in fuel over the longer term, which acts as a positive forcing on fire activity by reducing fuel limitation [9]. Fuel loading and connectivity are above historical levels due to a century of fire suppression and timber harvest by humans [9, 151, 152]. Because forest area burned in the western US is below historical levels, it does not significantly contribute to global climate change [150]. However, humans continue on a trajectory of aggressive burning of fossil fuels, contributing to positive forcing on area burned in this region [42, 50].

Over the next century, continued increases in temperature and decreases in fire season precipitation are likely as a result of anthropogenic fossil fuel emissions, acting as a positive feedback on area burned by further increasing the occurrence of windows of weather conducive to fire and increasing fuel aridity [153, 154] (Fig. 9b). Because fire suppression is less (if at all) effective during extreme weather and in forests with high fuel loading due to past management practices, suppression capability is likely to be overwhelmed and humans may not affect fire activity during extreme conditions [8, 155]. Unless significant changes from the status quo occur, various obstacles will continue to preclude the implementation of prescribed burning at the scale necessary to reduce fuel (which would constitute a negative (regulating) feedback on fire; obstacles include air quality regulations, lack of funding, and



**Fig. 9** The systems framework can be used to examine and understand feedbacks in fire activity under current (a) and future (b) predicted conditions. This example is for western US forests, broadly considered. Plus signs (+) indicate positive forcing, negative signs indicate negative

forcing (-); effects that are not expected to be significant are denoted with “NS,” and “?” indicates instance where the direction of the forcing cannot be predicted. Thicker arrows represent stronger drivers

strict prescriptions). If fire suppression is ineffectual as well, then humans might cease to have a significant effect on fuel in this region. However, in some regions, acceptance of managed fires has been increasing recently (“managed fires” refer to those on which land managers have a less-than-full-suppression objective), meaning that additional acreage may be burned by wildfires during less extreme weather conditions. At increased levels of fire activity, fire may become self-limiting in some areas [30]. Carbon emission levels would increase at higher levels of burning [156]; however, forests and woodlands may expand in areas where available moisture increases during the growing season, facilitated by the increase in water use efficiency from CO<sub>2</sub> fertilization or contract in areas where growing season moisture is expected to decrease and high vapor pressure deficits result in mortality of large woody vegetation [128, 157, 158]. Thus, the net effect on carbon and global climate is difficult to determine.

The net effect of future changes to the fire-climate system of the forested western US would likely be increased fire activity over the short term. Over the longer term, negative feedbacks resulting from increases in burned area and fuel limitation would constrain the system from experiencing further increases in forested burned area. However, the magnitude of these changes is uncertain, and the direction of these forcings would likely be different in other ecosystems.

that temperatures will increase, though the magnitude of the change is uncertain. Increasing temperatures can be expected to increase atmospheric moisture demand and increase fuel aridity [46]. This will have the effect of intensifying droughts, which frequently exacerbate fire. In addition, the length of the fire season is likely to increase [7]. In the short term, this may produce more rapid curing of herbaceous vegetation and contribute to water stress of all live vegetation, increasing fire activity. Moisture content and curing rate of live fuel greatly affect the probability of ignition and rate of fire spread [159, 160].

However, GCMs vary in predictions of windspeed and precipitation for a given region (direction, magnitude, and seasonality), posing challenges in predicting future fire activity [79, 161]. Warming and increased precipitation, which is likely for much of the globe, may produce greater increases in fire activity than warming and drying (for example, by promoting continuous vegetation in areas that are now xeric shrublands and semi-arid deserts) [162]. However, these increases may be punctuated by periods of reduced fire activity when fire causes fuel to be limited [14].

In addition, it is the variability in climatic conditions over decadal, annual, and seasonal scales that determine the spatial continuity and arrangement of fuel types and their availability to burn. Periods of extreme weather conducive to fire spread (hot, dry, and windy) are a prerequisite for the growth of large fires once ignition has occurred [43, 163]. Precipitation amount, seasonality, duration, and timing are all important factors in vegetation abundance and continuity, timing of vegetation curing, and whether/when dead fuels reach fuel moistures at which they are available for burning (e.g., [43, 46, 50]). The fact that the strongest correlations between burned area and climate/weather metrics are found in studies that use variables that track vegetation flammability rather than just temperature poses challenges for forecasting area burned under climate change scenarios, as flammability depends on short-term fluctuations in precipitation, temperature, and winds. Future lightning activity is another wild card, with some studies predicting increases [164]. Even if future lightning levels are similar to today’s, the location of storm tracks will greatly influence its spatial distribution. Thus, the

## Discussion: Major Challenges in Prediction of Future Fire Activity

Before human influences became globally significant, the climate-fire relationship was less complex. Currently, fire activity is driven by a system of controls and feedbacks affected not only by climate but also by human activity and its effects on vegetation and fuel, as well as human effects on climate itself (Fig. 8). The complexity of this system leads to a number of challenges in predicting future fire activity.

Many of these challenges are in the domain of forecasting future climate. Predictions from GCMs are in good agreement

relationship between fire and climate in the future depends greatly on weather conditions that are difficult to model. (GCMs do not simulate lightning directly and it must be inferred from other outputs [165].)

Another challenge in using climate conditions to predict future fire is that current fire-climate relationships cannot necessarily be used to understand the future. Extrapolations of current fire-climate relationships to no-analog conditions are likely to overpredict burned area, as recently burned areas are self-limiting, since fuel will not recover to the point where it can carry another fire for some time [14, 30, 166, 167]. The degree of self-limitation is a function of the vegetation type, time since fire, and biophysical setting related to climate [28, 29]. A second challenge is that different vegetation assemblages become flammable at different fuel moistures (e.g., [43]). As vegetation assemblages shift under future climate, extrapolating relationships between fuel moisture and burned area become dubious where novel species assemblages form, resulting in no-analog conditions (e.g., [61]).

Even if the trajectory of climate change was known, climate effects on vegetation are uncertain. At broad spatial and temporal scales, climate determines the biophysical setting for vegetation growth. While most of Earth's biomes have evolved in tandem with fire, they are adapted to specific fire regimes; changes in climate may result in departures from those regimes, with potentially disruptive effects ranging from introduction of invasive species to type conversion [21, 61]. Type conversions (e.g., a transition from forest to grassland due to widespread tree mortality) may occur due to the inability of a landscape to sustain the current vegetation due to increased water stress, for example, but more rapid and spatially extensive type conversions may occur due to disturbances such as fire (e.g., [61]). Type conversions may either increase burn probabilities, for example where closed forest is converted to shrub or grassland, or decrease burn probabilities where fuel on formerly burnable landscapes becomes too sparse. The combination of invasive grasses and fire can also be a significant factor in causing type conversions in a number of ecosystems by introducing or increasing the frequency of fire in systems poorly adapted to frequent fire [168, 169].

Challenges in modeling future fire activity are not limited to the climate and vegetation domains, but are rife in predictions of future human activities as well. Direct human influences on vegetation globally may be as powerful as those from climate, with these impacts producing both positive and negative feedbacks on global fire activity [21]. Human actions may drive increased fire levels in some cases (e.g., with fire-induced deforestation and agricultural burning) and decreased fire in others (e.g., through aggressive fire-suppression activities) (e.g., [11, 40]). At the global level, continued expansion of cropland and pasture may dampen climate-driven increases in landscape fire. Since the onset of the Medieval Climate Anomaly, humans have been able to disrupt and in some cases

uncouple the once robust relationship between biomass burning and temperature [32, 40]. Uncertainty in emissions scenarios is substantial, but predictions for the twenty-first century indicate temperature increases of double to quintuple those estimated during the Medieval Climate Anomaly and Little Ice Age. This raises a question: is there a threshold in the magnitude of climate change after which humans will no longer be able to affect this relationship? With instances of extreme weather occurring more frequently, and fire suppression being less effective under extreme weather, perhaps this threshold has already been surpassed in some areas.

An important human factor will be the trajectory of population growth. Since population density and wealth are negatively correlated with burned area [11], if society continues on a trend toward increasing population density, decreases in burned area might be expected. However, recent demographic shifts in some areas show rural populations moving to urban centers; thus, population density may be decreasing in many rural areas. When rural populations shift to urban areas, human ignitions and burned area may decrease, but when fires occur, they may grow larger due to increases in fuel connectivity [170, 171]. The strength and direction of these human effects in relationship to those in other parts of the system will ultimately determine whether fire activity increases or decreases regionally.

Projections of landscape fire and climate must necessarily rely on modeled projections of the major drivers of climate, vegetation, and human management, though these models can, to some extent, rely on empirically observed relationships and physical models [14]. Given uncertainties in climate, vegetation, and human drivers, uncertainty in predictions of future fire activity is substantial [14, 172, 173]. Currently, global fire models do not agree on whether climate change will increase or decrease the frequency of fires in individual locations, let alone on the magnitude of changes [174]. A challenge in comparing results across a suite of existing global fire models is that the set of drivers they incorporate (fuel moisture, fuel load, ignitions from lightning and anthropogenic sources, fire suppression, rate of spread, and burned area), the spatial scale at which they operate, and the level of complexity within each model vary considerably [174]. In addition, the short duration of global fire records makes validating the outputs of these predictive models difficult [14, 69]. Global fire models currently do not simulate fire with the level of complexity common in models used to render fire progression during wildfire incidents or those used for national-scale risk products (e.g., [163, 175]). However, global models incorporate broad-scale drivers and feedbacks such as the effects of CO<sub>2</sub> fertilization and projected changes to human demographics, factors that incident-level models do not currently handle (e.g., [73, 128]). The potential exists to couple more sophisticated fire models with existing vegetation models, but integrating these models will be complex and computationally intensive.

At this point, it is not possible to assign probabilities to changes in any of the human, climate, or vegetation drivers

of fire activity, meaning predictions can at best rely on scenario planning as a tool [172]. Simulations might be made in a factorial design, where the contributions of different scenarios can be assessed [14].

In summary, robust predictions of area burned under future climate require an understanding of the coupled dynamics of the major drivers of the landscape fire system (climate, fire, fuels, and humans), and how to feasibly simulate these dynamics quantitatively. The dynamics and strengths of these drivers vary not only regionally but also over time (Fig. 9), and there are likely severe limits on our abilities to project changing dynamics into the future. For example, culture and technology are major drivers of how humans interact with fire (e.g., the number of ignitions in the USA currently peaks on July 4 due to a cultural tradition and the technology that produced fireworks) but changes in culture and technology represent systems with enormous internal variability that cannot be modeled. The depth of study and the ability to simulate the landscape fire system vary regionally, with greater understanding in some well-studied regions such as the western US and Australia. Understanding is also, of course, the highest for the recent past and near-term future. For example, in the western US, the high correlation between fuel aridity and forested burned area in recent decades strongly suggests that as warming continues, burned area will continue to increase in tandem with fuel aridity into the near-term future (next few decades) (e.g., [42, 46]). However, the sensitivity of fire activity to climate in any time period is modulated by land-cover and human variables that vary across a range of time and spatial scales. The largest source of uncertainty inhibiting fire models is likely humans: how will they affect the landscape and fire? Uncertainty also arises due to a lack of understanding about how changes in climate, CO<sub>2</sub> levels, and fire will affect future vegetation communities (e.g., [128]). Uncertainty increases with time into the future [172]. In making predictions, forecasters must rely on a set of empirical and physical models, which are improving as research advances and the short timeframe of reliable observational records of fire and climate grows [14]. However, the landscape fire system is an integration of many highly coupled and complex physical, biological, and cultural systems, with hard limits likely to permanently inhibit accurate long-range projections of wildfire activity.

## Conclusions

Global fire activity in the future will be influenced by a system of climate, vegetation, and human drivers. The strength and sometimes even the direction of these drivers are challenging to predict, and system-wide feedbacks further complicate the picture.

We present here a systems framework to help make sense of this complexity. Climate, in tandem with vegetation, has been a strong driver of fire activity throughout the Earth's history and will remain so. However, human influences on

ecosystem condition/flammability and fire activity have increased over the past century due to the explosive population growth and technological advances. Humans now also exert an impact on the climate, via fossil fuel emissions, which in turn influences fire activity. The sign and magnitude of feedbacks among climate, vegetation, and fire may shift over time, depending on interactions among the various factors. For example, there are indications that changes in climate are lessening human ability to suppress fires in some regions (e.g., western Europe and the western US) but not in other regions (e.g., Africa). Looking to the future, the relative influences of the primary drivers may be deduced in some systems, at least over limited time horizons (such as a few decades). However, for other systems and over longer time horizons, how these drivers interact in this increasingly complex system is unclear.

Uncertainty associated with feedbacks, changes in precipitation, lightning, and winds translates into limited capabilities for quantitative assessment of what the future holds. Warming is likely to increase fire activity in some areas, but not everywhere. For example, warming in already arid environments may reduce fuel availability to the point where fire cannot spread, but warming in fuel-loaded areas such as western US forests may lead to increased fire at least in the short term.

In sum, landscape fire will continue to be a presence on the Earth's surface into the future, but it is difficult to say whether climate or humans will be a stronger influence on its prevalence.

## Compliance with Ethical Standards

**Conflict of Interest** On behalf of all authors, the corresponding author states that there is no conflict of interest.

**Human and Animal Rights and Informed Consent** This article does not contain any studies with human or animal subjects performed by any of the authors.

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

## References

- Bowman DMJS, Balch JK, Artaxo P, Bond WJ, Carlson JM, Cochrane MA, et al. Fire in the earth system. *Science*. 2009;324: 481–4.
- Pyne SJ. *World fire: the culture of fire on earth*. Seattle: University of Washington Press; 1995.
- Brown PM, Kaufmann MR, Sheppard WD. Long-term, landscape patterns of past fire events in a montane ponderosa pine forest of central Colorado. *Landsc Ecol*. 1999;14:513–32.
- Brooks ML, D'Antonio CM, Richardson DM, Grace JB, Keeley JE, DiTomaso JM, et al. Effects of invasive alien plants on fire regimes. *Bioscience*. 2004;54:677–88.
- Brooks ML, Matchett JR. Spatial and temporal patterns of wild fires in the Mojave Desert, 1980–2004. *J Arid Environ*. 2006;67: 148–64.

6. Morgan P, Hardy CC, Swetnam TW, Rollins MG, Long DG. Mapping fire regimes across time and space: understanding coarse and fine-scale fire patterns. *Int J Wildl Fire.* 2001;10:329–42.
7. Jolly WM, Cochrane MA, Freeborn PH, Holden ZA, Brown TJ, Williamson GJ, et al. Climate-induced variations in global wildfire danger from 1979 to 2013. *Nat Commun.* 2015;6:7537.
8. Williams J. Exploring the onset of high-impact mega-fires through a forest land management prism. *For Ecol Manage.* 2013;294:4–10. <https://doi.org/10.1016/j.foreco.2012.06.030>.
9. Calkin DE, Thompson MP, Finney MA. Negative consequences of positive feedbacks in US wildfire management. *For Ecosyst.* 2015;2:9.
10. Doerr SH, Santín C. Global trends in wildfire and its impacts: perceptions versus realities in a changing world. *Philos Trans R Soc B.* 2016;371:20150345. <https://doi.org/10.1098/rstb.2015.0345>.
11. Andela N, Morton DC, Giglio L, Chen Y, Van Der Werf GR, Kasibhatla PS, et al. A human-driven decline in global burned area. *Science.* 2017;356:1356–62.
12. Giglio L, Randerson JT, Van Der Werf GR. Analysis of daily, monthly, and annual burned area using the fourth-generation global fire emissions database (GFED4). *J Geophys Res Biogeosci.* 2013;118:317–28.
13. Theobald DM, Romme WH. Expansion of the US wildland-urban interface. *Lands Urban Plan.* 2007;83:340–54.
14. Williams AP, Abatzoglou JT. Recent advances and remaining uncertainties in resolving past and future climate effects on global fire activity. *Curr Clim Chang Rep.* 2016;2:1–14.
15. Liu J, Dietz T, Carpenter SR, Folke C, Alberti M, Redman CL, et al. Coupled human and natural systems. *Ambio.* 2009;36:639–49.
16. Liu J, Mooney H, Hull V, Davis SJ, Gaskell J, Hertel T, et al. Systems integration for global sustainability. *Science (80-).* 2015;347:963.
17. Spies TA, White EM, Kline JD, Paige Fischer A, Ager A, Bailey J, et al. Examining fire-prone forest landscapes as coupled human and natural systems. *Ecol Soc.* 2014;19:9.
18. Shindler B, Spies TA, Bolte JP, Kline JD. Integrating ecological and social knowledge: learning from CHANS research. *Ecol Soc.* 2017;22.
19. Thompson MP, Dunn CJ, Calkin DE. Systems thinking and wildland fire management. *Proc 60th Annu Meet Int Soc Syst Sci.* 2017;1(1):17.
20. Thompson MP, MacGregor DG, Dunn CJ, Calkin DE, Phipps J. Rethinking the wildland fire management system. *J For.* 2018;116:382–90.
21. Pausas JG, Keeley JE. A burning story: the role of fire in the history of life. *Bioscience.* 2009;59:593–601. <https://doi.org/10.1525/bio.2009.59.7.10>.
22. Scott AC, Glasspool IJ. The diversification of Paleozoic fire systems and fluctuations in atmospheric oxygen concentration. *Proc Natl Acad Sci.* 2006;103:10861–5. <https://doi.org/10.1073/pnas.0604090103>.
23. Glasspool IJ, Scott AC, Waltham D, Pronina N, Shao L. The impact of fire on the Late Paleozoic Earth system. *Front Plant Sci.* 2015;6:1–13. <https://doi.org/10.3389/fpls.2015.00756> abstract.
24. Fischer H, Schüpbach S, Gfeller G, Bigler M, Röhlisberger R, Erhardt T, et al. Millennial changes in North American wildfire and soil activity over the last glacial cycle. *Nat Geosci.* 2015;8:723–7.
25. Power MJ, Marlon J, Ortiz N, Bartlein PJ, Harrison SP, Mayle FE, et al. Changes in fire regimes since the last glacial maximum: an assessment based on a global synthesis and analysis of charcoal data. *Clim Dyn.* 2008;30:887–907.
26. Keeley JE, Rundel PW. Fire and the Miocene expansion of C4 grasslands. *Ecol Lett.* 2005;8:683–90.
27. Schwilk DW, Ackerly DD. Flammability and serotiny as strategies : correlated evolution in pines. *Oikos.* 2001;94:326–36.
28. Parks SA, Miller C, Holsinger LM, Baggett LS, Bird BJ. Wildland fire limits subsequent fire occurrence. *Int J Wildl Fire.* 2016;25:182–90.
29. Parks SA, Holsinger LM, Miller C, Nelson CR. Wildland fire as a self-regulating mechanism: the role of previous burns and weather in limiting fire progression. *Ecol Appl.* 2015;25:1478–92.
30. Riley KL, Thompson MP, Scott JH, Gilbertson-Day JW. A model-based framework to evaluate alternative wildfire suppression strategies. *Resources.* 2018;7. Available from: <http://www.mdpi.com/2079-9276/7/1/4>.
31. Kurtz AC, Kump LR, Arthur MA, Zachos JC, Paytan A. Early Cenozoic decoupling of the global carbon and sulfur cycles. *Paleoceanography.* 2003;18:1090.
32. Marlon JR, Bartlein PJ, Daniau AL, Harrison SP, Maezumi SY, Power MJ, et al. Global biomass burning: a synthesis and review of Holocene paleofire records and their controls. *Quat Sci Rev.* Elsevier Ltd. 2013;65:5–25. <https://doi.org/10.1016/j.quascirev.2012.11.029>.
33. Zennaro P, Kehrwald N, Marlon J, Ruddiman WF, Brucher T, Agostinelli C, et al. Europe on fire three thousand years ago: arson or climate ? *Geophys Res Lett.* 2015;42:5023–33.
34. Marlon JR, Bartlein PJ, Carcaillet C, Gavin DG, Harrison SP, Higuera PE, et al. Climate and human influences on global biomass burning over the past two millennia. *Nat Geosci.* 2008;1:697–702.
35. Kelly R, Chipman ML, Higuera PE, Stefanova I, Brubaker LB, Hu FS. Recent burning of boreal forests exceeds fire regime limits of the past 10,000 years. *Proc Natl Acad Sci.* 2013;110:13055–60. <https://doi.org/10.1073/pnas.1305069110>.
36. Barrett CM, Kelly R, Higuera PE, Hu FS. Climatic and land cover influences on the spatiotemporal dynamics of Holocene boreal fire regimes. *Ecology.* 2013;94:389–402.
37. Walsh MK, Marlon JR, Goring SJ, Brown KJ, Gavin DG. A regional perspective on Holocene fire–climate–human interactions in the Pacific northwest of North America. *Ann Assoc Am Geogr.* 2015;105:1135–57.
38. Calder WJ, Parker D, Stopka CJ, Jiménez-Moreno G, Shuman BN. Medieval warming initiated exceptionally large wildfire outbreaks in the Rocky Mountains. *Proc Natl Acad Sci.* 2015;112:13261–6. <https://doi.org/10.1073/pnas.1500796112>.
39. Power MJ, Mayle FE, Bartlein PJ, Marlon JR, Anderson RS, Behling H, et al. Climatic control of the biomass-burning decline in the Americas after AD 1500. *The Holocene.* 2013;23:3–13.
40. Taylor AH, Troutet V, Skinner CN, Stephens S. Socioecological transitions trigger fire regime shifts and modulate fire–climate interactions in the Sierra Nevada, USA, 1600–2015 CE. *Proc Natl Acad Sci.* 2016;113:13684–9.
41. Randerson JT, Chen Y, Van Der Werf GR, Rogers BM, Morton DC. Global burned area and biomass burning emissions from small fires. *J Geophys Res G Biogeosci.* 2012;117:G04012.
42. Abatzoglou JT, Williams AP. Impact of anthropogenic climate change on wildfire across western US forests. *Proc Natl Acad Sci.* 2016;113:11770–5.
43. Riley KL, Abatzoglou JT, Grenfell IC, Klene AE, Heinsch FA. The relationship of large fire occurrence with drought and fire danger indices in the western USA, 1984–2008: the role of temporal scale. *Int J Wildl Fire.* 2013;22:894–909.
44. Seager R, Hooks A, Williams AP, Cook B, Nakamura J, Henderson N. Climatology, variability, and trends in the U.S. vapor pressure deficit, an important fire-related meteorological quantity. *J Appl Meteorol Climatol.* 2015;54:1121–41.

45. Held IM, Soden BJ. Robust responses of the hydrological cycle to global warming. *J Clim.* 2006;19:5686–99.
46. Williams AP, Seager R, Macalady AK, Berkelhammer M, Crimmins MA, Swetnam TW, et al. Correlations between components of the water balance and burned area reveal new insights for predicting forest fire area in the southwest United States. *Int J Wildl Fire.* 2015;24:14–26.
47. Abatzoglou JT, Kolden CA. Relationships between climate and macroscale area burned in the western United States. *Int J Wildl Fire.* 2013;22:1003–20.
48. Sedano F, Randerson JT. Multi-scale influence of vapor pressure deficit on fire ignition and spread in boreal forest ecosystems. *Biogeosciences.* 2014;11:3739–55.
49. Morton DC, Collatz GJ, Wang D, Randerson JT, Giglio L, Chen Y. Satellite-based assessment of climate controls on US burned area. *Biogeosciences.* 2013;10:247–60.
50. Holden ZA, Swanson A, Luce CH, Jolly WM, Maneta M, Oyler JW, et al. Decreasing fire season precipitation increased recent western US forest wildfire activity. *Proc Natl Acad Sci.* 2018;115:E8349–57.
51. Barnett TP, Adam JC, Lettenmaier DP. Potential impacts of a warming climate on water availability in snow-dominated regions. *Nature.* 2005;438:303–9.
52. Westerling AL, Hidalgo HG, Cayan DR, Swetnam TW. Warming and earlier spring increase Western U.S. forest wildfire activity. *Science (80-).* 2006;313:940–3.
53. Woodhouse CA, Meko DM, MacDonald GM, Stahle DW, Cook ER. A 1,200-year perspective of 21st century drought in southwestern North America. *Proc Natl Acad Sci.* 2010;107:21283–8. <https://doi.org/10.1073/pnas.0911197107>.
54. Westerling ALR. Increasing western US forest wildfire activity: sensitivity to changes in the timing of spring. *Philos Trans R Soc B Biol Sci.* 2016;371:20150178.
55. Kitzberger T, Falk DA, Westerling AL, Swetnam TW. Direct and indirect climate controls predict heterogeneous early-mid 21st century wildfire burned area across western and boreal North America. *PLoS One.* 2017;12:e0188486.
56. Urbanski SP, Reeves MC, Corley R, Silverstein R, Hao WM. Contiguous United States wildland fire emission estimates during 2003–2015. *Earth Syst Sci Data Discuss.* 2018;10:2241–74 Available from: <https://www.earth-syst-sci-data-discuss.net/essd-2018-100/>.
57. Akagi SK, Yokelson RJ, Wiedinmyer C, Alvarado MJ, Reid JS, Karl T, et al. Emission factors for open and domestic biomass burning for use in atmospheric models. *Atmos Chem Phys.* 2011;11:4039–72.
58. Urbanski S. Wildland fire emissions, carbon, and climate: emission factors. *For Ecol Manag.* 2014;317:51–60.
59. van der Werf GR, Randerson JT, Giglio L, Van Leeuwen TT, Chen Y, Rogers BM, et al. Global fire emissions estimates during 1997–2016. *Earth Syst Sci Data.* 2017;9:697–720.
60. Boden TA, Marland G, Andres RJ. Global, regional, and national fossil-fuel CO<sub>2</sub> emissions [Internet]. Oak Ridge, TN; 2017. Available from: [http://cdiac.ess-dive.lbl.gov/trends/emis/tre\\_glob\\_2014.html](http://cdiac.ess-dive.lbl.gov/trends/emis/tre_glob_2014.html). Accessed 10/26/18.
61. Loehman RA, Reinhardt E, Riley KL. Wildland fire emissions, carbon, and climate: Seeing the forest and the trees - A cross-scale assessment of wildfire and carbon dynamics in fire-prone, forested ecosystems. *For Ecol Manag Elsevier B.V.* 2014;317:9–19. <https://doi.org/10.1016/j.foreco.2013.04.014>.
62. Vakkari V, Beukes JP, Maso MD, Aurela M, Miroslav J, van Zyl PG. Major secondary aerosol formation in southern African open biomass burning plumes. *Nat Geosci.* 2018;11:580–3 Available from: <https://www.nature.com/articles/s41561-018-0170-0>.
63. Langmann B, Duncan B, Textor C, Trentmann J, van der Werf GR. Vegetation fire emissions and their impact on air pollution and climate. *Atmos Environ.* 2009;43:107–16.
64. Fisk WJ, Chan WR. Health benefits and costs of filtration interventions that reduce indoor exposure to PM<sub>2.5</sub> during wildfires. *Indoor Air.* 2017;27:191–204.
65. Liu JC, Pereira G, Uhl SA, Bravo MA, Bell ML. A systematic review of the physical health impacts from non-occupational exposure to wildfire smoke. *Environ Res.* Elsevier. 2015;136:120–32. <https://doi.org/10.1016/j.envres.2014.10.015>.
66. Williamson GJ, Bowman DMJS, Price OF, Henderson SB, Johnston FH. A transdisciplinary approach to understanding the health effects of wildfire and prescribed fire smoke regimes. *Environ Res Lett.* IOP Publishing. 2016;11:125009.
67. Reisen F, Durán S, Flannigan M, Elliott C, Rideout K. Wildfire smoke and public health risk. *Int J Wildl Fire.* 2015;24:1029–44.
68. Johnston FH, Henderson SB, Chen Y, Randerson JT, Marlair M, DeFries RS, et al. Estimated global mortality attributable to smoke from landscape fires. *Environ Health Perspect.* 2012;120:695–701 Available from: <http://ehp.niehs.nih.gov/1104422>.
69. Knorr W, Jiang L, Arneth A. Climate, CO<sub>2</sub> and human population impacts on global wildfire emissions. *Biogeosciences.* 2016;13: 267–82.
70. Kloster S, Mahowald NM, Randerson JT, Lawrence PJ. The impacts of climate, land use, and demography on fires during the 21st century simulated by CLM-CN. *Biogeosciences.* 2012;9:509–25.
71. Riley KL, Loehman RA. Mid-21st century climate changes increase predicted fire occurrence and fire season length, northern Rocky Mountains, United States. *Ecosphere.* 2016;7.
72. Williams HL. Economic efficiency of fuel reduction treatments in the home ignition zone to mitigate wildfire risk in Montana, USA. University of Montana; 2015.
73. Le Page Y. Sensitivity of vegetation fires to climate , vegetation , and anthropogenic drivers in the HESFIRE model: consequences for fire modeling and projection uncertainties. In: Riley KL, Thompson MP, Webley P, editors. *Am Geophys Union Geophys Monogr 223 Nat Hazard Uncertain Assess Model Decis Support.* Hoboken, New Jersey: John Wiley & Sons; 2017. p. 277–85.
74. Woodall CW, Domke GM, Riley KL, Oswalt CM, Crocker SJ, Yohe GW. A framework for assessing global change risks to forest carbon stocks in the United States. *PLoS One.* 2013;8:e73222.
75. Brienen RJW, Phillips OL, Feldpausch TR, Gloor E, Baker TR, Lloyd J, et al. Long-term decline of the Amazon carbon sink. *Nature* Nature Publishing Group. 2015;519:344–8. <https://doi.org/10.1038/nature14283>.
76. Swann ALS, Laguë MM, Garcia ES, Field JP, Breshears DD, Moore DJP, et al. Continental-scale consequences of tree die-offs in North America: identifying where forest loss matters most. *Environ Res Lett.* 2018;13:055014.
77. Krawchuk MA, Moritz MA, Parisien MA, Van Dorn J, Hayhoe K. Global pyrogeography: the current and future distribution of wildfire. *PLoS One.* 2009;4:e5102.
78. Kirtman B, Power SB, Adedoyin AJ, Boer GJ, Bojariu R, Camilloni I, et al. Near-term climate change: projections and predictability. In: Stocker TF, Qin D, Plattner G-K, Tignor M, Allen SK, Boschung J, et al., editors. *Clim Chang 2013 Phys Sci Basis Contrib Work Gr I to Fifth Assess Rep Intergov Panel Clim Chang.* New York, New York: Cambridge University Press; 2013. p. 953–1028. Available from: <http://ebooks.cambridge.org/ref/id/CBO9781107415324A031>.
79. Knutti R, Sedláček J. Robustness and uncertainties in the new CMIP5 climate model projections. *Nat Clim Chang.* Nature Publishing Group. 2013;3:369–73. <https://doi.org/10.1038/nclimate1716>.
80. Gettelman A, Rood RB. Demystifying climate models: a users guide to earth system models. Springer Nature; 2016.

81. Flannigan M, Cantin AS, de Groot WJ, Wotton M, Newberry A, Gowman LM. Global wildland fire season severity in the 21st century. *For Ecol Manag.* 2013;294:54–61.
82. Bedia J, Herrera S, Gutiérrez JM, Benali A, Brands S, Mota B, et al. Global patterns in the sensitivity of burned area to fire-weather: Implications for climate change. *Agric For Meteorol.* Elsevier B.V. 2015;214–215:369–79. <https://doi.org/10.1016/j.agrformet.2015.09.002>.
83. Mankin JS, Smerdon JE, Cook BI, Williams AP, Seager R. The curious case of projected twenty-first-century drying but greening in the American West. *J Clim.* 2017;30:8689–710.
84. Bradstock RA. A biogeographic model of fire regimes in Australia: current and future implications. *Glob Ecol Biogeogr.* 2010;19:145–58.
85. Abatzoglou JT, Williams AP, Boschetti L, Zubkova M, Kolden CA. Global patterns of interannual climate-fire relationships. *Glob Chang Biol.* 2018;00:1–12. <https://doi.org/10.1111/gcb.14405>.
86. Krawchuk MA, Moritz MA. Constraints on global fire activity vary across a resource gradient. *Ecology.* 2011;92:121–32.
87. Boisvenue C, Running SW. Impacts of climate change on natural forest productivity - evidence since the middle of the 20th century. *Glob Chang Biol.* 2006;12:862–82.
88. Seager R, Naik N, Vecchi GA. Thermodynamic and dynamic mechanisms for large-scale changes in the hydrological cycle in response to global warming. *J Clim.* 2010;23:4651–68.
89. Jönsson AM, Bärring L. Future climate impact on spruce bark beetle life cycle in relation to uncertainties in regional climate model data ensembles. *Tellus Ser A Dyn Meteorol Oceanogr.* 2011;63A:158–73.
90. Lange H, Økland B, Krokene P. Thresholds in the life cycle of the spruce bark beetle under climate change. *Interjournal Complex Syst.* 2006;1648.
91. Bentz BJ, Regniere J, Fettig CJ, Hansen EM, Hayes JL, Hicke JA, et al. Climate change and bark beetles of the western United States and Canada: direct and indirect effects. *Bioscience.* 2010;60:602–13.
92. Carroll AL, Taylor SW, Régnière J, Safranyik L. Effect of climate change on range expansion by the mountain pine beetle in British Columbia. In: Shore TL, Brooks JE, Stone JE, editors. Mt Pine Beetle Symp Challenges Solut Oct 30–31, 2003 Kelowna, BC. Victoria, BC: Natural Resources Canada, Information Report BC-X-399; 2003. p. 223–32.
93. Raffa KF, Aukema BH, Bentz BJ, Carroll AL, Hicke JA, Turner MG, et al. Cross-scale drivers of natural disturbances prone to anthropogenic amplification: the dynamics of bark beetle eruptions. *Bioscience.* 2008;58:501–17 Available from: <http://academic.oup.com/bioscience/article/58/6/501/235938/Crossscale-Drivers-of-Natural-Disturbances-Prone>.
94. Anderegg WRL, Hicke JA, Fisher RA, Allen CD, Aukema J, Bentz BJ, et al. Tree mortality from drought, insects, and their interactions in a changing climate. *New Phytol.* 2015;208:674–83.
95. Trumbore S, Brando P, Hartmann H. Forest health and global change. *Science (80-).* 2015;349:814–8 Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26293952>.
96. Jactel H, Petit J, Desprez-Loustau ML, Delzon S, Piou D, Battisti A, et al. Drought effects on damage by forest insects and pathogens: a meta-analysis. *Glob Chang Biol.* 2012;18:267–76.
97. O'Connor CD, Lynch AM, Falk DA, Swetnam TW. Post-fire forest dynamics and climate variability affect spatial and temporal properties of spruce beetle outbreaks on a Sky Island mountain range. *For Ecol Manage.* Elsevier B.V. 2015;336:148–62. <https://doi.org/10.1016/j.foreco.2014.10.021>.
98. Harvey BJ, Donato DC, Turner MG. Recent mountain pine beetle outbreaks, wildfire severity, and postfire tree regeneration in the US Northern Rockies. *Proc Natl Acad Sci.* 2014;111:15120–5.
99. Hart SJ, Schoennagel T, Veblen TT, Chapman TB. Area burned in the western United States is unaffected by recent mountain pine beetle outbreaks. *Proc Natl Acad Sci.* 2015;112:4375–80. <https://doi.org/10.1073/pnas.1424037112>.
100. Pendergrass AG, Knutti R, Lehner F, Deser C, Sanderson BM. Precipitation variability increases in a warmer climate. *Sci Rep.* Springer US. 2017;7:1–9. <https://doi.org/10.1038/s41598-017-17966-y>.
101. Swain DL, Langenbrunner B, Neelin JD, Hall A. Increasing precipitation volatility in twenty-first-century California. *Nat Clim Chang.* Springer US. 2018;8:427–33. <https://doi.org/10.1038/s41558-018-0140-y>.
102. Allen CD, Macalady AK, Chenchouni H, Bachelet D, McDowell N, Vennetier M, et al. A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests. *For Ecol Manag.* 2010;259:660–84.
103. Cobb RC, Chan MN, Meentemeyer RK, Rizzo DM. Common factors drive disease and coarse woody debris dynamics in forests impacted by sudden oak death. *Ecosystems.* 2012;15:242–55.
104. Hansen WD, Braziunas KH, Rammer W, Seidl R, Turner MG. It takes a few to tango: changing climate and fire regimes can cause regeneration failure of two subalpine conifers. *Ecology.* 2018;99: 966–77.
105. Clark JS. Testing disturbance theory with long-term data: alternative life-history solutions to the distribution of events. 148. 1996;148:976–96.
106. Zhang X, Liu H, Zhang M. Double ITCZ in coupled ocean-atmosphere models: from CMIP3 to CMIP5. *Geophys Res Lett.* 2015;42:8651–9.
107. Bony S, Stevens B, Frierson DMW, Jakob C, Kageyama M, Pincus R, et al. Clouds, circulation and climate sensitivity. *Nat Geosci.* Nature Publishing Group. 2015;8:261–8. <https://doi.org/10.1038/ngeo2398>.
108. Burgman RJ, Kirtman BP, Clement AC, Vazquez H. Model evidence for low-level cloud feedback driving persistent changes in atmospheric circulation and regional hydroclimate. *Geophys Res Lett.* 2017;44:428–37.
109. Klein SA, Hall A, Norris JR, Pincus R. Low-cloud feedbacks from cloud-controlling factors: a review. *Surv Geophys* Springer Netherlands. 2017;38:1307–29.
110. van Vuuren DP, Edmonds J, Kainuma M, Riahi K, Thomson A, Hibbard K, et al. The representative concentration pathways: an overview. *Clim Chang.* 2011;109:5–31.
111. Schweizer VI, O'Neill BC. Systematic construction of global socioeconomic pathways using internally consistent element combinations. *Clim Chang.* 2014;122:431–45.
112. Cohen J, Screen JA, Furtado JC, Barlow M, Whittleston D, Coumou D, et al. Recent Arctic amplification and extreme mid-latitude weather. *Nat Geosci.* Nature Publishing Group. 2014;7: 627–37. <https://doi.org/10.1038/ngeo2234>.
113. Barnes EA, Screen JA. The impact of Arctic warming on the mid-latitude jet-stream: can it? Has it? Will it? *Wiley Interdiscip Rev Clim Chang.* 2015;6:277–86.
114. Knapp PA, Soulé PT. Spatio-temporal linkages between declining Arctic sea-ice extent and increasing wildfire activity in the western United States. *Forests.* 2017;8:1–13.
115. Francis JA, Vavrus SJ. Evidence linking Arctic amplification to extreme weather in mid-latitudes. *Geophys Res Lett.* 2012;39: L06801.
116. Dannenberg MP, Wise EK. Shifting Pacific storm tracks as stressors to ecosystems of western North America. *Glob Chang Biol.* 2017;23:4896–906.
117. Barnes EA. Revisiting the evidence linking Arctic amplification to extreme weather in midlatitudes. *Geophys Res Lett.* 2013;40: 4734–9.

118. Screen JA, Simmonds I. Exploring links between Arctic amplification and mid-latitude weather. *Geophys Res Lett.* 2013;40:959–64.
119. Woollings T, Czuchnicki C, Franzke C. Twentieth century North Atlantic jet variability. *Q J R Meteorol Soc.* 2014;140:783–91.
120. Walsh JE. Intensified warming of the Arctic: causes and impacts on middle latitudes. *Glob Planet Change.* Elsevier B.V. 2014;117:52–63. <https://doi.org/10.1016/j.gloplacha.2014.03.003>.
121. Barnes EA, Polvani L. Response of the midlatitude jets, and of their variability, to increased greenhouse gases in the CMIP5 models. *J Clim.* 2013;26:7117–35.
122. Barnes EA, Polvani LM. CMIP5 projections of arctic amplification, of the North American/North Atlantic circulation, and of their relationship. *J Clim.* 2015;28:5254–71.
123. Shaw TA, Voigt A. Tug of war on summertime circulation between radiative forcing and sea surface warming. *Nat Geosci.* 2015;8:560–6.
124. Fasullo JT, Otto-Bliesner BL, Stevenson S. ENSO's changing influence on temperature, precipitation, and wildfire in a warming climate. *Geophys Res Lett.* 2018;45:9216–25.
125. Schoennagel T, Veblen TT, Romme WH, Sibold JS, Cook ER. ENSO and PDO variability affect drought-induced fire occurrence in Rocky Mountain subalpine forests. *Ecol Appl.* 2005;15:2000–14.
126. Cai W, Santoso A, Wang G, Yeh S-W, An S-I, Cobb KM, et al. ENSO and greenhouse warming. *Nat Clim Chang.* Nature Publishing Group. 2015;5:849–59. <https://doi.org/10.1038/nclimate2743>.
127. Chen C, Cane MA, Wittenberg AT, Chen D. ENSO in the CMIP5 simulations: life cycles, diversity, and responses to climate change. *J Clim.* 2017;30:775–801. <https://doi.org/10.1175/JCLI-D-15-0901.1>.
128. Bachelet D, Sheehan T, Ferschweiler K, Abatzoglou JT. Simulating vegetation change, carbon cycling, and fire over the western United States using CMIP5 climate projections. In: Riley KL, Webley P, Thompson MP, editors. *Am Geophys Union Geophys Monogr 223 Nat Hazard Uncertain Assess Model Decis Support.* Hoboken, New Jersey: John Wiley & Sons; 2017. p. 257–75.
129. Ballantyne AP, Alden CB, Miller JB, Tans PP, White JWC. Increase in observed net carbon dioxide uptake by land and oceans during the past 50 years. *Nature.* Nature Publishing Group. 2012;488:70–3. <https://doi.org/10.1038/nature11299>.
130. Yin D, Roderick ML, Leech G, Sun F, Huang Y. The contribution of reduction in evaporative cooling to higher surface air temperatures during drought. *Geophys Res Lett.* 2014;41:7891–7.
131. Koster RD, Chang Y, Wang H, Schubert SD. Impacts of local soil moisture anomalies on the atmospheric circulation and on remote surface meteorological fields during boreal summer: a comprehensive analysis over North America. *J Clim.* 2016;29:7345–64.
132. Anderegg WRL, Ballantyne AP, Smith WK, Majkut J, Rabin S, Beaulieu C, et al. Tropical nighttime warming as a dominant driver of variability in the terrestrial carbon sink. *Proc Natl Acad Sci.* 2015;112:15591–6. <https://doi.org/10.1073/pnas.1521479112>.
133. Anderegg WRL, Schwalm C, Biondi F, Camarero JJ, Koch G, Litvak M, et al. Pervasive drought legacies in forest ecosystems and their implications for carbon cycle models. *Science (80-).* 2015;349:528–32. <https://doi.org/10.1126/science.aab1833>.
134. Schimel D, Stephens BB, Fisher JB. Effect of increasing CO<sub>2</sub> on the terrestrial carbon cycle. *Proc Natl Acad Sci.* 2015;112:436–41. <https://doi.org/10.1073/pnas.1407302112>.
135. Smith WK, Reed SC, Cleveland CC, Ballantyne AP, Anderegg WRL, Wieder WR, et al. Large divergence of satellite and earth system model estimates of global terrestrial CO<sub>2</sub> fertilization. *Nat Clim Chang.* 2016;6:306–10.
136. Wieder WR, Cleveland CC, Smith WK, Todd-Brown K. Future productivity and carbon storage limited by terrestrial nutrient availability. *Nat Geosci.* 2015;8:441–4.
137. Hartmann H, Moura CF, Anderegg WRL, Ruehr NK, Salmon Y, Allen CD, et al. Research frontiers for improving our understanding of drought-induced tree and forest mortality. *New Phytol.* 2018;218:15–28.
138. Pellegrini AFA, Ahlström A, Hobbie SE, Reich PB, Nieradzik LP, Staver AC, et al. Fire frequency drives decadal changes in soil carbon and nitrogen and ecosystem productivity. *Nature.* Nature Publishing Group. 2018;553:194–8. <https://doi.org/10.1038/nature24668>.
139. Quesada B, Arneth A, Robertson E, de Noblet-Ducoudré N. Potential strong contribution of future anthropogenic land-use and land-cover change to the terrestrial carbon cycle. *Environ Res Lett.* 2018;13:064023.
140. Trugman AT, Medvigy D, Mankin JS, Anderegg WRL. Soil moisture stress as a major driver of carbon cycle uncertainty. *Geophys Res Lett.* 2018;45:6495–503.
141. Houghton RA, Skole DL, Nobre CA, Hackler JL, Lawrence KT, Chomentowski WH. Annual fluxes of carbon from deforestation and regrowth in the Brazilian Amazon. *Nature.* 2000;403:301–4.
142. Kharabar N, Krasovskii A, Obersteiner M, Swart R, Dosio A, San-Miguel-Ayanz J, et al. Forest fires and adaptation options in Europe. *Reg Environ Chang.* 2016;16:21–30.
143. Thompson MP, Riley KL, Loeffler D, Haas JR. Modeling fuel treatment leverage: encounter rates, risk reduction, and suppression cost impacts. *Forests.* 2017;8.
144. Veira A, Lasslop G, Kloster S. Wildfires in a warmer climate: Emission fluxes, emission heights, and black carbon concentrations in 2090–2099. *J Geophys Res Atmos.* Wiley-Blackwell. 2016;121:3195–223. <https://doi.org/10.1002/2015JD024142>.
145. Knorr W, Dentener F, Lamarque J-F, Jiang L, Arneth A. Wildfire air pollution hazard during the 21st century. *Atmos Chem Phys.* 2017 [cited 2018 Jun 5];17:9223–36. <https://doi.org/10.5194/acp-17-9223-2017>.
146. Field RD, van der Werf GR, Shen SSP. Human amplification of drought-induced biomass burning in Indonesia since 1960. *Nat Geosci.* 2009;2:185–8.
147. Dennison PE, Brewer SC, Arnold JD, Moritz MA. Large wildfire trends in the western United States, 1984–2011. *Geophys Res Lett.* 2014;41:2928–33.
148. Fire Executive Council. Guidance for implementation of federal wildland fire management policy. 2009 [cited 2018 Oct 27]. Available from: [http://www.nifc.gov/policies/policies\\_documents/GIFWFMP.pdf](http://www.nifc.gov/policies/policies_documents/GIFWFMP.pdf).
149. Pyne SJ. *Fire in America: a cultural history of wildland and rural fire.* Seattle: University of Washington Press; 1997.
150. Parks SA, Miller C, Parisien M-A, Holsinger LM, Dobrowski SZ, Abatzoglou JT. Wildland fire deficit and surplus in the western United States, 1984–2012. *Ecosphere.* 2015;6:275.
151. North MP, Stephens SL, Collins BM, Agee JK, Aplet G, Franklin JF, et al. Reform forest fire management: agency incentives undermine policy effectiveness. *Science (80-).* 2015;349:1280–1.
152. Naficy C, Sala A, Keeling EG, Graham J, DeLuca TH. Interactive effects of historical logging and fire exclusion on ponderosa pine forest structure in the northern Rockies. *Ecol Appl.* 2010;20:1851–64.
153. Maloney ED, Camargo SJ, Chang E, Colle B, Fu R, Geil KL, et al. North American climate in CMIP5 experiments: part III: assessment of twenty-first-century projections. *J Clim.* 2014;27:2230–70.
154. Riley KL, Loehman RA. Mid-21st century climate changes increase predicted fire occurrence and fire season length, Northern Rocky Mountains, United States. *Ecosphere.* 2016;7:e01543.

155. Castellnou M, Prat-Guitart N, Arilla E, Larranaga A, Nebot E, Castellarnau X, et al. Empowering strategic decision making for wildfire management: avoiding the fear trap and creating a resilient landscape. *Fire Ecol.*
156. Calkin DE, Ager AA, Thompson MP, Finney MA, Lee DC, Quigley TM, et al. A comparative risk assessment framework for wildland fire management: the 2010 Cohesive Strategy science report [Internet]. Gen. Tech. Rep. RMRS-GTR-262. Fort Collins, CO; 2011. Available from: [https://www.fs.fed.us/rm/pubs/rmrs\\_gtr262.pdf](https://www.fs.fed.us/rm/pubs/rmrs_gtr262.pdf).
157. McDowell NG, Allen CD. Darcy's law predicts widespread forest mortality under climate warming. *Nat Clim Chang.* 2015;5:669–72.
158. Allen CD, Breshears DD, McDowell NG. On underestimation of global vulnerability to tree mortality and forest die-off from hotter drought in the Anthropocene. *Ecosphere.* 2015;6:129.
159. Andrews PL, Bevins CD. BehavePlus fire modeling system, version 2: overview. 2nd Int Wildl Fire Ecol Fire Manag Congr Novemb 16–20, 2003. Orlando, Florida; 2003. p. P5.11.
160. Scott JH, Burgan RE. Standard fire behavior fuel models: a comprehensive set for use with Rothermel's surface fire spread model. Fort Collins, Colorado; 2005.
161. Holtslag AAM, Svensson G, Baas P, Basu S, Beare B, Beljaars ACM, et al. Stable atmospheric boundary layers and diurnal cycles: challenges for weather and climate models. *Bull Am Meteorol Soc.* 2013;94:1691–706.
162. Batllori E, Parisien MA, Krawchuk MA, Moritz MA. Climate change-induced shifts in fire for Mediterranean ecosystems. *Glob Ecol Biogeogr.* 2013;22:1118–29.
163. Finney MA, McHugh CW, Grenfell IC, Riley KL, Short KC. A simulation of probabilistic wildfire risk components for the continental United States. *Stoch Environ Res Risk Assess.* 2011;25: 973–1000.
164. Romps DM, Seeley JT, Vollaro D, Molinari J. Projected increase in lightning strikes in the United States due to global warming. *Science (80-).* 2014;346:851–4.
165. Magi BI. Global lightning parameterization from CMIP5 climate model output. *J Atmos Ocean Technol.* 2015;32:434–52.
166. Littell JS. Drought and fire in the western USA: is climate attribution enough?. *Curr Clim Chang Reports. Current Climate Change Reports;* 2018;
167. Clark JA, Lochman RA, Keane RE. Climate changes and wildfire alter vegetation of Yellowstone National Park, but forest cover persists. *Ecosphere.* 2017;8:e01636.
168. O'Connor CD, Garfin GM, Falk DA, Swetnam TW. Human pyrogeography: a new synergy of fire, climate and people is reshaping ecosystems across the globe. *Geogr Compass.* 2011;5: 329–50.
169. Keeley JE. Fire management impacts on invasive plants in the western United States. *Conserv Biol.* 2006;20:375–84.
170. Turco M, Bedia J, Di Liberto F, Fiorucci P, Von Hardenberg J, Koutsias N, et al. Decreasing fires in Mediterranean Europe. *PLoS One.* 2016;11.
171. Chergui B, Fahd S, Santos X, Pausas JG. Socioeconomic factors drive fire-regime variability in the Mediterranean Basin. *Ecosystems Springer US.* 2018;21:619–28.
172. Riley K, Thompson M. An uncertainty analysis of wildfire modeling. In: Riley K, Webley P, Thompson M, editors. *Am Geophys Union Geophys Monogr 223 Nat Hazard Uncertain Assess Model Decis Support.* Hoboken, New Jersey: John Wiley & Sons; 2017. p. 193–213.
173. Kloster S, Lasslop G. Historical and future fire occurrence (1850 to 2100) simulated in CMIP5 Earth System Models. *Glob Planet Change.* Elsevier B.V. 2017;150:58–69. <https://doi.org/10.1016/j.gloplacha.2016.12.017>.
174. Hantson S, Arneth A, Harrison SP, Kelley DI, Prentice IC, Rabin SS, et al. The status and challenge of global fire modelling. *Biogeosciences.* 2016;13:3359–75.
175. Finney MA. FARSITE: fire area simulator – model development and evaluation. Ogden, UT; 2004.
176. Ehleringer JR, Monson RK. Evolutionary and ecological aspects of photosynthetic pathway variation. *Annu Rev Ecol Syst.* 1993;24:411–39.
177. Keeley JE, Rundel PW. Evolution of CAM and C4 carbon-concentrating mechanisms. *Int J Plant Sci.* 2003;164:S55–77. <https://doi.org/10.1086/374192>.