

Evolution in the Anthropocene: Informing Governance and Policy

Peter Sogaard Jørgensen,^{1,2} Carl Folke,^{1,2,3}
and Scott P. Carroll^{4,5}

¹Global Economic Dynamics and the Biosphere, Royal Swedish Academy of Sciences, SE104-05 Stockholm, Sweden; email: peter.sogaard.jorgensen@su.se, carl.folke@beijer.kva.se

²Stockholm Resilience Centre, Stockholm University, SE106-91 Stockholm, Sweden

³Beijer Institute of Ecological Economics, Royal Swedish Academy of Sciences, SE104-05 Stockholm, Sweden

⁴Institute for Contemporary Evolution, Davis, California 95616, USA;
email: spcarroll@ucdavis.edu

⁵Department of Entomology and Nematology, University of California, Davis, California 95616, USA

Annu. Rev. Ecol. Evol. Syst. 2019. 50:527–46

First published as a Review in Advance on
September 3, 2019

The *Annual Review of Ecology, Evolution, and
Systematics* is online at ecolsys.annualreviews.org

<https://doi.org/10.1146/annurev-ecolsys-110218-024621>

Copyright © 2019 by Annual Reviews.
All rights reserved

Keywords

eco-evolutionary dynamics, policy, management, sustainability,
evolutionary biology, complex adaptive systems

Abstract

The Anthropocene biosphere constitutes an unprecedented phase in the evolution of life on Earth with one species, humans, exerting extensive control. The increasing intensity of anthropogenic forces in the twenty-first century has widespread implications for attempts to govern both human-dominated ecosystems and the last remaining wild ecosystems. Here, we review how evolutionary biology can inform governance and policies in the Anthropocene, focusing on five governance challenges that span biodiversity, environmental management, food and other biomass production, and human health. The five challenges are: (a) evolutionary feedbacks, (b) maintaining resilience, (c) alleviating constraints, (d) coevolutionary disruption, and (e) biotechnology. Strategies for governing these dynamics will themselves have to be coevolutionary, as eco-evolutionary and social dynamics change in response to each other.

**ANNUAL
REVIEWS CONNECT**

www.annualreviews.org

- Download figures
- Navigate cited references
- Keyword search
- Explore related articles
- Share via email or social media

Anthropocene:

the term denotes the period in the history of the Earth characterized by human domination of the Earth's processes

Anthromes:

classification of the Earth's terrestrial surface according to type and degree of human interaction with ecosystems

Governance:

decision-making and power-sharing across all levels of society

1. INTRODUCTION

Humans dominate evolutionary dynamics on planet Earth (Carroll et al. 2014, Palumbi 2001). The size of the human population and the scale, connectivity, and speed of human actions are major forces in the Anthropocene biosphere (Folke et al. 2016), shaping evolution through anthropogenic environmental change (Hendry 2016). Further, at the turn of the twenty-first century, humans are increasingly in direct control of the evolution of species across the Earth (Bar-On et al. 2018). Using the concept of anthromes (Ellis et al. 2010), we can divide the degree of direct human evolutionary control into three types of evolutionary anthromes (**Figure 1**; **Supplemental Table 1**), in which humans (*a*) artificially select the dominant species covering the surface of the Earth, (*b*) select and to varying degrees control the reproduction of large grazing animals, and (*c*) influence evolution through varying levels of harvest pressure and selection. On land, humans control the propagation of dominant organisms in the almost 25% of the area made up by settlements and croplands (see “artificial” in **Figure 1**). We select the vertebrates that graze 25–34% of the land (Klein Goldewijk et al. 2011) (see “grazer” in **Figure 1**), and we have large impacts through harvesting in the remaining 45% of seminatural and wild land (see “harvest” in **Figure 1**) and, save the area covered by mariculture, in most of the ocean (see “aquatic harvest” in **Figure 1**). Through artificial selection and controlled reproduction of crops, livestock, trees, and microorganisms and by sculpting the new habitats that blanket the planet, humans—directly and indirectly—determine the constitution of species that succeed and fail. This anthropogenic impact on evolution contributes to at least three of five principles defining the Anthropocene biosphere (Williams et al. 2015).

Collectively and individually, then, *Homo sapiens* have set in motion myriad unplanned evolutionary experiments with large impacts at increasingly global scales (Smith et al. 2014). Now is the time to move away from uncoordinated and unmanaged human-driven evolution into conscious stewardship of anthropogenic evolution toward sustainability (**Figure 2**). The areas of governance





Figure 1

Anthropogenic biomes grouped by type of direct human control of evolution: artificial (*red*), where humans select the main species covering the surface; grazer (*yellow*), where humans control the evolution of the main (often domesticated) herbivorous species; and harvest (*green* on land and *blue* in the ocean), where humans select wildlife through harvesting activities. Darker colors indicate higher intensity of control. Land data reclassified from Ellis et al. (2010) (for further details, see **Supplemental Table 1**). In the ocean, hours of fisheries harvest in 2016 are shown (see also **Figure 3a**) (Kroodsma et al. 2018) on top of Exclusive Economic Zones (EEZs), in blue. EEZs indicate areas of potential longer-term harvest impacts and some areas not well covered by Kroodsma et al. (2018).

Figure 2

Governance challenges informed by evolutionary biology. (a) Dynamics of six governance challenges are sketched with links to human activities. Each example can lead to cross-scale eco-evolutionary dynamics and can be addressed through a variety of governance approaches that must also take their social context into account. (b) Color-coded examples of governance challenges and their links to relevant policy domains of the United Nations Sustainable Development Goals (SDGs). Evolutionary perspectives and insights from cultural evolution also inform governance and policies relating to partnerships, peace, inequality, and education but are not a topic of this review. Ecosystem services are the contributions of evolution to human well-being.



Policy: a course or principle of action adopted or proposed by an organization or individual

Resilience: the capacity of living systems to sustain key functions in the face of change

Evolutionary constraints: restrictions, limitations, or biases on the course or outcome of adaptive evolution

Mismatch: maladaptive environment for genotypes or phenotypes, referring to a mismatch between historical and current environments

Eco-evolutionary dynamics: the biological dynamics resulting from interaction between ecological and evolutionary processes

in which these Anthropogenic evolutionary dynamics are relevant are many. Policy areas influenced by anthropogenic evolutionary dynamics range from agendas for food security, biodiversity conservation, and natural resource management to those for good health for all and safe biotechnology. These areas are major goals or strategies for one of the most central policy objectives of the Anthropocene—namely, achieving sustainable development, as represented by the United Nations Sustainable Development Goals (Carroll et al. 2014) (**Figure 2**). Evolutionary biology is therefore an essential field of research if we are to navigate the Anthropocene successfully.

In this review, we highlight five overarching Anthropocene challenges for which evolutionary biology is central in informing governance (**Table 1**): (a) governing evolutionary feedbacks, (b) maintaining resilience and services, (c) alleviating evolutionary constraints, (d) managing consequences of disrupted coevolution, and (e) ensuring safe and successful biotechnology. Despite its importance, in many areas of governance, evolutionary biology has arguably not had an impact that reflects that importance (Carroll et al. 2014). This shortfall is likely due to a mismatch between focus areas of evolutionary biology research and the areas in need of governance and policies (Cook & Sgrò 2017). Due to that disjunction between the centrality of the field and current application needs, we start our review by gauging the development of a policy focus in the contemporary literature of evolutionary biology research and assess priorities and opportunities for increasing the relevance of that research within complex, real-world settings.

2. STATE OF APPLICATION IN GOVERNANCE

Previous reviews of applied evolutionary biology have highlighted a diverse set of applications, including directed evolution and algorithms (Bull & Wichman 2001); solutions for an array of global challenges in food, health, and environment (Carroll et al. 2014); and general applications of evolutionary principles (Hendry et al. 2011). Yet, they have been rather vague about governance and policy implications. The lack of governance and policy focus is also reflected in traditional research journals in evolutionary biology. For example, as per October 10, 2018, out of more than 1,000 published articles in two leading journals (*Evolution* and the *Journal of Evolutionary Biology*), less than a handful (3 and 1, respectively) had “management,” “policy,” or “governance” in the title. However, signs of attention to governance aspects are emerging, with several journals entirely or partially devoted to applications emerging in the last decade. Since the year 2008, the journal *Evolutionary Applications* has published 82 articles out of 990 meeting our criteria. Importantly, some of these articles are beginning to focus on the needs of decision makers in designing governance strategies and policies enlightened by evolutionary biology (e.g., Cook & Sgrò 2018, Ridley & Alexander 2016).

2.1. Evolution and Governance of Complex Adaptive Systems

Arguably, evolutionary biology has focused on detailing mechanisms for single species in controlled settings, but our general understanding of evolution across communities outside of experimental settings is limited (Barracough 2015). Several recent reviews have confronted this issue with calls for new types of theories and frameworks to elevate and expand the applications of evolutionary biology, contending with the high levels of uncertainty in complex real-world biological settings with low levels of monitoring (Barracough 2015, Santamaría & Méndez 2012). These ambitious pursuits are spurring some of the ongoing rapid development of evolutionary biology. In recent decades, the growth of theory around contemporary evolution (Palumbi 2001) and eco-evolutionary dynamics (Hendry 2016) illustrates that considerable evolutionary change can occur over few generations and interact with ecological dynamics.

Table 1 Ways that evolutionary biology informs central governance challenges of the Anthropocene

Dynamics in need of governance	Insight(s)	Governance strategies
Evolutionary feedbacks		
Selection response	Anthropogenic selection is widespread Harvest selection can lead to regime shifts and population collapse	Lower selectivity, monitor trait change, and establish refuges for harvested species
	Resistance to biocides should be expected and resistance management planned for from the start	Promote strategies that combine multiple types of selection (e.g., integrated pest management) Promote specific selection and regulate unspecific selection technologies
Anthrome colonization	Anthromes provide a new ecological opportunity that can facilitate adaptive radiations	Monitor anthrome boundaries with high probabilities of anthrome/biome shift Design settlements and production systems to promote benign species and increase resistance to harmful species
Loss of resilience and services		
Loss of resilience through loss of evolutionary potential	Anthromes are an evolutionary challenge to many species and ecosystems of value	Promote evolutionary potential or functional diversity through phylogenetic or genetic diversity Assist evolution through mixing, moving, or breeding populations
Evolutionary constraints		
Genetic constraints in novel environments	Many species will not be able to evolve through human intervention	Manage environments to resemble those of the evolutionary past Train phenotypes to respond to environment via adaptive immune and cognitive systems
Coevolutionary disruption		
Lost interactions	Many coevolved ecological interactions are being lost, with potential cascading consequences	Protect host species and keystone species by considering effects of policies on microbiomes and other symbionts
Novel interactions	New interactions are more likely to be virulent	Govern the density and connectivity of communities subject to new interactions to lower the probability of both colonization of virulent species and virulence evolution
Safe and successful biotechnology		
Gene drives and genome editing	Genome editing and gene drives potentially have great eco-evolutionary consequences	Regulate use of technologies to promote mutational specificity, evolutionary stability, and termination mechanisms
Synthetic biology and de-extinction	Resurrected species may not be eco-evolutionarily viable	Use extant proxies for extinct species until viability of de-extinct species is likely
Gain-of-function experiments with harmful organisms	Gene flow is a safety risk Evolutionary inquiry is possible through simulations and comparative approaches	Encourage evolutionary inquiry through safe methods rather than evolutionary experiments in harmful species

Complex adaptive systems (CAS):

systems in which macroscopic properties emerge from interactions among components and feed back on those lower-level interactions

Convergent evolution:

evolution of similar phenotypes or genotypes in multiple independent populations, in response to similar selection pressures, from different initial conditions

Evolutionary biology can also make explicit use of frameworks developed to deal with complexity, such as complex adaptive systems (CAS) approaches. In fact, biological systems are classic examples of CAS and display characteristic features of CAS by being path dependent and conditional yet exhibiting repeated higher-level patterns—for example, in the form of convergent evolution of species interactions (Bolnick et al. 2018). Although the type of predictability in CAS is of a different nature to that of deterministic linear systems, it may still be possible to distinguish alternative policies in terms of risk or probabilities of success, even if detailed outcomes cannot be predicted. CAS approaches may help predict diffuse coevolution in complex communities, for example, by focusing on a hub species that affects and connects many species interactions (Toju et al. 2017). The Anthropocene may in fact be viewed as a case of diffuse coevolution, with human society serving as a hub species through which eco-evolutionary dynamics of species-rich communities can be understood.

Adaptive governance is a general framework for governance of human environment systems confronted with complexity and change (Folke et al. 2005). Adaptive governance deals with fundamental uncertainties through managed experimentation and continuous learning, re-evaluation, and updating of strategies. Generalized evolutionary prescriptions for species conservation have been proposed using an adaptive governance framework (Smith et al. 2014). Design of vaccines for the seasonal flu (Morris et al. 2018) can also be seen as a concrete example of these principles put into practice.

2.2 Factoring in Social Complexity and Cultural Evolution

Governance in the Anthropocene is confronted with equally complex social dynamics (Polasky et al. 2011). In consequence, efforts to inform governance strategies with evolutionary biology must appreciate and take into account the cultural evolution of human behaviors, norms, technologies, and institutions that shape and are shaped by eco-evolutionary dynamics (Creanza et al. 2017) (**Figure 2a**). Without embracing this social side of the equation, policies will most likely be unsuccessful due to design failure. An example of how some evolutionarily informed governance strategies work in theory, but less well in practice, is agricultural pesticide use, in which a contributor to rapid evolution of pesticide resistance is the social transmission of farmer overreliance on pesticides (Dentzman et al. 2016). Hence, when resistance evolves, the response of farmers—complying with prevalent norms—matters for the future trajectory of evolution. Similarly, policies for pesticide resistance management that do not take regional differences in regulatory enforcement into account are less likely to live up to predictions of evolutionary optimal strategies (Living with Resistance project 2018). When understood well, such social dynamics can be a policy lever for governing evolutionary challenges, such as antibiotic resistance (Hallsworth et al. 2016).

3. GOVERNING EVOLUTIONARY FEEDBACKS

Evolutionary feedback to human activities is perhaps the most urgent governance challenge informed by evolutionary biology across policy domains (cf. “selection response” and “anthrome colonization” in **Figure 2b**). Here, we focus on two types of feedback: (*a*) responses to human-driven selection within Anthropocene habitats and (*b*) colonization of Anthropocene habitats (**Figure 3**). These two feedbacks often interact, but here we treat them separately for clarity (Reznick & Ghalambor 2001).

3.1. Responses to Widespread Selection

Human activities are major sources of selection and contemporary evolution for species that live in affected habitats, but these effects often go unnoticed (Alberti et al. 2017, Reznick &




Figure 3

Anthropocene selection and ecological opportunity in the twenty-first century. Panels display selection from (a) fisheries harvesting (hours per year; Kroodsma et al. 2018) and drivers of deforestation (Curtis et al. 2018), (b) livestock-associated antibiotic use (Van Boeckel et al. 2015), and (c) climate change [standardized temperature anomalies in standard deviations (SD) (Garcia et al. 2014)] and ecological opportunity presented by (d) human population (CIESIN 2017), (e) harvested crop biomass (Monfreda et al. 2008), and (f) livestock biomass (Robinson et al. 2014). All data sets, except climate change and forest loss, are presented at a logarithmic scale with base 10.

Biocide: a chemical substance intended to exert a controlling effect on any harmful organism; includes antimicrobials and pesticides

Ghalambor 2001) (**Figure 3a–c**). In particular need of governance are situations in which (a) valued species and communities of concern are unable to respond to selection (see Sections 4 and 5) or (b) respond in undesirable ways, and situations in which (c) species of potential harm respond in ways that increase their threat (Carroll et al. 2014). Evolutionary adaptation to human selection and to Anthropocene habitats by gains in defense functions—resistance evolution—are two major examples of the second and third situations, respectively.

3.1.1. Harvest evolution as early warning of change. Many species have evolved to evade human harvest, whether through behavioral evolution, evolution of smaller size to avoid traps, or evolution of features that make them less attractive to hunters (Darimont et al. 2009, Sih et al. 2011). The impact of this type of harvest evolution on society is—when documented—relatively small. For example, the economic impact of evolution toward smaller body size in harvested fish species has been argued to be of minor economic impact compared with the decline in population size induced by harvesting (Heino et al. 2015). Similarly, harvest-induced evolution in wild species has rarely been documented to have a negative impact on species that we want to protect but may instead actually improve their chances of survival—as in cases of smaller antler size or loss of tusks, which reduce desirability for trophy hunters (Chiyo et al. 2015). However, larger scale consequences of harvest-induced trait changes cannot be ruled out and may alter trophic cascades, nutrient cycling, and ecosystem stability, especially when harvesting targets ecosystem engineers or keystone species (Palkovacs et al. 2018). In such cases, evolution in harvested populations should be monitored closely and may act as early warning signals of population collapse or ecosystem regime shifts (Palkovacs et al. 2018). Strategies for minimizing these negative consequences include decreasing harvest rates and their selectivity and establishing refuges from which gene flow of nonharvested populations can mitigate change (Palkovacs et al. 2018).

3.1.2. De-escalating biocide resistance. Resistance evolution poses increasing risks to production ecosystems, human health, and environmental management, in particular through biocide resistance evolution to pesticides and antimicrobial compounds (Carroll et al. 2014). The scale of this governance challenge is significant. Antibiotic resistance is among the world's most significant health threats, cancer resistance is a contributing factor to relapse, antiviral and antimalarial resistance threatens treatments of major diseases, and pesticide resistance is a growing concern for agricultural systems, humans, and wildlife (Lipinski et al. 2016, Living with Resistance project 2018, Rev. Antimicrob. Resist. 2016). Evolution of resistance is itself associated with ecological properties, such as species richness and community composition, that help regulate ecosystem services, such as pollination, biological control, and immune and digestive function (Living with Resistance project 2018).

Resistance to biocidal technologies should always be anticipated, especially when targeted populations are large and diverse or connectivity is high to other treated environments (Living with Resistance project 2018). Preventing resistance by varying and diversifying chemical selection has been the object of studies for decades, but except in cases of strictly enforced regulation and simple ecosystems, this strategy can only delay resistance (Living with Resistance project 2018). In contrast, multimodal control strategies impose less directional selection for resistance and diversify the set of traits that are exposed to selection, thereby lowering rates of adaptation (Carroll et al. 2014). In agriculture, integrated pest management has become the modern exemplification of such strategies, making use of a range of tactics, including mechanical and chemical control, and increasing spatiotemporal variation in habitats and control mechanisms (Carrière et al. 2019) (**Figure 4a**).




Figure 4

Archetypical characterizations of selection dynamics in four major crop pest management strategies. (a) Diverse integrated pest management tactics impede pest adaptation. (b) Conventional toxic chemical sprays select efficiently for multiple resistance. (c) Narrow reliance on glyphosate application in glyphosate-resistant crops has generated a surge in resistance and escalated glyphosate dosing. (d) Endogenous insecticidal *Bacillus thuringiensis* (Bt) toxins create invariant selection, but mandated refuge plantings of susceptible crops have slowed resistance by breeding wild types to mate with and genetically neutralize the rare resistant individuals that mature on nearby Bt plants.

A key governance priority is the promotion of strategies that do not lead to escalation-prone arms races between mounting levels and modes of resistance and increasing dosing or shifts to less desirable biocides (Living with Resistance project 2018). The adoption of transgenic (tg) crops with insecticidal *Bacillus thuringiensis* (Bt) toxins or glyphosate-resistant (GR) genes illustrates how paying closer attention to simple principles of how selection is applied could help avoid unnecessary escalation in the future (Figure 4c–d). GR crops allow farmers to apply glyphosate more indiscriminately (Figure 4c). However, insecticidal Bt crops—crops with an insecticidal gene inserted—help increase the specificity of selection by exposing only insects feeding on specific parts of the crops (Carrière et al. 2019) (Figure 4d). Although both types of tg crops have seen resistance evolve, resistant weeds in fields of GR crops have spread particularly rapidly (Heap & Duke 2018). Insects have also evolved resistance to Bt crops in several regions but overall to a lesser extent, and when strictly enforced, non-Bt refuges provide a strategy to significantly slow or even avoid resistance evolution (Carrière et al. 2019). Similar governance alternatives exist for cancer treatment such as adaptive therapy and immunotherapy, which help lower selection from traditional chemotherapy and radiation therapy (Gatenby et al. 2009, Sharma et al. 2017).

3.2. Colonization of Anthromes

Although hostile to a majority of species, the large area covered by anthromes with a substantial degree of human transformation (Figure 1) also presents a vast ecological opportunity for colonizing species (Figure 3d–f). This opportunity is what evolutionary biology predicts to be the foundation for adaptive radiations (Stroud & Losos 2016). In fact, the prelude to the adaptive radiation that also succeeded all previous mass extinctions has already begun at a small scale in response to the current Anthropocene extinction crisis (Thomas 2015), although it will

Phylogenetic diversity (PD):

a measure of biodiversity that incorporates phylogenetic difference between species (e.g., measured as the sum of branch lengths)

presumably take millions of years for this radiation to rebuild past levels of diversity (Davis et al. 2018). Colonizing species can have substantial impact on society and are therefore of high governance priority. Emerging infectious diseases (EIDs) and invasive species, for example, have a major impact on human health, livestock, and wildlife (Allen et al. 2017).

Evolutionary biology offers an analytical toolkit for predicting the types of species that will succeed evolutionarily in Anthropocene habitats and where new species are likely to appear. One framework for predicting biome shifts in plants identifies the major traits of species and habitats most likely to be sources and destinations of colonization (Donoghue & Edwards 2014). Colonizing species are likely to have (a) large population size, (b) short generation time and high mutation rates, (c) preadaptation and access to these enabler adaptations, (d) been exposed to changing environments, and (e) exhibited previous biome shifts (Donoghue & Edwards 2014). Biome shifts can be expected (a) into larger biomes with ecological opportunity, such as few competitors or natural enemies; (b) out of large and biodiverse biomes; (c) between biomes that have been in direct contact for considerable time; or (d) between a retreating edge of an old biome and an expanding edge of a new, opportune biome (Donoghue & Edwards 2014). Importantly, in the Anthropocene, adjacency of biomes does not need to be through geographical proximity but can also be through trade and travel linkages (Living with Resistance project 2018). Incomplete lineage sorting, adaptive introgression, and hybridization are likely to be important processes, as they have been in other recent radiations (Henning & Meyer 2014, Thomas 2015).

Recent analyses of EIDs support several factors in the colonization framework given above. Zoonotic disease hot spots are likely to be tropical forest regions with high levels of mammal diversity and experiencing land use change (Allen et al. 2017, Jones et al. 2008). General evidence has also been found that zoonoses emerge at the nexus between natural habitats, human settlements, agriculture, and livestock (Jones et al. 2013, Parnell et al. 2017). Further, RNA viruses with high mutation rates are overrepresented among EIDs, indirectly supporting genetic access to a diversity of adaptations and large population size as predictors (Firth & Lipkin 2013). These results give reason for general optimism about the ability of refined frameworks to inform governance of anthrome colonization.

4. GOVERNING RESILIENCE AND SERVICES

We have so far dealt with the governance of rapid eco-evolutionary change, but evolutionary biology provides equally powerful insights and tools for managing resilience and ecosystem services of species and ecosystems that struggle to adapt in the Anthropocene (Carroll et al. 2014, Folke et al. 2004) (**Figure 2a**), including those under human control and human beings ourselves. Many strands of evolutionary biology predict that the species and ecosystems most vulnerable in the Anthropocene are those that have evolved in comparatively stable environments, have low population size, have little genetic variation, are spatially fragmented, are at the limit of fundamental genetic constraints, and are exposed to rapidly changing and fundamentally novel environments (Carroll et al. 2014, Hoffmann & Sgrò 2011, Lankau et al. 2011).

4.1. Genetic Variation as a Priority for Resilience

As many species will need to adapt evolutionarily to thrive in the Anthropocene (Bell 2017), a key principle for managing resilience has been to preserve genetic diversity (Sgrò et al. 2011, Smith et al. 2014). Similarly, at the ecosystem level, phylogenetic diversity (PD) of ecological assemblages has been proposed as a key metric for the resilience of ecosystem function (Faith et al. 2010) and as an indicator for assessing ecosystem vulnerability to environmental change (Díaz et al. 2013). However, the uncertainty around the correlation between PD, functional

diversity, and evolutionary potential poses challenges (Winter et al. 2013). So does developing the capacities to distinguish between general genetic variation and variation that will be critical for adaptation in the increasing wealth of genetic data (Pearse 2016). To preserve resilience at the macroevolutionary level, protected area allocation could aim to prioritize PD (Faith et al. 2010). At the intraspecific level, areas with high genetic diversity could be prioritized and, indeed, may sometimes be able to predict ranges of other threatened species (Smith et al. 2014). The use of phylogenetic analysis can also be extended to other policy and governance areas. Phylogenetic tools can help predict where discovery of new chemical compounds in the tree of life will be successful. For example, in the South African fynbos, PD predicts ecosystem services such as medically useful plants and economic value (Forest et al. 2007). In human production systems, considering PD or functional diversity is important in the selection of organisms for seeds banks, botanic gardens, and preservation in the wild (Castañeda-Álvarez et al. 2016).

Evolutionary potential: ability to adapt evolutionarily to one or multiple environments

4.2. Building Resilience Through Translocation and Assisted Evolution

Resilience can be manipulated through two basic types of intervention common in species under human control, namely moving and mixing populations in places that better match their evolved niches or through actively assisted evolution involving breeding or genetic engineering (Carroll et al. 2014). There are increasing calls for a more active role of conservation policy in enhancing resilience of wild species through these strategies (Frankham et al. 2017, Sgrò et al. 2011, Smith et al. 2014, van Oppen et al. 2015). For example, coral reef resilience may have to be strengthened through human-assisted evolution, given projections of insufficient in situ responses (van Oppen et al. 2015). Examples of assisted evolution include inducing phenotypic acclimatization through exposure to anthropogenic environmental conditions, introduction of stress-tolerant symbionts, and selectively bred or experimentally evolved species (van Oppen et al. 2015). Similarly, assisted migration and evolution of trees have been suggested to help build evolutionary resilience to climate change (Aitken & Bemmels 2016), diseases, and pests (Snieszko & Koch 2017) and will likely be central in efforts to meet the 1.5°C target (IPCC 2018). Ecosystem restoration increasingly needs to consider evolutionary dynamics to ensure long-term resilience of restored communities (Lau et al. 2019, Raimundo et al. 2018, Sgrò et al. 2011).

5. GOVERNING EVOLUTIONARY CONSTRAINTS

Sometimes the resilience of species and ecosystems cannot be enhanced by increasing their evolutionary potential, for reasons that include fundamental biological, as well as ethical and legal, constraints and concerns (**Figure 2a**). For example, fundamental boundaries of climate evolution appear to exist or be shared among distantly related species (Araújo et al. 2013), and the widespread exposure to novel environments can create both ecological and evolutionary traps (Robertson et al. 2013). A set of governance interventions is available to alleviate evolutionary constraints. Organisms or ecosystems can be moved to suitable environments (see Section 4), the environment can be changed on-site to better fit evolved niches, or the phenotypic response to the environment can be altered by training evolved phenotypic reaction norms—for example, through vaccination, physiological acclimatization, or learning (Carroll et al. 2014). In the context of new Anthropocene environments, we focus on altering the environment to better fit evolved niches, which is fundamental to conservation biology, public health, and medicine, as it underlies environmental management of threatened species and ecosystems (Ashley et al. 2003) as well as interventions to improve human health in the face of environmental pollution and lifestyle disease (Wells et al. 2017). For example, many human diseases are thought to originate from mismatches to the new Anthropocene environments, such as type 2 diabetes with high availability of glucose (Goh et al. 2014).

Nonparallel evolution:

spectrum spanning from convergent to divergent evolution, the latter consisting of evolution of increased phenotypic or genotypic distance between populations

Hygiene and biodiversity hypotheses:

two related hypotheses that reduced contact of people with the natural environment adversely affects the immunomodulatory capacity of human microbiota

Hologenome: the collective genomes of a host organism and its symbionts

Adaptive networks:

networks in which the feedback between the dynamics of species interaction structure and population-level processes shapes interactions, abundances, and traits

Three types of analysis can help predict species in need of management due to lack of adaptation and phenotype–environment mismatch: (a) analysis of evolutionary history (Ashley et al. 2003, Gluckman et al. 2011), (b) analysis of environmental exposures early in life (Gluckman et al. 2011), and (c) association analysis of genomic, phenotypic, and environmental exposure databases (Denny et al. 2013). Although the former two approaches are long established in evolutionary biology, the last has gained recent interest for disentangling environmental and evolutionary phenotypic contributions using, for example, public health register databases (Denny et al. 2013). These analyses can speed diagnosis, and even if a course of disease is not altered, attention can shift to treating the individual rather than continuing the diagnostics process and can help drive down rapidly increasing costs of developing therapeutics (Denny et al. 2016). However, some caution about the evolutionary insights that these association studies can provide is needed. In ecological settings, the complex regulation of many phenotypes, including through epistatic interactions between many genes and phenotypic plasticity, challenges the applicability of association studies (Shaw 2019). In human health settings, the many genetic conditions without a specific intervention limits options for acting on association studies (Strande & Berg 2016). Due to the prevalence of convergent and nonparallel evolution (Bolnick et al. 2018), it is important not to extrapolate beyond geographical populations. In humans, advancing knowledge of less-studied non-Western genotypes is needed (Adhikari et al. 2017). For example, lactose tolerance is monogenic in Europe but often polygenic and less well understood in other regions (Ségurel & Bon 2017).

In a cautionary example epitomizing the practical challenges of governing evolutionary constraints, the endangered Iberian lynx (*Felis pardinus*) cannot escape its remnant habitat to recolonize its former range. Assisted reintroduction is hampered by the decimation of its native rabbit prey by introduced biocontrol viruses (Ferreira & Delibes-Mateos 2011). A desperate annual vaccination program to adapt rabbit pups is difficult to complete before the annual summer virus outbreak, after which vaccinating exposed pups increases their mortality, leaving the efficacy of this intervention in question (Rouco et al. 2016).

6. GOVERNING COEVOLUTIONARY DISRUPTION

Ongoing coevolutionary processes underlie the stability of ecological communities (Lankau 2011) and hence the resilience of ecosystems and the Earth's biosphere. Yet, human transformation of the planet is disrupting and altering existing interactions and creating new ones that lack coevolutionary history (**Figure 2a**). Disruption occurs through local extinction of a variety of interactions (Valiente-Banuet et al. 2015) and the emergence of new interactions—for example, via colonization of anthromes (see Section 2). These developments have significant impact on topics of societal concern, including human health (Allen et al. 2017, von Hertzen et al. 2015), biodiversity conservation (Colwell et al. 2012, Vredenburg et al. 2010), and food and biomass production (Jones et al. 2013, Motta et al. 2018). Evolutionary biology informs these challenges through hypotheses, such as the hygiene and biodiversity hypotheses (Brooks et al. 2013, von Hertzen et al. 2015); concepts, such as the hologenome concept (Bordenstein & Theis 2015); and theory, such as the adaptive networks theory (Raimundo et al. 2018). A particular worry is that disruption of coevolved symbioses can lead to cascading ecosystem perturbations. Integration of microbiome data means that we increasingly think about most organisms, including humans, as multispecies symbioses, with the microbiome viewed as a second genome (Foster et al. 2017). Exposure of honeybee microbiomes to glyphosate is possibly contributing to honeybee colony collapse disorder (Motta et al. 2018), and disrupted human microbiomes likely play a role in autoimmune and inflammatory diseases (von Hertzen et al. 2015).

Novel interactions, which have not been subject to coevolution, are more likely to be hostile (i.e., virulent) to one or more species (Ewald 1994, 2011; Lebarbenchon et al. 2008). This is particularly the case in high-density, high-connectivity ecosystems as new species are under little selective pressure to adapt to the new easy-to-colonize habitat in ways that reduce their impact (Ewald 1994, Jones et al. 2013). This insight has broad implications for design and governance of many human production ecosystems (Jones et al. 2013) and human settlements (Ewald 2011) where these properties relax negative selection on virulence (Jones et al. 2013).

Gene editing:
targeted manipulation
of a specific site of the
genome of a living
organism by deletion,
replacement, or
insertion

7. GOVERNING BIOTECHNOLOGY

A defining principle of the Anthropocene is that the technology of one species, humans, is actively modifying a major portion of other living organisms on the planet (Gillings & Westoby 2014, Williams et al. 2015). Evolutionary biology informs the governance of biotechnology, including the identification of risks, benefits, and priority applications (Bevan et al. 2017). This process includes guiding the development of biotechnology to circumvent existing evolutionary trade-offs (Denison 2011). However, informing governance of current technologies has become increasingly difficult given rapidly advancing capabilities in construction and engineering (synthetic biology), precise editing (genome editing), and potential rapid spread (gene drives) (Drinkwater et al. 2014, Wright et al. 2013). These challenges include the lack of comparison with wild types, risk of genetic instability, and lateral gene transfer in the environment (Drinkwater et al. 2014).

7.1. Gene Editing and Gene Drives

Two new features illustrating the increasing scope for human biotechnology are the use of gene drives that can spread a mutation through an entire population of sexually reproducing organisms (Noble et al. 2017) and gene editing (with CRISPR/Cas9) that increases the array of traits that can be altered, inserted, or deleted with comparatively high precision (Singh et al. 2017). When combined, gene drives and gene editing technology have the potential to spread specific genes in populations at an unprecedented pace and level of control; therefore, concerns have been raised about their safe use and governance (Akbari et al. 2015, Jasanoff & Hurlbut 2018). Particular worries are that these systems may mutate, be transmitted to nontarget organisms, be used in a damaging way by promoting dangerous or impeding valuable species, or have hard-to-predict eco-evolutionary consequences in the wild (Drinkwater et al. 2014). Gene drives in mosquitos (Kyrou et al. 2018) and mice (Grunwald et al. 2019) have recently been tested in laboratory settings but are not immune to evolution of resistance in the wild (Unckless et al. 2017). Tests in larger confined areas will show if changes in relative fitness under competition for resources are large enough to select for resistance (Kyrou et al. 2018). A major priority before testing in the wild is the construction of evolutionarily stable mechanisms for such tg organisms to become inviable, so-called kill switches (Wright et al. 2013). With these gaps in the technology still missing, the world's governments have recently agreed to limit the use of gene drives in the wild (Callaway 2018). In humans and threatened wildlife, one of the main evolutionary challenges—besides many ethical concerns—is increasing the specificity of gene editing (Xiong et al. 2016).

7.2. Synthetic Biology and De-Extinction

Synthetic biology is the exercise of engineering new or partially new living systems. It is currently being discussed as a technology that could help halt and even reverse the loss of biodiversity, including through de-extinction, but little engagement exists between the conservation and the synthetic biology communities (Piaggio et al. 2017). From an evolutionary perspective, significant

Gain-of-function (GOF) experiments: broadly defined as experiments expected to increase the transmissibility or virulence of pathogens

Transdisciplinary research: knowledge coproduction between researchers and other individuals or organizations in society

concerns are that resurrected species are likely to have low evolutionary resilience (e.g., low genetic diversity) and be poorly adapted to both the biotic and abiotic changes that occurred since their extinction and the environment into which they are introduced (Robert et al. 2017). For example, the absence of coevolved mutualistic and parasitic microorganisms could be a problem for the viability of de-extinct species, as with time they would be exposed to colonization of new, perhaps virulent, species (Selbach et al. 2018). The practical challenges of bringing evolutionarily viable species back from extinction may be so large (even for recent extinctions) and potentially harmful for efforts to preserve threatened extant species (Bennett et al. 2017) that using functional proxies for extinct species may be a more viable, yet still challenging, approach (Steeves et al. 2017). The challenge of bringing back extinct species or replacing their functions only adds weight to the ethical argument for preventing species extinctions.

7.3. Safer Alternatives to Risky Experiments

A debate about the safety of so-called gain-of-function (GOF) experiments with potential pandemic pathogens (PPPs) emerged after the publication of two articles describing the creation of highly pathogenic viruses with airborne transmission and following reconstruction of the 1918 H1N1 pandemic influenza virus (Lipsitch 2018). Given the tremendous global risks with such experiments, it is desirable to use safer alternatives to inform governance, including evolutionary modeling and comparative approaches (Lipsitch 2018). Beyond the GOF PPP debate and stemming from the above discussion, evolutionarily safer applications of biotechnology are those that do not exert broad selection; do not disrupt or create new ecological interactions; are used at a limited extent; and have multiple mechanisms for termination.

8. ANTHROPOCENE (CO)EVOLUTION AND TRANSDISCIPLINARITY

As we have seen, evolutionary biology informs governance of both contemporary evolution and lack thereof, from the level of the gene to the level of the planet, across sectors of society. In our review, we have emphasized five common challenges of biosphere governance in the Anthropocene that evolutionary biology informs, namely (*a*) governing evolutionary feedbacks, (*b*) maintaining resilience and services, (*c*) alleviating evolutionary constraints, (*d*) managing consequences of disrupted coevolution, and (*e*) ensuring safe and successful biotechnology. In reality, these five challenges occur in parallel and interact, and decision-making will have to consider this added complexity in almost every situation. Doing so will be made easier by addressing three priorities.

First, developing a coherent theory of evolution in the Anthropocene that considers the eco-evolutionary dynamics humans are impacted by, are part of, and drive. This step entails better integration of the features defining the Anthropocene biosphere (Williams et al. 2015) and the adaptation of existing evolutionary theory to more explicitly consider the role of humans in evolution, instead of treating anthropogenic change and control as passive processes (Gillings & Westoby 2014).

Second, appreciating that governance of eco-evolutionary dynamics in the Anthropocene occurs in contexts of diverse social dynamics that can be used as leverage points for intervention (Brooks et al. 2018). This priority points toward the need for a field of coevolutionary governance of intertwined cultural and organic evolution to inform decision-making, management, and policy.

Third and finally, nurturing a culture of transdisciplinary research in evolutionary biology will help bridge the current policy gap by informing evolutionists about the complex governance context of decision makers and the critical information they need to inform decisions. Signs of such engagements are already emerging, whether in hospital therapy (Woods & Read 2015) or in design of conservation strategies and policies (Cook & Sgrò 2018, Ridley & Alexander 2016). Naturally,

this engagement will also help build trust and make decision makers more interested in the power of evolutionary inquiry for biosphere stewardship in the Anthropocene era.

SUMMARY POINTS

1. Evolutionary biology is an essential field for the successful governance of living systems in the Anthropocene. However, evolutionarily informed decision-making is limited by the lack of information and the lack of theory to inform governance of complex adaptive systems with high levels of uncertainty and complex social dynamics.
2. Evolutionary biology helps select de-escalatory strategies that avoid undesirable evolution and arms races in response to widespread anthropogenic selection and—when widespread selection cannot be avoided—helps vary and apply selection in ways that hinder unwanted responses, such as biocide resistance.
3. Evolutionary biology predicts where adaptive colonization of human transformed habitats is most likely. These insights are critical components informing design of monitoring strategies for emerging infectious diseases and managing general anthrome colonization.
4. Preservation of resilience and ecosystem services of many species and ecosystems can be achieved by incorporating indicators of evolutionary potential into governance or by creating interventions that enhance potential for, or directly assist, adaptive evolution.
5. When evolutionary adaptation is not possible or desirable, evolutionary constraints can be governed by managing the environment to better fit historical environments where species evolved or by training adaptive subsystems such as immune systems, physiological systems, or cognitive systems.
6. Disruption of coevolved interactions can have cascading consequences for ecosystem function and human health, and protecting host–microbiome interactions is an urgent priority in all areas of governance, for example, by considering adaptive networks. Novel interactions, likely to be more virulent, should be monitored for in high-risk zones, and system features, such as density and connectivity, should be governed in urban areas and production ecosystems to lower probabilities of new virulent interactions.
7. Evolutionarily informed governance of biotechnology’s rapidly expanding abilities to construct and manipulate living systems is urgently needed, including in the context of genome editing, gene drives, synthetic biology, and experiments with potential serious and widespread consequences. Mutational specificity, evolutionary stability, termination mechanisms, evolutionary safe alternatives, and eco-evolutionary viability are key principles to inform policies.

DISCLOSURE STATEMENT

The authors are not aware of any affiliations, memberships, funding, or financial holdings that might be perceived as affecting the objectivity of this review.

ACKNOWLEDGMENTS

This work was funded through FORMAS grant 2016-00451 (P.S.J.) and by The Erling-Persson Family Foundation (P.S.J., C.F.), the MISTRA foundation (P.S.J., C.F.), the Kjell and Märta Beijer’s Foundation (C.F.), and the Institute for Contemporary Evolution (S.P.C.).

LITERATURE CITED

- Adhikari K, Chacón-Duque JC, Mendoza-Revilla J, Fuentes-Guajardo M, Ruiz-Linares A. 2017. The genetic diversity of the Americas. *Annu. Rev. Genom. Hum. Genet.* 18:277–96
- Aitken SN, Bemmels JB. 2016. Time to get moving: assisted gene flow of forest trees. *Evol. Appl.* 9(1):271–90
- Akbari BOS, Bellen HJ, Bier E, Simon L, Burt A, et al. 2015. Safeguarding gene drive experiments in the laboratory. *Science* 349(6251):927–29
- Alberti M, Correa C, Marzluff JM, Hendry AP, Palkovacs EP, et al. 2017. Global urban signatures of phenotypic change in animal and plant populations. *PNAS* 144:8951–56
- Allen T, Murray KA, Zambrana-Torrel C, Morse SS, Rondinini C, et al. 2017. Global hotspots and correlates of emerging zoonotic diseases. *Nat. Commun.* 8(1):1–10
- Araújo MB, Ferri-Yáñez F, Bozinovic F, Marquet PA, Valladares F, Chown SL. 2013. Heat freezes niche evolution. *Ecol. Lett.* 16(9):1206–19
- Ashley MV, Willson MF, Pergams ORW, O'Dowd DJ, Gende SM, Brown JS. 2003. Evolutionarily enlightened management. *Biol. Conserv.* 111(2):115–23
- Bar-On YM, Phillips R, Milo R, Falkowski PG. 2018. The biomass distribution on Earth. *PNAS* 115(28):6506–11
- Barracough TG. 2015. How do species interactions affect evolutionary dynamics across whole communities? *Annu. Rev. Ecol. Evol. Syst.* 46:25–48
- Bell G. 2017. Evolutionary rescue. *Annu. Rev. Ecol. Evol. Syst.* 48:605–27
- Bennett JR, Maloney RF, Steeves TE, Brazill-Boast J, Possingham HP, Seddon PJ. 2017. Spending limited resources on de-extinction could lead to net biodiversity loss. *Nat. Ecol. Evol.* 1:0053
- Bevan MW, Uauy C, Wulff BBH, Zhou J, Krasileva K, Clark MD. 2017. Genomic innovation for crop improvement. *Nature* 543:346–54
- Bolnick DI, Barrett RDH, Oke KB, Rennison DJ, Stuart YE. 2018. (Non)parallel evolution. *Annu. Rev. Ecol. Evol. Syst.* 49:303–30
- Bordenstein SR, Theis KR. 2015. Host biology in light of the microbiome: ten principles of holobionts and hologenomes. *PLOS Biol.* 13(8):e1002226
- Brooks C, Pearce N, Douwes J. 2013. The hygiene hypothesis in allergy and asthma: an update. *Curr. Opin. Allergy Clin. Immunol.* 13(1):70–77
- Brooks JS, Waring TM, Borgerhoff Mulder M, Richerson PJ. 2018. Applying cultural evolution to sustainability challenges: an introduction to the special issue. *Sustain. Sci.* 13(1):1–8
- Bull JJ, Wichman HA. 2001. Applied evolution. *Annu. Rev. Ecol. Syst.* 32:183–217
- Callaway E. 2018. UN treaty agrees to limit gene drives but rejects a moratorium. *Nature*, Nov. 29. <https://doi.org/10.1038/d41586-018-07600-w>
- Carrière Y, Brown ZS, Downes SJ, Gujar G, Epstein G, et al. 2019. Governing evolution: a socioecological comparison of resistance management for insecticidal transgenic Bt crops among four countries. *Ambio*. <https://doi.org/10.1007/s13280-019-01167-0>
- Carroll SP, Jørgensen PS, Kinnison MT, Bergstrom CT, Denison RF, et al. 2014. Applying evolutionary biology to address global challenges. *Science* 346(6207):1245993
- Castañeda-Álvarez NP, Khoury CK, Achicanoy HA, Bernau V, Dempewolf H, et al. 2016. Global conservation priorities for crop wild relatives. *Nat. Plants* 2(4):16022
- Chiyo PI, Obanda V, Korir DK. 2015. Illegal tusk harvest and the decline of tusk size in the African elephant. *Ecol. Evol.* 5(22):5216–29
- CIESIN (Cent. Int. Earth Sci. Inf. Netw.), Columbia Univ. 2017. *Gridded Population of the World, Version 4 (GPWv4): population density, revision 10*. Rep., NASA Socioecon. Data Appl. Cent. (SEDAC), Palisades, NY, accessed Nov. 1, 2018. <https://doi.org/10.7927/H4VH5KS4>
- Colwell RK, Dunn RR, Harris NC. 2012. Coextinction and persistence of dependent species in a changing world. *Annu. Rev. Ecol. Evol. Syst.* 43:183–203
- Cook CN, Sgrò CM. 2017. Aligning science and policy to achieve evolutionarily enlightened conservation. *Conserv. Biol.* 31(3):501–12

- Cook CN, Sgrò CM. 2018. Understanding managers' and scientists' perspectives on opportunities to achieve more evolutionarily enlightened management in conservation. *Evol. Appl.* 11(8):1371–88
- Creanza N, Kolodny O, Feldman MW. 2017. Cultural evolutionary theory: how culture evolves and why it matters. *PNAS* 114(30):7782–89
- Curtis PG, Slay CM, Harris NL, Tyukavina A, Hansen MC. 2018. Classifying drivers of global forest loss. *Science* 361(6407):1108–11
- Darimont CT, Carlson SM, Kinnison MT, Paquet PC, Reimchen TE, Wilmer CC. 2009. Human predators outpace other agents of trait change in the wild. *PNAS* 106(3):952–54
- Davis M, Faurby S, Svenning J-C. 2018. Mammal diversity will take millions of years to recover from the current biodiversity crisis. *PNAS* 115(44):11262–67
- Denison RF. 2011. Past evolutionary tradeoffs represent opportunities for crop genetic improvement and increased human lifespan. *Evol. Appl.* 4(2):216–24
- Denny JC, Bastarache L, Ritchie MD, Carroll RJ, Zink R, et al. 2013. Systematic comparison of phenome-wide association study of electronic medical record data and genome-wide association study data. *Nat. Biotechnol.* 31(12):1102–11
- Denny JC, Bastarache L, Roden DM. 2016. Phenome-wide association studies as a tool to advance precision medicine. *Annu. Rev. Genom. Hum. Genet.* 17:353–73
- Dentzman K, Gunderson R, Jussaume R. 2016. Techno-optimism as a barrier to overcoming herbicide resistance: comparing farmer perceptions of the future potential of herbicides. *J. Rural Stud.* 48:22–32
- Díaz S, Purvis A, Cornelissen JHC, Mace GM, Donoghue MJ, et al. 2013. Functional traits, the phylogeny of function, and ecosystem service vulnerability. *Ecol. Evol.* 3(9):2958–75
- Donoghue MJ, Edwards EJ. 2014. Biome shifts and niche evolution in plants. *Annu. Rev. Ecol. Evol. Syst.* 45:547–72
- Drinkwater K, Kuikem T, Lightfoot S, McNamara J, Oye K. 2014. *Creating a research agenda for the ecological implications of synthetic biology*. Rep., Synthetic Biology Project, SYN BIO 7, Washington, DC
- Ellis EC, Klein Goldewijk K, Siebert S, Lightman D, Ramankutty N. 2010. Anthropogenic transformation of the biomes, 1700 to 2000. *Glob. Ecol. Biogeogr.* 19:589–606
- Ewald PW. 1994. *Evolution of Infectious Disease*. Oxford, UK: Oxford Univ. Press
- Ewald PW. 2011. Evolution of virulence, environmental change, and the threat posed by emerging and chronic diseases. *Ecol. Res.* 26(6):1017–26
- Faith DP, Magallón S, Hendry AP, Conti E, Yahara T, Donoghue MJ. 2010. Ecosystem services: an evolutionary perspective on the links between biodiversity and human well-being. *Curr. Opin. Environ. Sustain.* 2(1–2):66–74
- Ferreira C, Delibes-Mateos M. 2011. Wild rabbit management in the Iberian Peninsula: state of the art and future perspectives for Iberian lynx conservation. *Wildl. Biol. Pract.* 6(3):48–66
- Firth C, Lipkin WI. 2013. The genomics of emerging pathogens. *Annu. Rev. Genom. Hum. Genet.* 14:281–300
- Folke C, Biggs R, Norström AV, Reyers B, Rockström J. 2016. Social-ecological resilience and biosphere-based sustainability science. *Ecol. Soc.* 21(3):41
- Folke C, Carpenter S, Walker B, Scheffer M, Elmqvist T, et al. 2004. Regime shifts, resilience, and biodiversity in ecosystem management. *Annu. Rev. Ecol. Evol. Syst.* 35:557–81
- Folke C, Hahn T, Olsson P, Norberg J. 2005. Adaptive governance of social-ecological systems. *Annu. Rev. Environ. Resour.* 30:441–73
- Forest F, Grenyer R, Rouget M, Davies TJ, Cowling RM, et al. 2007. Preserving the evolutionary potential of floras in biodiversity hotspots. *Nature* 445(7129):757–60
- Foster KR, Schluter J, Coyte KZ, Rakoff-Nahoum S. 2017. The evolution of the host microbiome as an ecosystem on a leash. *Nature* 548(7665):43–51
- Frankham R, Ballou JD, Ralls K, Eldridge M, Dudash MR, et al. 2017. *Genetic Management of Fragmented Animal and Plant Populations*. Oxford, UK: Oxford Univ. Press
- Garcia RA, Cabeza M, Rahbek C, Araújo MB. 2014. Multiple dimensions of climate change and their implications for biodiversity. *Science* 344(6183):1247579
- Gatenby RA, Silva AS, Gillies RJ, Frieden BR. 2009. Adaptive therapy. *Cancer Res.* 69(11):4894–4903

- Gillings MR, Westoby M. 2014. DNA technology and evolution of the Central Dogma. *Trends Ecol. Evol.* 29(1):1–2
- Gluckman PD, Low FM, Buklijas T, Hanson MA, Beedle AS. 2011. How evolutionary principles improve the understanding of human health and disease. *Evol. Appl.* 4(2):249–63
- Goh K-P, Koh A, Wackerhage H. 2014. Human evolution, type 2 diabetes mellitus and exercise. In *Molecular Exercise Physiology*, ed. H Wackerhage, pp. 205–27. New York: Routledge
- Grunwald HA, Gantz VM, Poplawski G, Xu XS, Bier E, Cooper KL. 2019. Super-Mendelian inheritance mediated by CRISPR–Cas9 in the female mouse germline. *Nature* 566:105–9
- Hallsworth M, Chadborn T, Sallis A, Sanders M, Berry D, et al. 2016. Provision of social norm feedback to high prescribers of antibiotics in general practice: a pragmatic national randomised controlled trial. *Lancet* 387(10029):1743–52
- Heap I, Duke SO. 2018. Overview of glyphosate-resistant weeds worldwide. *Pest Manag. Sci.* 74(5):1040–49
- Heino M, Díaz Pauli B, Dieckmann U. 2015. Fisheries-induced evolution. *Annu. Rev. Ecol. Syst.* 46:461–80
- Hendry AP. 2016. Introduction and conceptual framework. In *Eco-Evolutionary Dynamics*, pp. 1–23. Princeton, NJ: Princeton Univ. Press
- Hendry AP, Kinnison MT, Heino M, Day T, Smith TB, et al. 2011. Evolutionary principles and their practical application. *Evol. Appl.* 4(2):159–83
- Henning F, Meyer A. 2014. The evolutionary genomics of cichlid fishes: explosive speciation and adaptation in the postgenomic era. *Annu. Rev. Genom. Hum. Genet.* 15:417–41
- Hoffmann AA, Sgrò CM. 2011. Climate change and evolutionary adaptation. *Nature* 470(7335):479–85
- IPCC (Intergov. Panel Clim. Change). 2018. *Global warming of 1.5°C: summary for policymakers*. Rep., IPCC, Geneva. https://www.ipcc.ch/site/assets/uploads/sites/2/2019/05/SR15_SPM_version_report_LR.pdf
- Jasanoff S, Hurlbut BJ. 2018. A global observatory for gene editing. *Nature* 555:435–37
- Jones BA, Grace D, Kock R, Alonso S, Rushton J, Said MY. 2013. Zoonosis emergence linked to agricultural intensification and environmental change. *PNAS* 110(21):8399–8404
- Jones KE, Patel NG, Levy MA, Storeygard A, Balk D, et al. 2008. Global trends in emerging infectious diseases. *Nature* 451(7181):990–93
- Klein Goldewijk K, Beusen A, van Drecht G, de Vos M. 2011. The HYDE 3.1 spatially explicit database of human-induced global land-use change over the past 12,000 years. *Glob. Ecol. Biogeogr.* 20(1):73–86
- Kroodsmä DA, Mayorga J, Hochberg T, Miller NA, Boerder K, et al. 2018. Tracking global footprint of fisheries. *Science* 359(6378):904–8
- Kyrou K, Hammond AM, Galizi R, Kranjc N, Burt A, et al. 2018. A CRISPR–Cas9 gene drive targeting *doublesex* causes complete population suppression in caged *Anopheles gambiae* mosquitoes. *Nat. Biotechnol.* 36(11):1062–66
- Lankau RA. 2011. Rapid evolutionary change and the coexistence of species. *Annu. Rev. Ecol. Syst.* 42:335–54
- Lankau R, Jørgensen PS, Harris DJ, Sih A. 2011. Incorporating evolutionary principles into environmental management and policy. *Evol. Appl.* 4(2):315–25
- Lau JA, Magnoli SM, Zirbel CR, Brudvig LA. 2019. The limits to adaptation in restored ecosystems and how management can help overcome them. *Ann. Mo. Bot. Gard.* 104(3):441–54
- Lebarbenchon C, Brown SP, Poulin R, Gauthier-Clerc M, Thomas F. 2008. Evolution of pathogens in a man-made world. *Mol. Ecol.* 17(1):475–84
- Lipinski KA, Barber LJ, Davies MN, Ashenden M, Sottoriva A, Gerlinger M. 2016. Cancer evolution and the limits of predictability in precision cancer medicine. *Trends Cancer.* 2(1):49–63
- Lipsitch M. 2018. Why do exceptionally dangerous gain-of-function experiments in influenza? In *Influenza Virus: Methods and Protocols*, ed. Y Yamauchi, pp. 589–608. New York: Springer
- Living with Resistance project. 2018. Antibiotic and pesticide susceptibility and the Anthropocene operating space. *Nat. Sustain.* 1(11):632–41
- Monfreda C, Ramankutty N, Foley JA. 2008. Farming the planet: 2. Geographic distribution of crop areas, yields, physiological types, and net primary production in the year 2000. *Glob. Biogeochem. Cycles* 22(1):1–19

- Morris DH, Gostic KM, Pompei S, Bedford T, Marta L, et al. 2018. Predictive modeling of influenza shows the promise of applied evolutionary biology. *Trends Microbiol.* 26(2):102–18
- Motta EVS, Raymann K, Moran NA. 2018. Glyphosate perturbs the gut microbiota of honey bees. *PNAS* 115(41):10305–10
- Noble C, Olejarz J, Esvelt KM, Church GM, Nowak MA. 2017. Evolutionary dynamics of CRISPR gene drives. *Sci. Adv.* 3(4):e1601964
- Palkovacs EP, Moritsch MM, Contolini GM, Pelletier F. 2018. Ecology of harvest-driven trait changes and implications for ecosystem management. *Front. Ecol. Environ.* 16(1):20–28
- Palumbi SR. 2001. Humans as the world's greatest evolutionary force. *Science* 293(5536):1786–90
- Parnell S, van den Bosch F, Gottwald T, Gilligan CA. 2017. Surveillance to inform control of emerging plant diseases: an epidemiological perspective. *Annu. Rev. Phytopathol.* 55:591–610
- Pearse DE. 2016. Saving the spandrels? Adaptive genomic variation in conservation and fisheries management. *J. Fish Biol.* 89(6):2697–716
- Piaggio AJ, Segelbacher G, Seddon PJ, Alphey L, Bennett EL, et al. 2017. Is it time for synthetic biodiversity conservation? *Trends Ecol. Evol.* 32(2):97–107
- Polasky S, Carpenter SR, Folke C, Keeler B. 2011. Decision-making under great uncertainty: environmental management in an era of global change. *Trends Ecol. Evol.* 26(8):398–404
- Raimundo RLG, Guimarães PR, Evans DM. 2018. Adaptive networks for restoration ecology. *Trends Ecol. Evol.* 33(9):664–75
- Rev. Antimicrob. Resist. 2016. *Tackling drug-resistant infections globally: final report and recommendations*. Rep., May 19, Rev. Antimicrob. Resist., London. https://amr-review.org/sites/default/files/160518_Final%20paper_with%20cover.pdf
- Reznick DN, Ghalambor CK. 2001. The population ecology of contemporary adaptations: what empirical studies reveal about the conditions that promote adaptive evolution. *Genetica* 112:183–98
- Ridley CE, Alexander LC. 2016. Applying gene flow science to environmental policy needs: a boundary work perspective. *Evol. Appl.* 9(7):924–36
- Robert A, Thévenin C, Princé K, Sarrazin F, Clavel J. 2017. De-extinction and evolution. *Funct. Ecol.* 31(5):1021–31
- Robertson BA, Rehage JS, Sih A. 2013. Ecological novelty and the emergence of evolutionary traps. *Trends Ecol. Evol.* 28(9):552–60
- Robinson TP, William Wint GR, Conchedda G, Van Boeckel TP, Ercoli V, et al. 2014. Mapping the global distribution of livestock. *PLOS ONE* 9(5):e96084
- Rouco C, Moreno S, Santoro S. 2016. A case of low success of blind vaccination campaigns against myxomatosis and rabbit haemorrhagic disease on survival of adult European wild rabbits. *Prev. Vet. Med.* 133:108–13
- Santamaría L, Méndez PF. 2012. Evolution in biodiversity policy—current gaps and future needs. *Evol. Appl.* 5(2):202–18
- Ségurel L, Bon C. 2017. On the evolution of lactase persistence in humans. *Annu. Rev. Genom. Hum. Genet.* 18:297–319
- Selbach C, Seddon PJ, Poulin R. 2018. Parasites lost: neglecting a crucial element in de-extinction. *Trends Parasitol.* 34(1):9–11
- Sgrò CM, Lowe AJ, Hoffmann AA. 2011. Building evolutionary resilience for conserving biodiversity under climate change. *Evol. Appl.* 4(2):326–37
- Sharma P, Hu-Lieskovan S, Wargo JA, Ribas A. 2017. Primary, adaptive, and acquired resistance to cancer immunotherapy. *Cell* 168(4):707–23
- Shaw RG. 2019. From the past to the future: considering the value and limits of evolutionary prediction. *Am. Nat.* 193(1):1–10
- Sih A, Ferrari MCO, Harris DJ. 2011. Evolution and behavioural responses to human-induced rapid environmental change. *Evol. Appl.* 4(2):367–87
- Singh V, Braddick D, Dhar PK. 2017. Exploring the potential of genome editing CRISPR-Cas9 technology. *Gene* 599:1–18

- Smith TB, Kinnison MT, Strauss SY, Fuller TL, Carroll SP. 2014. Prescriptive evolution to conserve and manage biodiversity. *Annu. Rev. Ecol. Evol. Syst.* 45:1–22
- Snieszko RA, Koch J. 2017. Breeding trees resistant to insects and diseases: putting theory into application. *Biol. Invasions* 19(11):1–24
- Steeves TE, Johnson JA, Hale ML. 2017. Maximising evolutionary potential in functional proxies for extinct species: a conservation genetic perspective on de-extinction. *Funct. Ecol.* 31(5):1032–40
- Strande NT, Berg JS. 2016. Defining the clinical value of a genomic diagnosis in the era of next-generation sequencing. *Annu. Rev. Genom. Hum. Genet.* 17:303–32
- Stroud JT, Losos JB. 2016. Ecological opportunity and adaptive radiation. *Annu. Rev. Ecol. Evol. Syst.* 47:507–32
- Thomas CD. 2015. Rapid acceleration of plant speciation during the Anthropocene. *Trends Ecol. Evol.* 30(8):448–55
- Toju H, Yamamichi M, Guimarães PR, Olesen JM, Mougi A, et al. 2017. Species-rich networks and eco-evolutionary synthesis at the metacommunity level. *Nat. Ecol. Evol.* 1(2):1–11
- Unckless RL, Clark AG, Messer PW. 2017. Evolution of resistance against CRISPR/Cas9 gene drive. *Genetics* 205(2):827–41
- Valiente-Banuet A, Aizen MA, Alcántara JM, Arroyo J, Cocucci A, et al. 2015. Beyond species loss: The extinction of ecological interactions in a changing world. *Funct. Ecol.* 29(3):299–307
- Van Boeckel TP, Brower C, Gilbert M, Grenfell BT, Levin SA, et al. 2015. Global trends in antimicrobial use in food animals. *PNAS* 112(18):5649–54
- van Oppen MJH, Oliver JK, Putnam HM, Gates RD. 2015. Building coral reef resilience through assisted evolution. *PNAS* 112(8):2307–13
- von Hertzen L, Beutler B, Bienenstock J, Blaser M, Cani PD, et al. 2015. Helsinki alert of biodiversity and health. *Ann. Med.* 47(3):218–25
- Vredenburg VT, Knapp RA, Tunstall TS, Briggs CJ. 2010. Dynamics of an emerging disease drive large-scale amphibian population extinctions. *PNAS* 107(21):9689–94
- Wells JCK, Nesse RM, Sear R, Johnstone RA, Stearns SC. 2017. Evolutionary public health: introducing the concept. *Lancet* 390(10093):500–9
- Williams M, Zalasiewicz J, Haff P, Schwägerl C, Barnosky AD, Ellis EC. 2015. The Anthropocene biosphere. *Anthropocene Rev.* 2(3):196–219
- Winter M, Devictor V, Schweiger O. 2013. Phylogenetic diversity and nature conservation: Where are we? *Trends Ecol. Evol.* 28(4):199–204
- Woods RJ, Read AF. 2015. Clinical management of resistance evolution in a bacterial infection: a case study. *Evol. Med. Public Health* 2015(1):281–88
- Wright O, Stan G-B, Ellis T. 2013. Building-in biosafety for synthetic biology. *Microbiology* 159(7):1221–35
- Xiong X, Chen M, Lim WA, Zhao D, Qi LS. 2016. CRISPR/Cas9 for human genome engineering and disease research. *Annu. Rev. Genom. Hum. Genet.* 17:131–54



Contents

<i>AREES at 50: A Semicentennial Celebration</i> <i>Douglas J. Futuyma</i>	1
Cultural Evolution in Animals <i>Andrew Whiten</i>	27
Somatic Mutation and Evolution in Plants <i>Daniel J. Schoen and Stewart T. Schultz</i>	49
Beyond Reproductive Isolation: Demographic Controls on the Speciation Process <i>Michael G. Harvey, Sonal Singhal, and Daniel L. Rabosky</i>	75
An Integrative Framework for Understanding the Mechanisms and Multigenerational Consequences of Transgenerational Plasticity <i>Alison M. Bell and Jennifer K. Hellmann</i>	97
Origins and Assembly of Malesian Rainforests <i>Robert M. Kooyman, Robert J. Morley, Darren M. Crayn,</i> <i>Elizabeth M. Joyce, Maurizio Rossetto, J.W. Ferry Slik,</i> <i>Joeri S. Strijk, Tao Su, Jia-Yee S. Yap, and Peter Wilf</i>	119
More Than the Sum of Its Parts: Microbiome Biodiversity as a Driver of Plant Growth and Soil Health <i>Mubammad Saleem, Jie Hu, and Alexandre Jousset</i>	145
Consequences of Multispecies Introductions on Island Ecosystems <i>James C. Russell and Christopher N. Kaiser-Bunbury</i>	169
Importance of Pollinator-Mediated Interspecific Pollen Transfer for Angiosperm Evolution <i>Juan Isaac Moreira-Hernández and Nathan Muchhala</i>	191
Haploid Selection in “Diploid” Organisms <i>Simone Immler</i>	219
Mycorrhizal Fungi as Mediators of Soil Organic Matter Dynamics <i>Serita D. Frey</i>	237

What Have Long-Term Field Studies Taught Us About Population Dynamics?	
<i>Beth A. Reinke, David A.W. Miller, and Fredric J. Janzen</i>	261
History and Geography of Neotropical Tree Diversity	
<i>Christopher W. Dick and R. Toby Pennington</i>	279
Climate Change in the Tropics: Ecological and Evolutionary Responses at Low Latitudes	
<i>Kimberly S. Sheldon</i>	303
Experimental Studies of Evolution and Eco-Evo Dynamics in Guppies (<i>Poecilia reticulata</i>)	
<i>David N. Reznick and Joseph Travis</i>	335
The Invasion Hierarchy: Ecological and Evolutionary Consequences of Invasions in the Fossil Record	
<i>Alycia L. Stigall</i>	355
Interactive Effects of Global Change on Forest Pest and Pathogen Dynamics	
<i>Allison B. Simler-Williamson, David M. Rizzo, and Richard C. Cobb</i>	381
Phylogenetic Comparative Methods and the Evolution of Multivariate Phenotypes	
<i>Dean C. Adams and Michael L. Collyer</i>	405
Spatial Population Genetics: It's About Time	
<i>Gideon S. Bradburd and Peter L. Ralph</i>	427
Evolutionary and Ecological Consequences of Gut Microbial Communities	
<i>Nancy A. Moran, Howard Ochman, and Tobin J. Hammer</i>	451
A Bird's-Eye View of Pollination: Biotic Interactions as Drivers of Adaptation and Community Change	
<i>Anton Pauw</i>	477
Life Ascending: Mechanism and Process in Physiological Adaptation to High-Altitude Hypoxia	
<i>Jay F. Storz and Graham R. Scott</i>	503
Evolution in the Anthropocene: Informing Governance and Policy	
<i>Peter Sogaard Jørgensen, Carl Folke, and Scott P. Carroll</i>	527
Revisiting the Fate of Dead Leaves That Fall into Streams	
<i>Jane C. Marks</i>	547
The Paradox Behind the Pattern of Rapid Adaptive Radiation: How Can the Speciation Process Sustain Itself Through an Early Burst?	
<i>Christopher H. Martin and Emilie J. Richards</i>	569

Related Articles

From the *Annual Review of Animal Biosciences*, Volume 7 (2019)

Functional Annotation of Animal Genomes (FAANG): Current Achievements
and Roadmap

Elisabetta Giuffra, Christopher K. Tuggle, and The FAANG Consortium

Mammalian Sex Chromosome Structure, Gene Content, and Function in Male
Fertility

Wan-Sheng Liu

Multiple Facets of Marine Invertebrate Conservation Genomics

Jose V. Lopez, Bisboy Kamel, Mónica Medina, Timothy Collins, and Iliana B. Baums

The Role of Reproductive Technologies in Amphibian Conservation Breeding
Programs

Aimee J. Silla and Phillip G. Byrne

Tigers of the World: Genomics and Conservation

Shu-Jin Luo, Yue-Chen Liu, and Xiao Xu

From the *Annual Review of Earth and Planetary Sciences*, Volume 47 (2019)

Seawater Chemistry Through Phanerozoic Time

Alexandra V. Turchyn and Donald J. DePaolo

Flood Basalts and Mass Extinctions

Matthew E. Clapham and Paul R. Renne

Soil Functions: Connecting Earth's Critical Zone

*Steven A. Banwart, Nikolaos P. Nikolaidis, Yong-Guan Zhu, Caroline L. Peacock,
and Donald L. Sparks*

Marsh Processes and Their Response to Climate Change and Sea-Level Rise

Duncan M. FitzGerald and Zoe Hughes

The Mesozoic Biogeographic History of Gondwanan Terrestrial Vertebrates:
Insights from Madagascar's Fossil Record

*David W. Krause, Joseph J.W. Sertich, Patrick M. O'Connor, Kristina Curry Rogers,
and Raymond R. Rogers*

Droughts, Wildfires, and Forest Carbon Cycling: A Pantropical Synthesis
*Paulo M. Brando, Lucas Paolucci, Caroline C. Ummenhofer, Elsa M. Ordway,
Henrik Hartmann, Megan E. Cattau, Ludmila Rattis, Vincent Medjibe,
Michael T. Coe, and Jennifer Balch*

From the *Annual Review of Environment and Resources*, Volume 44 (2019)

The State and Future of Antarctic Environments in a Global Context
Steven L. Chown and Cassandra M. Brooks

Island Biodiversity in the Anthropocene
James C. Russell and Christoph Kueffer

Mammal Conservation: Old Problems, New Perspectives, Transdisciplinarity,
and the Coming of Age of Conservation Geopolitics
David W. Macdonald

The State of the World's Mangrove Forests: Past, Present, and Future
*Daniel A. Friess, Kerrylee Rogers, Catherine E. Lovelock, Ken W. Krauss,
Stuart E. Hamilton, Shing Yip Lee, Richard Lucas, Jurgenne Primavera,
Anusha Rajkaran, and Subua Shi*

Status, Institutions, and Prospects for Global Capture Fisheries
Christopher Costello and Daniel Ovando

Illegal Wildlife Trade: Scale, Processes, and Governance
*Michael 't Sas-Rolfes, Daniel W.S. Challender, Amy Hinsley, Diogo Veríssimo,
and E. J. Milner-Gulland*

Ecotourism for Conservation?
Amanda L. Stronza, Carter A. Hunt, and Lee A. Fitzgerald

Co-Producing Sustainability: Reordering the Governance of Science, Policy,
and Practice
*Carina Wyborn, Amber Datta, Jasper Montana, Melanie Ryan, Peat Leith,
Brian Chaffin, Clark Miller, and Lorrae van Kerkhoff*

Social Synergies, Tradeoffs, and Equity in Marine Conservation Impacts
*David A. Gill, Samantha H. Cheng, Louise Glew, Ernest Aigner, Nathan J. Bennett,
and Michael B. Mascia*

From the *Annual Review of Entomology*, Volume 64 (2019)

The Ecology of Collective Behavior in Ants
Deborah M. Gordon

Invasion Success and Management Strategies for Social *Vespula* Wasps
Philip J. Lester and Jacqueline R. Beggs

Invasive Cereal Aphids of North America: Ecology and Pest Management
Michael J. Brewer, Frank B. Peairs, and Norman C. Elliott

Movement and Demography of At-Risk Butterflies: Building Blocks for Conservation

Cheryl B. Schultz, Nick M. Haddad, Erica H. Henry, and Elizabeth E. Crone

Epigenetics in Insects: Genome Regulation and the Generation of Phenotypic Diversity

Karl M. Glastad, Brendan G. Hunt, and Michael A.D. Goodisman

Molecular Evolution of the Major Arthropod Chemoreceptor Gene Families

Hugh M. Robertson

Systematics, Phylogeny, and Evolution of Braconid Wasps: 30 Years of Progress

Xue-xin Chen and Cornelis van Achterberg

Water Beetles as Models in Ecology and Evolution

David T. Bilton, Ignacio Ribera, and Andrew Edward Z. Short

Phylogeography of Ticks (Acari: Ixodida)

Lorenza Beati and Hans Klompen

From the *Annual Review of Genetics*, Volume 53 (2019)

Crossover Interference: Shedding Light on the Evolution of Recombination

Sarah P. Otto and Bret A. Payseur

Evolutionary Ecology of *Wolbachia* Releases for Disease Control

Perran A. Ross, Michael Turelli, and Ary A. Hoffmann

Living with Two Genomes: Grafting and Its Implications for Plant Genome-to-Genome Interactions, Phenotypic Variation, and Evolution

Brandon S. Gaut, Allison J. Miller, and Danelle K. Seymour

Standard Deviations: The Biological Bases of Transmission Ratio Distortion

Lila Fishman and Mariah McIntosh

The Microbiome and Aging

Bianca Bana and Filipe Cabreiro

Zebrafish Pigment Pattern Formation: Insights into the Development and Evolution of Adult Form

Larissa B. Patterson and David M. Parichy

From the *Annual Review of Genomics and Human Genetics*, Volume 20 (2019)

The Genetics of Human Skin and Hair Pigmentation

William J. Pavan and Richard A. Sturm

Measuring Clonal Evolution in Cancer with Genomics

Marc J. Williams, Andrea Sottoriva, and Trevor A. Graham

The Causes and Consequences of Genetic Interactions (Epistasis)

Júlia Domingo, Pablo Baeza-Centurion, and Ben Lebner

Thinking About the Evolution of Complex Traits in the Era of Genome-Wide Association Studies

Guy Sella and Nicholas H. Barton

From the *Annual Review of Marine Science*, Volume 11 (2019)

Planktonic Marine Archaea

Alyson E. Santoro, R. Alexander Richter, and Christopher L. Dupont

Arctic and Antarctic Sea Ice Change: Contrasts, Commonalities, and Causes

Ted Maksym

Biologically Generated Mixing in the Ocean

Eric Kunze

Climate Change, Coral Loss, and the Curious Case of the Parrotfish Paradigm: Why Don't Marine Protected Areas Improve Reef Resilience?

John F. Bruno, Isabelle M. Côté, and Lauren T. Toth

Marine Metazoan Modern Mass Extinction: Improving Predictions by Integrating Fossil, Modern, and Physiological Data

Piero Calosi, Hollie M. Putnam, Richard J. Twitchett, and Fanny Vermandele

From the *Annual Review of Microbiology*, Volume 73 (2019)

Ecology and Evolution of Plant Microbiomes

Viviane Cordovez, Francisco Dini-Andreote, Víctor J. Carrión, and Jos M. Raaijmakers

Algal Sex Determination and the Evolution of Anisogamy

James Umen and Susana Coelbo

The Ultimate Guide to Bacterial Swarming: An Experimental Model to Study the Evolution of Cooperative Behavior

Jinyuan Yan, Hilary Monaco, and Joao B. Xavier

Biogeography of the Oral Microbiome: The Site-Specialist Hypothesis

Jessica L. Mark Welch, Floyd E. Dewhirst, and Gary G. Boris

Diversity, Genomics, and Distribution of Phytoplankton-Cyanobacterium Single-Cell Symbiotic Associations

Rachel A. Foster and Jonathan P. Zehr

Paleomicrobiology: Diagnosis and Evolution of Ancient Pathogens

Kirsten I. Bos, Denise Kühnert, Alexander Herbig, Luis Roger Esquivel-Gomez, Aida Andrades Valtueña, Rodrigo Barquera, Karen Giffin, Aditya Kumar Lankapalli, Elizabeth A. Nelson, Susanna Sabin, Maria A. Spyrou, and Johannes Krause

From the *Annual Review of Plant Biology*, Volume 70 (2019)

Molecular Interactions Between Plants and Insect Herbivores

Matthias Erb and Philippe Reymond

A Molecular View of Plant Local Adaptation: Incorporating Stress-Response Networks

Acer VanWallendael, Ali Soltani, Nathan C. Emery, Murilo M. Peixoto, Jason Olsen, and David B. Lowry

Comparative and Functional Algal Genomics

Crysten E. Blaby-Haas and Sabeeha S. Merchant

CRISPR/Cas Genome Editing and Precision Plant Breeding in Agriculture

Kunling Chen, Yanpeng Wang, Rui Zhang, Huawei Zhang, and Caixia Gao

Risk Assessment and Regulation of Plants Modified by Modern Biotechniques: Current Status and Future Challenges

Joachim Schiemann, Antje Dietz-Pfeilstetter, Frank Hartung, Christian Kohl, Jörg Romeis, and Thorben Sprink

From the *Annual Review of Phytopathology*, Volume 57 (2019)

Revisiting the Concept of Host Range of Plant Pathogens

Cindy E. Morris and Benoît Moury

Durability of Quantitative Resistance in Crops: Greater Than We Know?

Christina Cowger and James K.M. Brown

Molecular Dialog Between Parasitic Plants and Their Hosts

Christopher R. Clarke, Michael P. Timko, John I. Yoder, Michael J. Axtell, and James H. Westwood

Ecology and Evolution of the Sudden Oak Death Pathogen *Phytophthora ramorum*

Niklaus J. Grünwald, Jared M. LeBoldus, and Richard C. Hamelin

Understanding Adaptation, Coevolution, Host Specialization, and Mating System in Castrating Anther-Smut Fungi by Combining Population and Comparative Genomics

Fanny E. Hartmann, Ricardo C. Rodríguez de la Vega, Fantin Carpentier, Pierre Gladieux, Amandine Cornille, Michael E. Hood, and Tatiana Giraud

Surviving in a Hostile World: Plant Strategies to Resist Pests and Diseases

Samuel W. Wilkinson, Melissa H. Magerøy, Ana López Sánchez, Lisa M. Smith, Leonardo Furci, T.E. Anne Cotton, Paal Krokene, and Jurriaan Ton