#### Four.II Geometry of Determinants


Linear Algebra
Jim Hefferon

http://joshua.smcvt.edu/linearalgebra

# Determinants as size functions

#### Box


This parallelogram is defined by the two vectors.



1.1 Definition In  $\mathbb{R}^n$  the box (or parallelepiped) formed by  $\langle \vec{v}_1, \dots, \vec{v}_n \rangle$  is the set  $\{t_1 \vec{v}_1 + \dots + t_n \vec{v}_n \mid t_1, \dots, t_n \in [0 \dots 1]\}$ .


The box is like the span, except that the scalars are from the unit interval.

#### Area of a two dimensional box



box area = rectangle area - area of 
$$A - \cdots$$
 - area of  $F$   
=  $(x_1 + x_2)(y_1 + y_2) - x_2y_1 - x_1y_1/2$   
 $- x_2y_2/2 - x_2y_2/2 - x_1y_1/2 - x_2y_1$   
=  $x_1y_2 - x_2y_1$ 

#### Area of a two dimensional box



box area = rectangle area - area of 
$$A - \cdots$$
 - area of  $F$   
=  $(x_1 + x_2)(y_1 + y_2) - x_2y_1 - x_1y_1/2$   
 $- x_2y_2/2 - x_2y_2/2 - x_1y_1/2 - x_2y_1$   
=  $x_1y_2 - x_2y_1$ 

That area equals the value of the determinant.

$$\begin{vmatrix} x_1 & x_2 \\ y_1 & y_2 \end{vmatrix} = x_1 y_2 - x_2 y_1$$

We will argue here that the properties of determinants make good postulates for a function that gives the size of the box formed by the columns of the matrix.

We will argue here that the properties of determinants make good postulates for a function that gives the size of the box formed by the columns of the matrix.

In line with the above use of column vectors, in this section we consider the determinant as a function of the columns of the matrix. Note that because the determinant of a matrix equals the determinant of its transpose, we can change the determinant properties from statements about rows to statements about columns. For instance, the first property as given earlier says that a determinant is unaffected by a row operation  $k\rho_i+\rho_j$  (with  $i\neq j$ ). By transposing we have the property that a determinant does not change under combinations of columns.

## Definition of determinant reinterpreted

Recall property (3) from the definition of determinant, that rescaling a column rescales the entire determinant  $\det(\ldots,k\vec{v}_i,\ldots)=k\det(\ldots,\vec{v}_i,\ldots)$ . This fits with the idea that the determinant gives the size of the box formed by the columns of the matrix: if we scale a column by a factor k then the size of the box scales by that factor.




# Definition of determinant reinterpreted

Recall property (3) from the definition of determinant, that rescaling a column rescales the entire determinant  $\det(\dots,k\vec{v}_i,\dots)=k\det(\dots,\vec{v}_i,\dots)$ . This fits with the idea that the determinant gives the size of the box formed by the columns of the matrix: if we scale a column by a factor k then the size of the box scales by that factor.



Property (1) says that the determinant is unaffected by combining columns. The picture



shows that the box formed by  $\vec{v}$  and  $k\vec{v} + \vec{w}$  is more slanted than the original box but the two have the same base and height, and hence the same area.


As we noted after the definition, property (2) is a consequence of the other properties so we leave it aside for the moment.


Property (4) says that the determinant of the identity matrix is 1.



## Aside: Orientation in two space

1.2 Remark Although property (2) is redundant, it says something notable. Consider these two.







Swapping changes the sign; on the left we take  $\vec{u}$  first in the matrix and then follow the counterclockwise arc to  $\vec{v}$ , following the counterclockwise arc and and get a positive size, while on the right following the clockwise arc gives a negative size. The sign returned by the size function, the determinant, reflects the *orientation* or *sense* of the box.

## Aside: Orientation in two space

1.2 Remark Although property (2) is redundant, it says something notable. Consider these two.





Swapping changes the sign; on the left we take  $\vec{u}$  first in the matrix and then follow the counterclockwise arc to  $\vec{v}$ , following the counterclockwise arc and and get a positive size, while on the right following the clockwise arc gives a negative size. The sign returned by the size function, the determinant, reflects the *orientation* or *sense* of the box. We see the same thing if we use Property (3) with a negative scalar.

## Aside: Orientation in three space

Starting with these two vectors we want to form a box with positive size.

$$\vec{v}_1 = \begin{pmatrix} 1 \\ 4 \\ 1 \end{pmatrix}, \vec{v}_2 = \begin{pmatrix} -2 \\ 3 \\ 1 \end{pmatrix}$$

$$\vec{v}_3$$

$$\vec{v}_1$$

Those two vectors span a plane, which divides three-space in two. The  $\vec{v}_3$  shown is on the side of the plane containing vectors with this property:

## Aside: Orientation in three space

Starting with these two vectors we want to form a box with positive size.


$$\vec{v}_1 = \begin{pmatrix} 1 \\ 4 \\ 1 \end{pmatrix}, \vec{v}_2 = \begin{pmatrix} -2 \\ 3 \\ 1 \end{pmatrix} \qquad \vec{v}_3$$

Those two vectors span a plane, which divides three-space in two. The  $\vec{v}_3$  shown is on the side of the plane containing vectors with this property: if a person at the tip of  $\vec{v}_3$  looks down at the box and traces out the parallelogram from  $\vec{0}$ , to  $\vec{v}_1$ , to  $\vec{v}_1 + \vec{v}_2$ , to  $\vec{v}_2$ , and back to  $\vec{0}$  then their trace looks to them to be counterclockwise. Any such vector will make a positive-sized box.

$$\vec{v}_3 = \begin{pmatrix} 0 \\ -1 \\ 2 \end{pmatrix} \qquad \begin{vmatrix} 1 & -2 & 0 \\ 4 & 3 & -1 \\ 1 & 1 & 2 \end{vmatrix} = 25$$


$$\vec{v}_3 = \begin{pmatrix} 0 \\ -1 \\ 2 \end{pmatrix} \qquad \begin{vmatrix} 1 & -2 & 0 \\ 4 & 3 & -1 \\ 1 & 1 & 2 \end{vmatrix} = 25$$

$$\begin{vmatrix} 1 & -2 & 0 \\ 4 & 3 & 1 \\ 1 & 1 & -2 \end{vmatrix} = -25$$




$$\vec{v}_3 = \begin{pmatrix} 0 \\ -1 \\ 2 \end{pmatrix} \qquad \begin{vmatrix} 1 & -2 & 0 \\ 4 & 3 & -1 \\ 1 & 1 & 2 \end{vmatrix} = 25$$

$$\begin{vmatrix} 1 & -2 & 0 \\ 4 & 3 & 1 \\ 1 & 1 & -2 \end{vmatrix} = -25$$




$$\vec{v}_3 = \begin{pmatrix} 0 \\ -1 \\ 2 \end{pmatrix} \qquad \begin{vmatrix} 1 & -2 & 0 \\ 4 & 3 & -1 \\ 1 & 1 & 2 \end{vmatrix} = 25$$

$$\begin{vmatrix} 1 & -2 & 0 \\ 4 & 3 & 1 \\ 1 & 1 & -2 \end{vmatrix} = -25$$




$$\vec{v}_3 = \begin{pmatrix} 0 \\ -1 \\ 2 \end{pmatrix} \qquad \begin{vmatrix} 1 & -2 & 0 \\ 4 & 3 & -1 \\ 1 & 1 & 2 \end{vmatrix} = 25$$

$$\begin{vmatrix} 1 & -2 & 0 \\ 4 & 3 & 1 \\ 1 & 1 & -2 \end{vmatrix} = -25$$




$$\vec{v}_3 = \begin{pmatrix} 0 \\ -1 \\ 2 \end{pmatrix} \qquad \begin{vmatrix} 1 & -2 & 0 \\ 4 & 3 & -1 \\ 1 & 1 & 2 \end{vmatrix} = 25$$

$$\begin{vmatrix} 1 & -2 & 0 \\ 4 & 3 & 1 \\ 1 & 1 & -2 \end{vmatrix} = -25$$




$$\vec{v}_3 = \begin{pmatrix} 0 \\ -1 \\ 2 \end{pmatrix} \qquad \begin{vmatrix} 1 & -2 & 0 \\ 4 & 3 & -1 \\ 1 & 1 & 2 \end{vmatrix} = 25$$

$$\begin{vmatrix} 1 & -2 & 0 \\ 4 & 3 & 1 \\ 1 & 1 & -2 \end{vmatrix} = -25$$



$$\vec{v}_3 = \begin{pmatrix} 0 \\ -1 \\ 2 \end{pmatrix} \qquad \begin{vmatrix} 1 & -2 & 0 \\ 4 & 3 & -1 \\ 1 & 1 & 2 \end{vmatrix} = 25$$

$$\begin{vmatrix} 1 & -2 & 0 \\ 4 & 3 & 1 \\ 1 & 1 & -2 \end{vmatrix} = -25$$



#### Determinants are multiplicative

.5 Theorem A transformation  $t \colon \mathbb{R}^n \to \mathbb{R}^n$  changes the size of all boxes by the same factor, namely the size of the image of a box |t(S)| is |T| times the size of the box |S|, where T is the matrix representing t with respect to the standard basis.

That is, the determinant of a product is the product of the determinants  $|TS| = |T| \cdot |S|$ .

#### Determinants are multiplicative

.5 Theorem A transformation  $t: \mathbb{R}^n \to \mathbb{R}^n$  changes the size of all boxes by the same factor, namely the size of the image of a box |t(S)| is |T| times the size of the box |S|, where T is the matrix representing t with respect to the standard basis.

That is, the determinant of a product is the product of the determinants  $|TS| = |T| \cdot |S|$ .

**Proof** First consider the |T| = 0 case, the case that T is singular and does not have an inverse. Observe that if TS is invertible then there is an M such that (TS)M = I, so T(SM) = I, and so T is invertible. The contrapositive of that observation is: if T is not invertible then neither is TS - if |T| = 0 then |TS| = 0.

#### Determinants are multiplicative

.5 Theorem A transformation  $t: \mathbb{R}^n \to \mathbb{R}^n$  changes the size of all boxes by the same factor, namely the size of the image of a box |t(S)| is |T| times the size of the box |S|, where T is the matrix representing t with respect to the standard basis.

That is, the determinant of a product is the product of the determinants  $|TS| = |T| \cdot |S|$ .

**Proof** First consider the |T| = 0 case, the case that T is singular and does not have an inverse. Observe that if TS is invertible then there is an M such that (TS)M = I, so T(SM) = I, and so T is invertible. The contrapositive of that observation is: if T is not invertible then neither is TS - if |T| = 0 then |TS| = 0.

Now consider the case that T is nonsingular. Any nonsingular matrix factors into a product of elementary matrices  $T = E_1 E_2 \cdots E_r$ . To finish this argument we will verify that  $|ES| = |E| \cdot |S|$  for all matrices S and elementary matrices E. The result will then follow because  $|TS| = |E_1 \cdots E_r S| = |E_1| \cdots |E_r| \cdot |S| = |E_1 \cdots E_r| \cdot |S| = |T| \cdot |S|$ .

There are three types of elementary matrix. We will cover the  $M_i(k)$  case; the  $P_{i,j}$  and  $C_{i,j}(k)$  checks are similar. The matrix  $M_i(k)S$  equals S except that row i is multiplied by k. The third condition of determinant functions then gives that  $|M_i(k)S| = k \cdot |S|$ . But  $|M_i(k)| = k$ , again by the third condition because  $M_i(k)$  is derived from the identity by multiplication of row i by k. Thus  $|ES| = |E| \cdot |S|$  holds for  $E = M_i(k)$ .

*Example* The transformation  $t_{\theta} \colon \mathbb{R}^2 \to \mathbb{R}^2$  that rotates all vectors through a counterclockwise angle  $\theta$  is represented by this matrix.

$$\mathsf{T}_{\theta} = \mathsf{Rep}_{\mathcal{E}_2}(\mathsf{t}_{\theta}) = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$$

Observe that  $t_{\theta}$  doesn't change the size of any boxes, it just rotates the entire box as a rigid whole. Note that  $|T_{\theta}| = 1$ .

*Example* The transformation  $t_{\theta} \colon \mathbb{R}^2 \to \mathbb{R}^2$  that rotates all vectors through a counterclockwise angle  $\theta$  is represented by this matrix.

$$T_{\theta} = \operatorname{Rep}_{\mathcal{E}_{2}}(t_{\theta}) = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$$


Observe that  $t_{\theta}$  doesn't change the size of any boxes, it just rotates the entire box as a rigid whole. Note that  $|T_{\theta}| = 1$ .

*Example* The linear transformation  $s: \mathbb{R}^2 \to \mathbb{R}^2$  represented with respect to the standard basis by this matrix

$$S = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$$


will, by the theorem, change the size of a box by a factor of |S| = -2. Here is s acting on a typical box.

The box defined by the two vectors  $\vec{v}_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$  and  $\vec{v}_2 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$  is transformed by s to the box defined by the two vectors  $s(\vec{v}_1) = \begin{pmatrix} 1 \\ 3 \end{pmatrix}$  and  $s(\vec{v}_2) = \begin{pmatrix} 3 \\ 7 \end{pmatrix}$ .



Note the change in orientation.

The box defined by the two vectors  $\vec{v}_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$  and  $\vec{v}_2 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$  is transformed by s to the box defined by the two vectors  $s(\vec{v}_1) = \begin{pmatrix} 1 \\ 3 \end{pmatrix}$  and  $s(\vec{v}_2) = \begin{pmatrix} 3 \\ 7 \end{pmatrix}$ .



Note the change in orientation.

The two sizes are easy.

$$\begin{vmatrix} 1 & 1 \\ 0 & 1 \end{vmatrix} = 1 \qquad \begin{vmatrix} 1 & 3 \\ 3 & 7 \end{vmatrix} = -2$$

# Determinant of the inverse is the inverse of the determinant

1.7 Corollary If a matrix is invertible then the determinant of its inverse is the inverse of its determinant  $|T^{-1}| = 1/|T|$ .

# Determinant of the inverse is the inverse of the determinant

1.7 Corollary If a matrix is invertible then the determinant of its inverse is the inverse of its determinant  $|T^{-1}| = 1/|T|$ .

*Proof* 
$$1 = |I| = |TT^{-1}| = |T| \cdot |T^{-1}|$$

QED

#### Volume

1.3 *Definition* The *volume* of a box is the absolute value of the determinant of a matrix with those vectors as columns.

Example The box formed by thevectors

$$\vec{v}_1 = \begin{pmatrix} -1\\1 \end{pmatrix} \quad \vec{v}_2 = \begin{pmatrix} 1\\1 \end{pmatrix}$$

gives this determinant

$$\begin{vmatrix} -1 & 1 \\ 1 & 1 \end{vmatrix} = -2$$

so its volume is 2.

