
Holding: all archived pre-peak ranges


Corrected 10-second midpoint results • all 7 cutoff-available scenes • absolute D; no imputed baseline • times UTC+08

Order 1 · Order 2 · Order 3 · Order 4 · Order 5 · Mean of available orders · × Rejected D fit (bottom marks)


A Sep 01, 09:00 | N = 32
archived range 0–22; median order ρ = 0.898




B Sep 04, 16:00 | N = 40
archived range 0–32; median order ρ = 0.828




C Sep 10, 14:00 | N = 36
archived range 0–20; median order ρ = 0.845




D Sep 15, 14:00 | N = 46
archived range 0–21; median order ρ = 0.850




E Sep 20, 12:00 | N = 33
archived range 0–17; median order ρ = 0.830




F Sep 26, 07:00 | N = 31
archived range 0–7; median order ρ = 0.762



G Sep 28, 15:00 | N = 34
archived range 0–16; median order ρ = 0.901



H Every retained-range coefficient
32/33 positive; median ρ = 0.845



Hour-based source selection; complete injected tails retained. Historical cutoffs are outcome-selected; no claim of a validated operating limit.

All 12 planned scenes and full ranges are retained in the report. Four displayed scenes have no accepted baseline; these panels show absolute D.

Panel y-scales differ. Available-order means are descriptive; order correlations use each order's valid doses. Missing values break lines; no smoothing.

$R^2 \geq 0.98$ unchanged. Five orders share the same aircraft and are not independent scenes. Equal-duration control remains separate evidence.