Vectoring dose response | all four scenes and both saved 15 km shapes
Accepted D on a common y-axis; no truncation or scene selection



09-14 14:00 UTC+08 | N=38
3 candidates | D fits: 20/20
rho NA: only 4 doses (<5 required)

09-15 15:00 UTC+08 | N=47
6 candidates | D fits: 35/35
Median order rho=0.964; positive 5/5

09-19 15:00 UTC+08 | N=40
5 candidates | D fits: 30/30
Median order rho=0.714; positive 4/5

09-27 13:00 UTC+08 | N=32
5 candidates | D fits: 30/30
Median order rho=0.600; positive 4/5



09-14 14:00 UTC+08 | N=38
3 candidates | D fits: 20/20
rho NA: only 4 doses (<5 required)

09-15 15:00 UTC+08 | N=47
6 candidates | D fits: 35/35
Median order rho=0.929; positive 5/5

09-19 15:00 UTC+08 | N=40
5 candidates | D fits: 30/30
Median order rho=-0.486; positive 2/5

09-27 13:00 UTC+08 | N=32
5 candidates | D fits: 30/30
Median order rho=-0.086; positive 2/5



All 115 D fits per shape pass $R^2 \geq 0.94$; the first scene lacks enough dose levels for correlation, not for D. Negative-order responses remain visible.
Segmented: schematic stimulus without the smooth motion certificate. Smooth: screened control. Five orders are dependent, not independent scenes.