



Elementary First Aid (EFA)

90 Practice Questions and Answers for STCW Exams

Updated for DG Shipping India Exams | Source: brightmariner.com

Practice Set 1

Q1. What is called the voice box?

- A) Larynx
- B) Trachea
- C) Epiglottis
- D) None of the above

Correct Answer: A (Larynx)

Explanation: The larynx, commonly known as the voice box, contains the vocal cords and is responsible for producing sound and speech.

Q2. In topographic anatomy, the term lateral means:

- A) Towards the midline of the body
- B) Away from the midline of the body
- C) Above the midline of the body
- D) Below the midline of the body

Correct Answer: B (Away from the midline of the body)

Explanation: In topographic anatomy, the term 'lateral' refers to a position away from the midline of the body or toward the side.

Q3. Where does the partly digested food in liquid form go after it leaves the stomach?

- A) Small intestine
- B) Large intestine
- C) Liver
- D) Pancreas

Correct Answer: A (Small intestine)

Explanation: After partially digesting in the stomach, food moves into the small intestine where most of the digestion and nutrient absorption occurs.

Q4. Urine enters the bladder through which structure?

- A) Urethra
- B) Ureter
- C) Nephron
- D) Bowman's capsule

Correct Answer: B (Ureter)

Explanation: Urine travels from the kidneys down to the bladder through tubes called ureters.



- A) Sympathetic nervous system
- B) Parasympathetic nervous system
- C) Central nervous system
- D) Endocrine system

Correct Answer: A (Sympathetic nervous system)

Explanation: The sympathetic nervous system is responsible for the 'fight-or-flight' response, preparing the body to react to perceived threats by increasing heart rate and diverting blood flow.

Q6. What is the correct sequence for the primary survey?

- A) D-R-A-B-C
- B) A-B-C-D-R
- C) R-A-B-C-D
- D) D-A-B-C-R

Correct Answer: A (D-R-A-B-C)

Explanation: The correct sequence for the primary survey in first aid is Danger, Response, Airway, Breathing, Circulation (DRABC).

Q7. Which of the following are considered crucial protective equipment?

- A) Gloves
- B) Mask
- C) Eye shield
- D) All of the above

Correct Answer: D (All of the above)

Explanation: Gloves, masks, and eye shields are all considered crucial protective equipment to prevent the transmission of infections and protect against bodily fluids.

Q8. You are taking blood pressure by palpation. A radial pulse indicates a blood pressure of at least:

- A) 60 mmHg
- B) 70 mmHg
- C) 80 mmHg
- D) 90 mmHg

Correct Answer: C (80 mmHg)

Explanation: When taking blood pressure by palpation, the presence of a radial pulse typically indicates a systolic blood pressure of at least 80 mmHg.

Q9. An unconscious seaman is lying on his back and not breathing. What is the first thing you do?

- A) Check for a pulse
- B) Give rescue breaths
- C) Tilt his head back as far as it will go
- D) Call for help

Correct Answer: C (Tilt his head back as far as it will go)

Explanation: When an unconscious seaman is not breathing, the first step to open the airway is to tilt the head back and lift the chin, ensuring the tongue is pulled away from the throat.



- A) Move the victim as quickly as possible
- B) Support the head and neck in the position found
- C) Try to straighten the victim's neck
- D) None of the above

Correct Answer: B (Support the head and neck in the position found)

Explanation: If a head or neck injury is suspected, it is crucial to support the head and neck in the position they are found to prevent further damage to the spinal cord.

Q11. When a victim is having a neck injury, you should:

- A) Hold his head still
- B) Move his head gently
- C) Try to align his head and neck
- D) None of the above

Correct Answer: A (Hold his head still)

Explanation: When a victim has a suspected neck injury, the primary action is to hold their head still to avoid exacerbating any spinal trauma.

Q12. You suspect a head and neck injury in a victim who is unresponsive and not breathing. How would you open the airway to give breaths?

- A) Tilt the head back and lift the chin
- B) Jaw thrust technique
- C) Use a suction device
- D) None of the above

Correct Answer: B (Jaw thrust technique)

Explanation: For a victim with suspected head or neck injury who is unresponsive and not breathing, the jaw thrust technique is used to open the airway, as it minimally disturbs the cervical spine.

Q13. Check the carotid artery for a pulse on:

- A) The side of his neck
- B) The wrist
- C) The groin
- D) The ankle

Correct Answer: A (The side of his neck)

Explanation: The carotid artery, used for checking a pulse in an unresponsive adult, is located on the side of the neck.

Q14. Fainting happens when:

- A) The heart stops beating
- B) The brain does not get enough blood with oxygen
- C) The person is dehydrated
- D) The person is anxious

Correct Answer: B (The brain does not get enough blood with oxygen)

Explanation: Fainting, or syncope, occurs when the brain temporarily does not receive enough oxygenated blood, leading to a brief loss of consciousness.



- A) Call 108 and return to the crew
- B) Give rescue breaths
- C) Check for a pulse
- D) None of the above

Correct Answer: A (Call 108 and return to the crew)

Explanation: If there is no response from the crew during the primary survey, the immediate next step is to call for emergency medical assistance (e.g., 108 or ship's medical officer) and then return to the victim.

Q16. Shock is the condition where:

- A) The body loses a lot of blood
- B) The person is dehydrated
- C) The cardiovascular system fails to deliver blood to all parts of the body
- D) The person is in pain

Correct Answer: C (The cardiovascular system fails to deliver blood to all parts of the body)

Explanation: Shock is a critical condition where the cardiovascular system fails to deliver adequate blood supply to all parts of the body, leading to a lack of oxygen in tissues.

Q17. What is the CPR procedure for a six-year-old boy?

- A) 30 compressions, 2 breaths
- B) 15 compressions, 2 breaths
- C) 15 compressions, 4 breaths
- D) 30 compressions, 4 breaths

Correct Answer: C (15 compressions, 4 breaths)

Explanation: For CPR on a child between puberty and adulthood (or a larger child), the recommended ratio is 15 compressions followed by 2 breaths.

Q18. What is the proper rate of chest compressions in CPR?

- A) 60 per minute
- B) 80 per minute
- C) 100 per minute
- D) 120 per minute

Correct Answer: C (100 per minute)

Explanation: The proper rate for chest compressions during CPR, for both adults and children, is 100 to 120 compressions per minute.

Q19. If an AED is available but does not have child pads or child energy setting, what should you do for a six-year-old non-responsive victim?

- A) Use the AED with adult pads
- B) Do not use the AED
- C) Call for help
- D) None of the above

Correct Answer: A (Use the AED with adult pads)

Explanation: If an AED is available without child pads or a child setting, it is recommended to use the adult pads for a child victim, ensuring they do not touch each other.



- A) Apply direct pressure
- B) Elevate the limb
- C) Apply a tourniquet
- D) Call for help

Correct Answer: A (Apply direct pressure)

Explanation: When managing external bleeding, the immediate action is to apply direct pressure to the wound to control blood loss.

Q21. For heavy bleeding, you need to apply pressure points to reduce bleeding.

- A) True
- B) False

Correct Answer: A (True)

Explanation: Applying pressure to specific points on arteries that run close to the bone helps to reduce blood flow to an injured area, which is a crucial technique for managing severe bleeding when direct pressure alone is insufficient.

Q22. You feel a sudden short pain in the bottom of your left foot. You look at the bottom of your left shoe and see a tack sticking in your shoe. What type of wound do you probably have?

- A) Abrasion
- B) Laceration
- C) Puncture
- D) Avulsion

Correct Answer: C (Puncture)

Explanation: A tack sticking into the foot through the shoe describes a puncture wound, which is caused by a sharp object penetrating the skin. This is distinct from abrasions (superficial scraping) or lacerations (tearing of tissue).

Q23. A tourniquet can be used for first resort to stop bleeding.

- A) True
- B) False

Correct Answer: B (False)

Explanation: A tourniquet is a last resort for life-threatening bleeding that cannot be controlled by other means. It should not be the first resort due to the risk of permanent damage, including nerve damage and tissue loss.

Q24. A tourniquet can be removed as and when necessary.

- A) True
- B) False

Correct Answer: B (False)

Explanation: A tourniquet should only be removed by trained medical personnel. Premature removal can lead to renewed severe bleeding and shock. Its application and removal must be carefully documented and managed.

Q25. What are splints made of?

- A) Wood
- B) Cardboard
- C) Clean plastic
- D) All of the above

Correct Answer: C (Clean plastic)



Q26. What should you do if you think a crew has internal bleeding?

- A) Give them something to drink
- B) Apply pressure to their abdomen
- C) Call for help
- D) None of the above

Correct Answer: C (Call for help)

Explanation: Internal bleeding is serious and requires immediate medical attention. Signs like severe pain, dizziness, or a rigid abdomen necessitate calling for professional medical help promptly; do not give the victim anything to eat or drink.

Q27. Burns that involve all layers of skin are:

- A) First-degree burns
- B) Second-degree burns
- C) Third-degree burns
- D) Full-thickness burns

Correct Answer: D (Full-thickness burns)

Explanation: Full-thickness burns, also known as third-degree burns, involve damage to all layers of the skin, and potentially underlying tissues. They are characterized by a white, leathery, or charred appearance and may be painless due to nerve damage.

Q28. For second-degree burns, you can break the blister and serve.

- A) True
- B) False

Correct Answer: B (False)

Explanation: Blisters should not be broken as they protect the underlying skin from infection. Keeping blisters intact is part of proper burn wound management to promote healing and reduce the risk of complications.

Q29. If 15% of the body is affected by second-degree burns, you should take the victim to the hospital.

- A) True
- B) False

Correct Answer: A (True)

Explanation: Second-degree burns covering 15% or more of the body surface area are considered severe and require immediate professional medical evaluation and hospital treatment to manage fluid loss and prevent infection.

Q30. First aid for frostbite is:

- A) Give some hot coffee
- B) Give some warm water
- C) Rub the affected area
- D) None of the above

Correct Answer: B (Give some warm water)

Explanation: For frostbite, the affected area should be gently rewarmed in warm (not hot) water, typically around 40 degreesC (104 degreesF). Rubbing the area can cause further damage, and hot drinks can be detrimental as they can redistribute blood flow away from extremities.



Practice Set 2

Q1. An unconscious person should not be placed in a position with the head lower than the body.

- A) Given an inhalation stimulant
- B) Given something to drink

Correct Answer: B (Given something to drink)

Explanation: An unconscious person should be placed in the recovery position to maintain an open airway and prevent aspiration if they vomit. Positioning the head lower than the body could impede circulation and breathing.

Q2. Which should not be a treatment for a person who has received a head injury and is groggy or unconscious?

- A) Given a stimulant
- B) Elevate his head
- C) Stop severe bleeding

Correct Answer: A (Given a stimulant)

Explanation: Stimulants should not be given to a person with a head injury, especially if they are groggy or unconscious, as they can mask symptoms or increase intracranial pressure. Focus should be on airway, breathing, circulation, and preventing further injury.

Q3. A person who gets battery acid in an eye should immediately wash the eye with:

- A) Boric acid solution
- B) Water
- C) Baking soda solution

Correct Answer: A (Boric acid solution)

Explanation: Battery acid is highly corrosive; immediate and copious irrigation with plain water is the first and most critical step. While a weak baking soda solution might be suggested for some acids, water is the universal first-aid treatment for chemical eye exposures.

Q4. A person who gets something in his or her eye and you see that it is not embedded, you can:

- A) Get them to rub their eye until the object is gone
- B) Remove it with a match or toothpick
- C) Remove it with a moist cotton-tipped applicator

Correct Answer: C (Remove it with a moist cotton-tipped applicator)

Explanation: If a foreign object is visible and not embedded in the eye, gently removing it with a moist cotton-tipped applicator is the safest method. Rubbing the eye or using sharp objects can cause further damage or embed the object deeper.

Q5. A victim is coughing and wheezing from a partial obstruction of the airway. An observer should:

- A) Perform the Heimlich maneuver
- B) Give back blows and something to drink
- C) Allow the person to continue coughing and dislodge the obstruction on his own

Correct Answer: C (Allow the person to continue coughing and dislodge the obstruction on his own)

Explanation: A person with a partial airway obstruction who is coughing and wheezing should be encouraged to continue coughing. This indicates they can still breathe and may be able to clear the obstruction themselves without intervention.

Q6. A shipmate chokes suddenly, cannot speak, and starts to turn blue. You should:

brightmariner.com



- A) Perform the Heimlich maneuver
- B) Make the victim lie down with the feet elevated to get blood to the brain
- C) Immediately administer CPR

Correct Answer: A (Perform the Heimlich maneuver)

Explanation: Sudden choking with inability to speak and cyanosis indicates a complete airway obstruction. The Heimlich maneuver (abdominal thrusts) is the immediate appropriate action to dislodge the object.

Q7. When starting CPR on a drowning victim, you should:

- A) Start chest compressions before the victim is removed from the water
- B) Drain water from the lungs before ventilating
- C) Begin mouth-to-mouth ventilations as soon as possible

Correct Answer: C (Begin mouth-to-mouth ventilations as soon as possible)

Explanation: For a drowning victim, it is crucial to begin rescue breathing as soon as possible to provide oxygen, even before removing them from the water if safe to do so. Chest compressions are initiated if no pulse is detected after ventilations.

Q8. When administering artificial respiration to an adult, the breathing cycle should be repeated about:

- A) 12 to 15 times per minute
- B) 18 to 20 times per minute
- C) As fast as possible

Correct Answer: A (12 to 15 times per minute)

Explanation: The recommended rate for artificial respiration in adults is about 12 to 15 breaths per minute, ensuring adequate oxygenation without causing hyperventilation or gastric distension.

Q9. If the person vomits during mouth-to-mouth resuscitation, the rescuer should first:

- A) Ignore it and continue mouth-to-mouth ventilation
- B) Switch to mouth-to-nose ventilation
- C) Turn the patient's body to the side, sweep out the mouth, and resume mouth-to-mouth resuscitation

Correct Answer: C (Turn the patient's body to the side, sweep out the mouth, and resume mouth-to-mouth resuscitation)

Explanation: If vomiting occurs during resuscitation, the rescuer must immediately turn the victim to their side, clear the mouth of vomit, and then resume CPR to prevent aspiration into the lungs.

Q10. Bleeding from a vein is:

- A) Dark red and has a steady flow
- B) Bright red and slow
- C) Bright red and squirting

Correct Answer: A (Dark red and has a steady flow)

Explanation: Venous bleeding is characterized by a dark red color and a steady, continuous flow, as veins carry deoxygenated blood back to the heart under lower pressure.



- A) Dark red with a steady flow
- B) Bright red with a steady flow
- C) Bright red and squirting

Correct Answer: C (Bright red and squirting)

Explanation: Arterial bleeding is bright red and typically spurts rhythmically with each heartbeat due to the high pressure within arteries.

Q12. The major cause of shock in burn victims is:

- A) The high level of pain
- B) Emotional stress
- C) Massive loss of fluid to the burnt area

Correct Answer: C (Massive loss of fluid to the burnt area)

Explanation: The primary cause of shock in burn victims is the significant loss of body fluid through the damaged skin, leading to hypovolemia and reduced circulating blood volume.

Q13. First, second, and third-degree burns are classified according to the:

- A) Area of the body involved
- B) Source of heat causing the burn
- C) Layers of skin affected

Correct Answer: C (Layers of skin affected)

Explanation: Burns are classified by depth based on the layers of skin affected: first-degree (epidermis), second-degree (epidermis and dermis), and third-degree (all skin layers and potentially deeper tissues).

Q14. Symptoms of heatstroke are:

- A) Cold and moist skin, high body temperature
- B) Cold and dry skin, low body temperature
- C) Hot and moist skin, high body temperature

Correct Answer: B (Cold and dry skin, low body temperature)

Explanation: Heatstroke is a medical emergency characterized by a high body temperature (often above 104 degreesF or 40 degreesC), hot and often dry skin (though it can be moist), and altered mental status. The skin is not typically cold in heatstroke.

Q15. A crew member suffering from hypothermia should be given:

- A) A small dose of alcohol
- B) Treatment for shock
- C) A large meal

Correct Answer: B (Treatment for shock)

Explanation: A victim of hypothermia needs gentle rewarming and management of shock. Alcohol can worsen hypothermia by causing vasodilation, and rubbing can cause further tissue damage.



- A) Revive the person with smelling salts
- B) Loosen the clothing
- C) Give pain reliever

Correct Answer: C (Give pain reliever)

Explanation: Pain relievers should not be given to a person who has fainted without medical advice, as they could mask symptoms or interact with potential underlying conditions. Focus should be on restoring circulation and ensuring airway patency.

Q17. First aid means:

- A) Medical treatment of an accident
- B) Setting of broken bones
- C) Emergency treatment at the scene of the injury

Correct Answer: C (Emergency treatment at the scene of the injury)

Explanation: First aid refers to the immediate, emergency care given to an injured or ill person before professional medical help arrives, aiming to preserve life and prevent further harm.

Q18. While administering artificial respiration, it is most important to:

- A) Monitor blood pressure
- B) Clear airways
- C) Use the rhythmic pressure method

Correct Answer: B (Clear airways)

Explanation: Maintaining a clear airway is the most critical step when administering artificial respiration, ensuring that air can reach the lungs effectively.

Q19. A shipmate suffers a heart attack and stops breathing. You must:

- A) Immediately give a stimulant by force if necessary
- B) Make the victim comfortable in a bunk
- C) Immediately start CPR

Correct Answer: C (Immediately start CPR)

Explanation: Sudden cardiac arrest where breathing stops requires immediate Cardiopulmonary Resuscitation (CPR) to maintain blood flow and oxygenation until professional help or an AED is available.

Q20. The best way to prevent a critical incident stress situation from becoming worse is to:

- A) Ignore it
- B) Be familiar with the signs and symptoms
- C) Ensure early use of proper medicine

Correct Answer: B (Be familiar with the signs and symptoms)

Explanation: Recognizing the signs and symptoms of critical incident stress is paramount for effective management. Early awareness allows for appropriate interventions and support, preventing the situation from escalating.



- A) Always use restraint
- B) Give the patient a choice in the treatment
- C) Play along with auditory or visual hallucinations

Correct Answer: C (Play along with auditory or visual hallucinations)

Explanation: When dealing with behavioral disorders, it's important not to argue with hallucinations. Playing along or acknowledging their perceived reality, while ensuring safety, can help de-escalate the situation and build trust.

Q22. You are taking blood pressure by palpation. A radial pulse indicates a blood pressure of at least:

- A) 110 mmHg
- B) 100 mmHg
- C) 90 mmHg

Correct Answer: A (110 mmHg)

Explanation: Palpating a radial pulse indicates that the systolic blood pressure is at least 80 mmHg, and it is generally considered to be at least 90 mmHg for a palpable femoral pulse and higher for other pulses.

Q23. The skeletal covering for the brain is provided by the bones of the cranium. The overall skeletal structure covering the spinal cord is the vertebral column or spine.

- A) True
- B) False

Correct Answer: A (True)

Explanation: This statement accurately describes the protective skeletal structures for the brain (cranium) and spinal cord (vertebral column), fundamental concepts in anatomy relevant to injury assessment.

Q24. You have tried to control a victim's forearm bleeding with direct pressure and elevation, but the bleeding does not stop. Where would you apply pressure?

- A) Outside the arm midway between the shoulder and the elbow
- B) On the inside of the elbow
- C) Inside the arm between the shoulder and elbow

Correct Answer: B (On the inside of the elbow)

Explanation: For forearm bleeding not controlled by direct pressure and elevation, the brachial artery pressure point is located on the inside of the arm, midway between the shoulder and the elbow. This point is compressed against the humerus bone.

Q25. What is the CPR procedure for a six-year-old boy?

- A) 15 compressions to 1 breath, 5 cycles in one minute
- B) 15 compressions to 2 breaths, 4 cycles in one minute
- C) 5 compressions to 1 breath, 12 cycles in one minute

Correct Answer: B (15 compressions to 2 breaths, 4 cycles in one minute)

Explanation: For a child aged puberty or younger, CPR guidelines recommend a compression-to-ventilation ratio of 30:2 for single rescuers and 15:2 for two rescuers, with approximately 5 cycles (about 2 minutes) per minute.



- A) The respiratory system fails to deliver air to the lungs
- B) The cardiovascular system fails to deliver blood to all parts of the body
- C) All of the above

Correct Answer: C (All of the above)

Explanation: Shock is a complex condition where the body's vital organs do not receive adequate blood flow. This can stem from various causes, including respiratory failure or a failing cardiovascular system.

Q27. If an AED is available but does not have child pads or child energy settings, what should you do for a six-year-old non-responsive victim?

- A) Perform CPR but do not use the AED
- B) Use the AED but only apply one pad
- C) Use the AED in adult mode

Correct Answer: A (Perform CPR but do not use the AED)

Explanation: If an AED is not equipped for pediatric use, it is safer to perform CPR and not use the AED at all for a child, as using adult settings or pads could deliver a potentially harmful electrical shock.

Q28. Irrigate an eye with:

- A) Diluted vinegar
- B) Saline solution
- C) An appropriate chemical antidote

Correct Answer: B (Saline solution)

Explanation: Irrigating an eye with a foreign object or chemical irritant should always be done with a clean, isotonic saline solution or plain water to avoid further irritation or damage.

Q29. In topographic anatomy, the term "lateral" means:

- A) Away from the midline of the body
- B) Nearer the head
- C) Away from the head

Correct Answer: A (Away from the midline of the body)

Explanation: In anatomical terms, 'lateral' refers to a position that is away from the midline or center of the body. Conversely, 'medial' means towards the midline.

Q30. The onset of shock in infants and children:

- A) May be sudden and severe
- B) Is no different than in adults
- C) Usually results from cardiac problems

Correct Answer: A (May be sudden and severe)

Explanation: Infants and children can develop shock more rapidly and severely than adults due to their smaller blood volume and less developed compensatory mechanisms. Early recognition and intervention are critical.

Practice Set 3



- A) Carotid pulse for 5-10 seconds
- B) Radial pulse for 5-10 seconds

Correct Answer: A (Carotid pulse for 5-10 seconds)

Explanation: The carotid pulse, located in the neck, is the preferred site for pulse assessment in adults during CPR or emergency assessment as it is generally more accessible and reliable than the radial pulse in compromised individuals. Assessing for 5-10 seconds is standard practice to confirm the presence or absence of a pulse.

Q2. What is the recommended amount of time to wash your hands?

- A) 20 seconds
- B) 30 seconds
- C) 40 seconds
- D) 5 minutes

Correct Answer: D (5 minutes)

Explanation: Washing hands for a minimum of 20 seconds is crucial for effectively removing transient microorganisms and preventing the spread of infections, aligning with public health recommendations for hand hygiene. Longer durations, like 30 or 40 seconds, may offer slightly better results, but 5 minutes is excessive for routine handwashing.

Q3. You can best control the swelling and pain of an ankle sprain by:

- A) Rest, immobilization, application of cold, and elevation (RICE)
- B) Applying a compression bandage
- C) Taking over-the-counter pain medication
- D) All of the above

Correct Answer: A (Rest, immobilization, application of cold, and elevation (RICE))

Explanation: The RICE protocol (Rest, Immobilization, Cold, Elevation) is the foundational treatment for sprains and strains to reduce swelling, pain, and inflammation. While compression bandages and pain medication can offer symptomatic relief, they are adjuncts to the primary RICE management strategy.

Q4. Mouth-to-nose respiration should be used only when:

- A) The victim is unconscious
- B) The victim has a severe head injury
- C) The victim has a broken jaw
- D) It is impossible to open the victim's mouth

Correct Answer: D (It is impossible to open the victim's mouth)

Explanation: Mouth-to-nose respiration is a crucial alternative technique indicated when the victim's mouth cannot be opened due to significant facial trauma, jaw injuries, or foreign body obstruction. This ensures ventilation is still possible, maintaining oxygenation until advanced medical care is available.

Q5. Toxic substances can affect the body in the following ways:

- A) Through the lungs
- B) Through the skin
- C) Through the eyes
- D) All of the above

Correct Answer: D (All of the above)

Explanation: Toxic substances can enter the body via multiple routes, including inhalation through the lungs, dermal absorption through the skin, and ingestion or direct contact with the eyes. Recognizing these entry points is vital for appropriate first aid and



~~Q6. What should you do if you think a crew member has internal bleeding?~~

- A) Monitor the victim's vital signs and seek medical attention if they worsen
- B) Apply a cold compress to the abdomen
- C) Give the victim pain medication
- D) Call your local emergency phone number for help

Correct Answer: D (Call your local emergency phone number for help)

Explanation: Internal bleeding is a serious condition that requires immediate professional medical intervention. While monitoring vital signs is important, definitive management necessitates seeking emergency medical services promptly rather than attempting home care like cold compresses or pain medication.

Q7. Which of the following is a sign of severe airway obstruction?

- A) Forceful cough
- B) Wheezing
- C) High-pitched inhalation noise
- D) All of the above

Correct Answer: D (All of the above)

Explanation: Severe airway obstruction is characterized by the inability to breathe effectively, leading to signs such as paradoxical breathing, high-pitched noises on inhalation (stridor), wheezing, and potentially cyanosis if oxygenation is compromised. A forceful cough can indicate a partial obstruction, but these other signs point to a more critical situation.

Q8. Infant CPR is administered to any child under the age of:

- A) 6 months
- B) 12 months
- C) 18 months
- D) 24 months

Correct Answer: B (12 months)

Explanation: Infant CPR guidelines typically define an infant as a child under one year of age, excluding newborns. This age distinction is important for applying appropriate techniques, such as two-finger chest compressions and ensuring correct airway positioning.

Q9. An atraumatic splint could be used for which of the following injuries?

- A) A mid-shaft femur fracture
- B) An ankle sprain
- C) A wrist fracture
- D) A rib fracture

Correct Answer: A (A mid-shaft femur fracture)

Explanation: An atraumatic splint, such as a Hare or Sager splint, is designed for immobilizing mid-shaft long bone fractures, particularly the femur, without causing further injury. Ankle sprains, wrist fractures, or rib fractures typically do not require this specialized type of splinting.



- A) Higher than the unaffected eye
- B) Lower than the unaffected eye
- C) At the same level as the unaffected eye
- D) It doesn't matter

Correct Answer: B (Lower than the unaffected eye)

Explanation: When flushing a chemical from the eye, holding the affected eye lower than the unaffected one allows the contaminated water to drain away from the unaffected eye. This prevents cross-contamination and ensures thorough irrigation of the exposed eye.

Q11. You find an adult who is unresponsive and not breathing. The scene is safe. What is your next step?

- A) Check for a pulse
- B) Activate the emergency response system
- C) Perform CPR
- D) Administer rescue breaths

Correct Answer: B (Activate the emergency response system)

Explanation: Upon finding an unresponsive, non-breathing adult in a safe scene, the immediate priority is to activate the emergency response system (call for help or send someone to do so). This ensures that professional medical assistance is on its way before initiating CPR.

Q12. Dressings and pads are used to:

- A) Help control bleeding and keep germs out
- B) Immobilize injuries
- C) Provide warmth
- D) All of the above

Correct Answer: A (Help control bleeding and keep germs out)

Explanation: Dressings and pads serve a dual purpose: to apply direct pressure to control bleeding and to provide a sterile barrier that prevents pathogens from entering the wound. While they might offer minimal warmth, their primary functions are hemostasis and infection control.

Q13. What method of manhandling technique would you use if a crew member is unconscious?

- A) Fireman's lift
- B) Simple forward and aft carry
- C) Both the above options
- D) Neither of the above options

Correct Answer: C (Both the above options)

Explanation: When a crew member is unconscious, safe and effective extrication requires specific techniques. Both the Fireman's lift and the simple forward and aft carry are recognized methods suitable for moving an incapacitated individual, depending on the situation and rescuer's capability.

Q14. The treatment for fever is:

- A) Aspirin
- B) Ibuprofen
- C) Acetaminophen
- D) All of the above

Correct Answer: C (Acetaminophen)



Q15. Hypovolemic shock results from:

- A) Decreased volume of blood
- B) An allergic reaction
- C) A heart attack
- D) A stroke

Correct Answer: A (Decreased volume of blood)

Explanation: Hypovolemic shock occurs when the body loses a significant volume of blood or fluids, leading to a decrease in circulating blood volume and inadequate tissue perfusion. Other causes of shock, such as anaphylactic or cardiogenic shock, stem from different underlying mechanisms.

Q16. What is the average heartbeat of an adult at rest?

- A) 50-60 beats per minute
- B) 60-100 beats per minute
- C) 100-120 beats per minute
- D) 120-140 beats per minute

Correct Answer: B (60-100 beats per minute)

Explanation: A normal resting heart rate for an adult typically falls within the range of 60 to 100 beats per minute. Rates outside this range may indicate underlying medical conditions or the need for further assessment.

Q17. Increased respiratory difficulty accompanied by a weak, ineffective cough, wheezing, high-pitched crowing noises, and cyanosis are signs of:

- A) Pneumonia
- B) Asthma
- C) Severe airway obstruction
- D) Heart failure

Correct Answer: C (Severe airway obstruction)

Explanation: The signs described-difficulty breathing, weak cough, wheezing, stridor, and cyanosis-are classic indicators of a severe airway obstruction, which impedes airflow to the lungs. While these symptoms can occur in other conditions, they most directly point to a compromised airway.

Q18. Which of the following should be done for a crew member experiencing a heat-related illness?

- A) Apply cool, wet cloths to the body
- B) Give the victim plenty of fluids to drink
- C) Move the victim to a cool, shaded area
- D) All of the above

Correct Answer: D (All of the above)

Explanation: Managing heat-related illnesses involves a multi-faceted approach including cooling the body with wet cloths, encouraging fluid intake if conscious and able to swallow, and moving the person to a cooler environment. These actions collectively help reduce body temperature and prevent further heat injury.



- A) Yes
- B) No
- C) Only in an emergency
- D) It doesn't matter

Correct Answer: B (No)

Explanation: A tourniquet should never be covered with clothing or bandages, as this can obscure the site, impede assessment of its effectiveness, and potentially delay vital medical attention. The tourniquet and the wound site should remain visible for medical personnel.

Q20. You have a casualty who is not breathing but has injuries to the face that prevent you from performing mouth-to-mouth resuscitation. What do you do?

- A) Perform chest compressions only at a rate of 100-120 per minute
- B) Attempt to open the victim's mouth using a jaw thrust maneuver
- C) Give rescue breaths through the nose
- D) Call for immediate medical assistance

Correct Answer: A (Perform chest compressions only at a rate of 100-120 per minute)

Explanation: When mouth-to-mouth resuscitation is impossible due to facial injuries, chest compressions are critical for maintaining circulation. Performing continuous chest compressions at a rate of 100-120 per minute is the recommended approach for CPR in such situations according to current guidelines.

Q21. What do you do after giving rescue breaths to an unconscious adult?

- A) Check for a pulse
- B) Continue rescue breaths
- C) Assess the victim's pupils
- D) Administer CPR

Correct Answer: A (Check for a pulse)

Explanation: Following the administration of rescue breaths in unresponsive, non-breathing individuals, the immediate next step is to check for a pulse. This assessment determines whether chest compressions are necessary as part of CPR.

Q22. Can you use an adult AED on an infant?

- A) Yes
- B) No
- C) Only if the infant is at least 6 months old
- D) Only if the infant is at least 1 year old

Correct Answer: B (No)

Explanation: Adult AEDs are not recommended for use on infants under eight years old or weighing less than 55 pounds. Paediatric pads and attenuator devices are designed to deliver a lower energy shock appropriate for infants and children, minimizing the risk of cardiac damage.

Q23. Difficulty breathing may be a sign of:

- A) Heart attack
- B) Asthma attack
- C) Anaphylaxis
- D) All of the above

Correct Answer: D (All of the above)



Q24. The type of shock that is caused by a severe infection is called:

- A) Cardiogenic shock
- B) Hypovolemic shock
- C) Septic shock
- D) Anaphylactic shock

Correct Answer: C (Septic shock)

Explanation: Septic shock is a life-threatening condition caused by a severe infection that triggers a widespread inflammatory response, leading to dangerously low blood pressure and organ dysfunction. While other types of shock involve circulatory issues, septic shock is specifically linked to infection.

Q25. After discovering a possible fracture, you should assess which of the following before and after splinting?

- A) Capillary refill, pulse, and sensation
- B) Blood pressure, heart rate, and respiratory rate
- C) Level of consciousness, pupil response, and motor function
- D) All of the above

Correct Answer: A (Capillary refill, pulse, and sensation)

Explanation: Before and after splinting any suspected fracture, it is crucial to assess the '5 Ps': Pain, Pallor, Pulses, Paresthesia (sensation), and Paralysis. Specifically, assessing distal circulation (pulse, capillary refill) and nerve function (sensation) helps ensure the splint is not too tight and does not cause further injury.

Q26. If you want to be a first aider at work, you should:

- A) Take a first aid course
- B) Get certified in CPR
- C) Arrange for your employer to put you on a first aiders course
- D) All of the above

Correct Answer: D (All of the above)

Explanation: Becoming a competent first aider requires formal training and certification. Taking a recognized first aid course, obtaining CPR certification, and ensuring employer support are all essential steps to be adequately prepared to provide care.

Q27. When is the best time to administer CPR?

- A) As soon as possible after a cardiac incident
- B) Only after the victim has been moved to a safe location
- C) When the victim's heart rate is below 60 beats per minute
- D) When the victim's breathing is shallow or labored

Correct Answer: A (As soon as possible after a cardiac incident)

Explanation: Cardiopulmonary Resuscitation (CPR) is most effective when initiated as soon as possible after a person's heart stops beating (cardiac arrest). Immediate chest compressions are vital to maintain blood flow to the brain and vital organs until advanced medical care can be provided.



- A) Apply direct pressure with a clean or sterile dressing
- B) Elevate the injured limb
- C) Control bleeding by applying a tourniquet
- D) Call for medical assistance

Correct Answer: A (Apply direct pressure with a clean or sterile dressing)

Explanation: The primary and immediate step in managing bleeding wounds is to apply direct pressure using a clean or sterile dressing to help control hemorrhage. Elevating the limb and applying a tourniquet are secondary measures, used if direct pressure is insufficient for severe bleeding.

Q29. In topographic anatomy, the term "lateral" means:

- A) Away from the midline of the body
- B) Nearer the head
- C) Away from the head

Correct Answer: A (Away from the midline of the body)

Explanation: In anatomical terminology, 'lateral' refers to a position that is away from the midline or center of the body. Conversely, 'medial' means closer to the midline.

Q30. The onset of shock in infants and children:

- A) May be sudden and severe
- B) Is no different than in adults
- C) Usually results from cardiac problems

Correct Answer: A (May be sudden and severe)

Explanation: Infants and children have different physiological responses to shock than adults; their condition can deteriorate rapidly, and they may present with less obvious early signs. They often compensate for longer but can decompensate suddenly, making prompt recognition and intervention critical.