



Medical First Aid (MFA)

196 Practice Questions and Answers for STCW Exams

Updated for DG Shipping India Exams | Source: brightmariner.com

Practice Set 1

Q1. Which one of the following should the first provider do when helping a crew with a snake bite?

- A) Do not remove jewelry or clothing in the area of the bite
- B) If a crew is bitten on the arm, elevate the arm above the level of the heart
- C) If the snake is still there, go back and approach from another direction to offer help
- D) Suck the venom from the snake bite

Correct Answer: B (If a crew is bitten on the arm, elevate the arm above the level of the heart)

Q2. What is the best position for a conscious casualty suffering from internal bleeding?

- A) Laying down with legs raised
- B) Lying flat down
- C) Half-sitting position
- D) Recovery position

Correct Answer: A (Laying down with legs raised)

Q3. The correct ratio of compression to ventilation for two rescuer CPR on a child is:

- A) 5:1
- B) 5:2
- C) 15:1
- D) 15:2

Correct Answer: D (15:2)

Q4. Shock is best defined as:

- A) Inadequate circulation to the body's tissues
- B) Inadequate delivery of oxygen to the lungs
- C) Inadequate elimination of waste products from the body
- D) Inadequate amount of insulin

Correct Answer: A (Inadequate circulation to the body's tissues)

Q5. When assessing the crew, you should feel for a pulse for:

brightmariner.com



- A) 3 seconds
- B) 5 to 10 seconds
- C) 15 to 20 seconds
- D) 20 to 30 seconds

Correct Answer: B (5 to 10 seconds)

Q6. For any victim, the correct compression rate is:

- A) At least 120 per minute
- B) At least 80 per minute
- C) At least 90 per minute
- D) 100 to 120 per minute

Correct Answer: D (100 to 120 per minute)

Q7. One sign that a bandage may be too tight after completing compress bandage is:

- A) Blue color of the skin above the bandage
- B) Swelling below the bandage and heavy pain
- C) Heartbeat increases and a dizzy feeling
- D) Swelling above the bandage

Correct Answer: A (Blue color of the skin above the bandage)

Q8. Sun rays and light reflected from a bright surface (e.g., sea) can cause damage to the skin and eyes. What is this type of burn called?

- A) Dry burns
- B) Radiation burns
- C) Cold burns
- D) Electrical burns

Correct Answer: B (Radiation burns)

Q9. The treatment of electrical burns is:

- A) Remove loose skin and apply ointment. Don't secure with a bandage
- B) Apply lotions and ointment to the injured area and secure with a bandage
- C) Place sterile dressing over the burn and secure with a bandage
- D) Break blisters and secure with a bandage

Correct Answer: C (Place sterile dressing over the burn and secure with a bandage)

Q10. Asthma attacks can be triggered by:

- A) Loud music
- B) Too much fresh air activity
- C) Not keeping to the diet
- D) Nervous tension, allergy, or non-obvious cause

Correct Answer: D (Nervous tension, allergy, or non-obvious cause)

Medical First Aid (MFA) Exit Exam Prep

Q11. If a person is in a state of shock, the correct thing to do is:

brightmariner.com



- A) Apply hot-water bottles to keep the patient warm
- B) Move the casualty as much as possible
- C) Be kind to the casualty and give anything to eat or drink at the first opportunity
- D) Treat and reassure the casualty and stay with the person at all times

Correct Answer: D (Treat and reassure the casualty and stay with the person at all times)

Q12. When you suspect a casualty has fractured a bone, you should:

- A) Raise the affected portion of the body above the level of the casualty's head
- B) Massage the affected area to prevent stiffness
- C) Rinse the area with cold water
- D) Immobilize the affected area

Correct Answer: D (Immobilize the affected area)

Q13. If the heart of a casualty has stopped, brain damage is likely to occur after:

- A) Beyond 6 minutes
- B) Beyond 20 minutes
- C) Beyond 1 minute
- D) Beyond 10 seconds

Correct Answer: A (Beyond 6 minutes)

Q14. The most effective method of artificial respiration is:

- A) Silvester method
- B) Mouth-to-mouth method
- C) Heath Robinson method
- D) Mouth-to-nose method

Correct Answer: B (Mouth-to-mouth method)

Q15. A resuscitator is:

- A) An insulated and heated bag that is used to wrap around a casualty suffering from hypothermia
- B) An oxygen tank, with a demand valve and mask
- C) An electrical device with two paddles that can be used to restart the heart
- D) A plastic tube that fits over the casualty's throat to keep an airway open

Correct Answer: B (An oxygen tank, with a demand valve and mask)

Q16. The best way to control severe bleeding is:

- A) Raise the bleeding part above the level of the head
- B) Direct pressure over the wound
- C) Application of a tourniquet
- D) Direct pressure on a pressure point

Correct Answer: B (Direct pressure over the wound)



- A) The canvas pole stretcher
- B) The Hart Imco stretcher
- C) The SOLAS stretcher
- D) The Neil Robertson stretcher

Correct Answer: D (The Neil Robertson stretcher)

Q18. The unconscious or recovery position is used to:

- A) Ease the pain of broken bones
- B) Minimize nose bleeding
- C) Prevent the casualty from drowning in his own vomit
- D) Correct for any spinal injury

Correct Answer: C (Prevent the casualty from drowning in his own vomit)

Q19. In the absence of a doctor, artificial respiration and heart compression should be continued until:

- A) When the casualty turns pale and starts to go cold CPR can be stopped
- B) Until the heart starts beating or the rescuer is unable to continue because of fatigue
- C) When the casualty shows no response to CPR after 20 minutes, it is useless to continue
- D) When the casualty has fixed and dilated pupils for 15 minutes you should stop CPR

Correct Answer: B (Until the heart starts beating or the rescuer is unable to continue because of fatigue)

Q20. The Recovery Position is:

- A) The patient is seated in a position with the head kept as low as possible
- B) The patient is placed in a 'face-to-the-floor' position with arms and legs arranged in order to stabilize this position
- C) The patient is placed flat on a bed
- D) The patient is seated in an upright position and with the arms and legs arranged in order to keep this position stable

Correct Answer: B (The patient is placed in a 'face-to-the-floor' position with arms and legs arranged in order to stabilize this position)

Q21. What do you call the method used for bone-soft part injuries?

- A) REHAB-method
- B) ABC-method
- C) First Aid-method
- D) ICE-method

Correct Answer: D (ICE-method)

Q22. You witness a man getting electric current through his body and is stuck to the dangerous area. It is impossible to switch off the current by any main switch. How do you break the current safely?

- A) Stand on dry insulating material and pull the person away with isolating material
- B) Just take the casualty in your arms and pull the person away
- C) Apply fish oil on your hands and cut the cords by the use of any metallic pliers
- D) Call the electrician immediately

Correct Answer: A (Stand on dry insulating material and pull the person away with isolating material)



Q23. Abdominal Thrust is a technique that involves applying a series of thrusts to the upper abdomen to air out of a choking casualty's lungs. How do you perform this technique?

- A) Lay the casualty on a hard surface, e.g., deck, and press firmly and rapidly on the middle of the lower half of the breastbone
- B) Remove the obstruction and restore normal breathing
- C) Let the casualty grab a list and hang upside down for a period of a minimum of 5 minutes
- D) Stand behind the casualty. Clench your fist with the thumb inwards in the center of the upper abdomen. Grasp your fist with your other hand and pull quickly inwards

Correct Answer: D (Stand behind the casualty. Clench your fist with the thumb inwards in the center of the upper abd

Q24. When acting as a watcher or lookout at a cargo hold and men below show signs of distress, what must you do?

- A) Try to rescue them yourself
- B) Raise the alarm immediately
- C) Lower additional breathing equipment
- D) Don a B.A. set and enter the space

Correct Answer: B (Raise the alarm immediately)

Q25. A severe blow to or a heavy fall on the upper part of the abdomen (solar plexus) can upset the regularity of breathing. What are the symptoms and signs?

- A) The casualty is speaking in a very loud manner
- B) The casualty may start sweating profusely and develop a fever
- C) The casualty feels very hungry
- D) Difficulty in breathing in, and the casualty may be unable to speak

Correct Answer: D (Difficulty in breathing in, and the casualty may be unable to speak)

Q26. In which way may intake of poisonous material occur?

- A) By inhalation
- B) Swallowing
- C) Skin penetration and skin absorption
- D) All mentioned

Correct Answer: D (All mentioned)

Q27. Due to exposure to heat fatigue, heat stroke, and dehydration, what is the maximum recommended effective temperature (ET) for a full workload in enclosed spaces?

- A) 30.5 degreesC
- B) 27.5 degreesC
- C) 29.0 degreesC
- D) 35.0 degreesC

Correct Answer: B (27.5 degreesC)

Q28. A casualty suddenly loses consciousness and falls to the ground letting out a strange cry. The patient is pale, red-blue in the face and froth may appear around the mouth. You are witnessing a major epileptic attack.



should you do?

- A) Move the casualty into a sit-up position and put something in the person's mouth to protect the tongue
 - B) Forcibly restrain and try to wake the casualty
 - C) Loosen tight clothing, ask all unnecessary bystanders to leave, and carefully place something soft under the head.
- If the casualty is unconscious, place the person in the Recovery position
- D) Give the casualty a lot to drink and keep talking to the person at all times

Correct Answer: C (Loosen tight clothing, ask all unnecessary bystanders to leave, and carefully place something soft under the head.)

Q29. Treatment of burns and scalds depends on the severity of the injury. What is the correct thing to do for minor burns and scalds?

- A) Break blisters, remove any loose skin or foreign objects from the injured area
- B) Place the injured part under slowly running cold water for at least 10 minutes, but preferably until the pain is gone. If no water is available, use any cold, harmless liquid
- C) Remove all sticky clothing from the casualty
- D) Apply lotions, ointments, or fat to the injury

Correct Answer: B (Place the injured part under slowly running cold water for at least 10 minutes, but preferably until the pain is gone.)

Q30. When acting as a watcher or lookout at a cargo hold and men below show signs of distress, what must you do?

- A) Raise the alarm immediately
- B) Try to rescue them yourself
- C) Lower additional breathing equipment
- D) Don a B.A. set and enter the space

Correct Answer: A (Raise the alarm immediately)

Practice Set 2

Q1. The acronym SAMPLE stand for signs and symptoms, allergies, medications, past history, _____ and events.

- A) Location of pain.
- B) Last bowel movement.
- C) Last meal.
- D) Latest related injury.

Correct Answer: C (Last meal.)

Q2. Which one of the following is not a system of the body:

- A) Respiratory.
- B) Musculoskeletal.
- C) Endocrine.
- D) Follicular.

Correct Answer: D (Follicular.)

Q3. Increased respiratory difficulty accompanied by a weak ineffective cough, wheezing, high-pitched crackles and cyanosis are signs of:



- A) Mild airway obstruction
- B) C.O.P.D.
- C) Severe airway obstruction.
- D) Complete airway obstruction.

Correct Answer: C (Severe airway obstruction.)

Q4. The most important step in managing shock is to:

- A) Keep the patient warm.
- B) Give CPR as soon as possible.
- C) Give first aid for the illness or injury.
- D) Elevate the lower extremities.

Correct Answer: C (Give first aid for the illness or injury.)

Q5. A small percentage of casualties with chronic obstructive pulmonary disease have hypoxic drive. These patients breathe because of a:

- A) High oxygen level.
- B) Low oxygen level.
- C) High carbon dioxide level.
- D) Low carbon dioxide level.

Correct Answer: B (Low oxygen level.)

Q6. The emergency responder is one link in the chain of services known as the:

- A) Emergency Patient Care (EPC) system.
- B) Emergency Medical Services (EMS) system.
- C) Professional Emergency Care (PEC) system.
- D) Community Medical Care (CMC) system.

Correct Answer: B (Emergency Medical Services (EMS) system.)

Q7. The acronym used to assist the emergency responder assessing the patient's level of responsiveness is:

- A) SAMPLE.
- B) EMCA.
- C) OPQRST.
- D) AVPU.

Correct Answer: D (AVPU.)

Q8. An industrial worker sustains a severe laceration to his forearm. Direct pressure to the wound fails to control the bleeding. The correct arterial pressure point to control the bleeding is the:

- A) Carotid.
- B) Radial.
- C) Femoral.
- D) Brachial.

Correct Answer: D (Brachial.)



- A) Normal blood circulation to that part.
- B) The presence of fractured fingers.
- C) Pain in the area.
- D) A lack of oxygen in the blood.

Correct Answer: A (Normal blood circulation to that part.)

Q10. Your primary survey of a casualty involved in a serious car collision shows only that he is confused. Later you find his pulse rate at 140 and weak, his skin cold and clammy and his breathing irregular and gasping. These signs, along with the mechanism of injury, indicate:

- A) An oncoming faint.
- B) Emotional stress.
- C) Internal bleeding.
- D) Onset of diabetic coma.

Correct Answer: C (Internal bleeding.)

Q11. For which one of the following infectious diseases is a vaccine presently available:

- A) Herpes.
- B) AIDS.
- C) Mononucleosis.
- D) Hepatitis B.

Correct Answer: D (Hepatitis B.)

Q12. Based on current research, which of the following statements about the AIDS virus is correct?

- A) It can be found in blood and semen.
- B) It can be transmitted by sharing eating utensils.
- C) It can be transmitted by shaking hands.
- D) It can be found in perspiration.

Correct Answer: A (It can be found in blood and semen.)

Q13. In a hazmat situation airway management and immobilization are carried out in the:

- A) hot zone
- B) warm zone
- C) cold zone
- D) neutral zone

Correct Answer: B (warm zone)

Q14. During the primary assessment of a responsive adult patient, you detect a breathing rate of 28 breaths per minute. You would categorize this as:

- A) Above normal.
- B) Below normal.
- C) Normal.
- D) Indeterminate.

Correct Answer: A (Above normal.)



- A) Remove the dressing and check the wound.
- B) Apply pressure to the femoral artery.
- C) Place a clean dressing on top and apply more pressure.
- D) Apply a tourniquet.

Correct Answer: C (Place a clean dressing on top and apply more pressure.)

Q16. The type of shock that is caused by a severe infection is called:

- A) Septic.
- B) Psychogenic.
- C) Cardiogenic.
- D) Hemorrhagic.

Correct Answer: A (Septic.)

Q17. Which one of the following is considered a breach of duty:

- A) failure to obtain consent.
- B) failure to wear your name tag.
- C) inappropriate use of lights and siren.
- D) insubordination.

Correct Answer: A (failure to obtain consent.)

Q18. Immediately before a seizure, the patient experiences an unpleasant odour. This phase is referred to as:

- A) Clonic.
- B) Aura.
- C) Tonic.
- D) Postical.

Correct Answer: B (Aura.)

Q19. The emergency responder should assume a head/spinal injury in any unwitnessed situation where the patient is:

- A) Alert.
- B) Responsive to pain.
- C) Not breathing.
- D) Unresponsive.

Correct Answer: D (Unresponsive.)

Q20. The secondary assessment of the patient consists of a head to toe examination and a check of the:

- A) Pressure points.
- B) Procedures for administering CPR.
- C) Manual stabilization of the head.
- D) Vital signs.

Correct Answer: D (Vital signs.)

Q21. You are taking blood pressure by palpation. A radial pulse indicates a blood pressure of at least:

Medical First Aid (MFA) Exit Exam Prep

Brightmariner.com



- A) 110 mmHg
- B) 100 mmHg
- C) 90 mmHg
- D) 80 mmHg

Correct Answer: D (80 mmHg)

Q22. You are called for an asthma attack, treatments may include all except:

- A) Metered dose inhaler
- B) Bronchodilators
- C) Nitroglycerin
- D) Inhaled steroids

Correct Answer: C (Nitroglycerin)

Q23. In topographic anatomy, the term "lateral" means:

- A) Nearer the midline of the body.
- B) Away from the midline of the body.
- C) Nearer the head.
- D) Away from the head.

Correct Answer: B (Away from the midline of the body.)

Q24. Which one of the following conditions may mimic the signs of acute alcohol intoxication?

- A) Hypoglycemia
- B) Congestive heart failure.
- C) Absence seizures.
- D) Anaphylactic shock.

Correct Answer: A (Hypoglycemia)

Q25. Which one of the following breathing diseases is included in C.O.P.D.:

- A) Croup.
- B) Hyperventilation
- C) Emphysema.
- D) Dyspnea

Correct Answer: C (Emphysema.)

Q26. A condition where air builds up in the pleural space, collapses the lung and puts pressure on the heart is called:

- A) Closed pneumothorax
- B) Tension pneumothorax
- C) Hemothorax
- D) Open pneumothorax

Correct Answer: B (Tension pneumothorax)



- A) 15 minutes.
- B) 30 minutes.
- C) 60 minutes.
- D) 90 minutes.

Correct Answer: B (30 minutes.)

Q28. You find a male patient with obvious difficulty breathing. He is using his neck muscles, and is cyanotic. There are red blotches on his chest, and his neck is swelling. You suspect :

- A) Anaphylaxis.
- B) Bronchitis.
- C) Emphysema.
- D) Asthma.

Correct Answer: A (Anaphylaxis.)

Q29. When you are giving mouth-to-nose AR, you should:

- A) Hold the casualty's mouth closed.
- B) Pinch the nostrils closed before blowing air into the casualty.
- C) Tilt the head back less than for the mouth-to-mouth method of AR.
- D) Keep the mouth and nose closed between breaths.

Correct Answer: A (Hold the casualty's mouth closed.)

Q30. During one-rescuer CPR for adults, the ratio of compressions to ventilations should be:

- A) 5:1
- B) 15:2
- C) 30:2
- D) 35:2

Correct Answer: C (30:2)

Practice Set 3

Q1. A guideline for normal systolic blood pressure in an adult male would be:

- A) 50 plus the man's age up to 150 mm. Hg.
- B) 65 plus the man's age up to 120 mm. Hg.
- C) 80 plus the man's age up to 130 mm. Hg.
- D) 100 plus the man's age up to 150 mm. Hg.

Correct Answer: D (100 plus the man's age up to 150 mm. Hg.)

Q2. A rapid body survey should take no longer than:



- A) 30 seconds
- B) 45 seconds
- C) 60 seconds
- D) 90 seconds

Correct Answer: A (30 seconds)

Q3. To correctly size an oropharyngeal airway place the flange at the corner of the mouth with the tip reaching:

- A) The angle of the patient's jaw
- B) The top of the patient's ear
- C) The patient's earlobe
- D) Two fingerswidth from the flange.

Correct Answer: C (The patient's earlobe)

Q4. ASA should be offered to the patient with chest pain who is taking nitroglycerin:

- A) Before the first dose of nitro
- B) After the first dose of nitro
- C) After the second dose of nitro
- D) After the third dose of nitro

Correct Answer: B (After the first dose of nitro)

Q5. When suctioning a non-breathing adult casualty, a first aider should:

- A) Apply suction as soon as the suction tip touches the mouth.
- B) Insert the suction tip deep into the larynx.
- C) Suction for no more than 15 seconds.
- D) Insert the tip with the curved side towards the tongue.

Correct Answer: C (Suction for no more than 15 seconds.)

Q6. Which of the following devices provides the highest percentage of O2 delivery?

- A) Nasal cannula.
- B) Plastic face mask.
- C) Partial rebreathing mask.
- D) Non-rebreathing mask.

Correct Answer: D (Non-rebreathing mask.)

Q7. You suspect a patient's pain is due to a myocardial infarction rather than angina because the pain:

- A) Is under the sternum.
- B) Radiates to the neck, jaw and arms.
- C) Is not relieved by rest or medication.
- D) Lasts for about 3 minutes.

Correct Answer: C (Is not relieved by rest or medication.)



- A) Three ribs are broken on each side of the chest.
- B) Part of the spine becomes separated from the ribs.
- C) The breastbone is broken in three places.
- D) Several ribs are broken in more than one place.

Correct Answer: D (Several ribs are broken in more than one place.)

Q9. Status epilepticus refers to:

- A) Normal seizure pattern in epilepsy.
- B) Seizures caused by a high fever.
- C) Continuous seizure activity.
- D) Seizures lasting longer than 2 minutes.

Correct Answer: C (Continuous seizure activity.)

Q10. OPQRST is used to assess:

- A) Pain
- B) Level of consciousness
- C) Vision
- D) Hearing

Correct Answer: A (Pain)

Q11. The recommended oxygen flow rate for nasal cannulae (nasal prongs) is:

- A) 1-6 litres/min.
- B) 6-8 litres/min.
- C) 8-12 litres/min.
- D) 10-15 litres/min.

Correct Answer: A (1-6 litres/min.)

Q12. When aligning the head in the eyes-front neutral position, watch for:

- A) Pulse, movement, sensation.
- B) Crepitus, pain, resistance.
- C) Pain, circulation, retention.
- D) Tingling, numbness, paralysis.

Correct Answer: B (Crepitus, pain, resistance.)

Q13. Manual support of the head and neck of the patient with suspected spinal injury can be released:

- A) Once the breathing has been checked.
- B) When a hard collar has been applied.
- C) If the patient is unconscious.
- D) When the patient is fully immobilized to a backboard.

Correct Answer: D (When the patient is fully immobilized to a backboard.)

Q14. To prepare a completely amputated part of the body for transportation to hospital with the casualty, you should:



- A) Place it in a clean plastic bag filled with cold water.
- B) Wrap it in clean, moist dressings and keep it cool.
- C) Wash it off and place it into a bag of crushed ice.
- D) Wrap it in clean, moist dressings and keep it at body temperature.

Correct Answer: B (Wrap it in clean, moist dressings and keep it cool.)

Q15. Hypovolemic shock results from:

- A) The body's reaction to a foreign protein.
- B) An overdose of insulin.
- C) Toxins produced by a severe infection.
- D) Decreased volume of the blood.

Correct Answer: D (Decreased volume of the blood.)

Q16. A conscious casualty who has suffered a stroke becomes unconscious. To keep his airway open and help him breathe, you should:

- A) Turn him into the recovery position with his unaffected side down.
- B) Place him in a semisitting position.
- C) Turn him into the recovery position with his unaffected side up.
- D) Place him on his back.

Correct Answer: A (Turn him into the recovery position with his unaffected side down.)

Q17. A worker has fallen from a six-foot ladder onto a concrete floor. Your first action should be.

- A) Take a SAMPLE history.
- B) Perform a secondary survey.
- C) Take charge and perform a scene survey.
- D) Assess the rate and quality of breathing.

Correct Answer: C (Take charge and perform a scene survey.)

Q18. To determine whether a c-spine injury has occurred, you should think about the Mechanism of Injury. This can best be described as:

- A) The force that causes the injury and the way it acts on the body.
- B) The circumstances after the injury.
- C) The weather conditions at the time of the incident.
- D) The area of the body that is injured

Correct Answer: A (The force that causes the injury and the way it acts on the body.)

Q19. The correct ratio of compressions to ventilations for two-rescuer CPR on a child is:

- A) 5:1
- B) 15:2
- C) 5:2
- D) 15:1

Correct Answer: B (15:2)



- A) Inadequate circulation to the body's tissues
- B) Inadequate delivery of oxygen to the lungs
- C) Inadequate elimination of waste products from the body.
- D) Inadequate amount of insulin.

Correct Answer: A (Inadequate circulation to the body's tissues)

Q21. The most critical factor in defibrillation is:

- A) The time from collapse to defibrillation.
- B) The skill of the AED responder.
- C) The patient's previous cardiac history.
- D) The type of defibrillator used.

Correct Answer: A (The time from collapse to defibrillation.)

Q22. The AED will shock a patient:

- A) With a sinus rhythm.
- B) In asystole.
- C) In ventricular fibrillation.
- D) With pulseless electrical activity (PEA).

Correct Answer: C (In ventricular fibrillation.)

Q23. In the hypothermic patient you should limit the number of shocks to a maximum of:

- A) 1.
- B) 2.
- C) 3.
- D) 4.

Correct Answer: A (1.)

Q24. When transporting a patient, a shock should be administered:

- A) Only in a vehicle that is properly grounded.
- B) Whenever the machine indicates shock necessary.
- C) When the vehicle is stopped.
- D) When the vehicle is travelling at less than 20 km per hour.

Correct Answer: C (When the vehicle is stopped.)

Q25. When using an AED on a patient who is wearing a pacemaker, place the electrode pads:

- A) Directly on pacemaker.
- B) Two inches away from pacemaker.
- C) Within one inch of the pacemaker.
- D) At least one inch away from pacemaker.

Correct Answer: D (At least one inch away from pacemaker.)

Q26. You have shocked a casualty and there are signs of circulation. You should:

brightmariner.com



- A) Remove the AED from the scene.
- B) Leave the AED attached to the casualty.
- C) Disconnect the cables from the AED.
- D) Remove the pads from the chest. 15 Burns

Correct Answer: B (Leave the AED attached to the casualty.)

Q27. Burns that involve all layers of the skin are:

- A) Superficial.
- B) Partial thickness.
- C) Full thickness.
- D) First degree.

Correct Answer: C (Full thickness.)

Q28. In determining the amount of body surface burned, the area of the palm of the hand represents ____ % of the body surface.

- A) One
- B) Two
- C) Five
- D) Nine

Correct Answer: A (One)

Q29. The most immediate serious complication associated with burns is:

- A) Shock
- B) Infection
- C) Scarring
- D) Hypothermia

Correct Answer: A (Shock)

Q30. A man is filling a gas tank on a generator when it bursts into flames. The casualty has a hoarse voice, he is blistered around the mouth and nose and both arms are reddened. You should classify this burn as:

- A) Critical
- B) Superficial
- C) Moderate
- D) Minor 16 Wound Management

Correct Answer: A (Critical)

Practice Set 4



- A) deep break in the skin involving significant bleeding.
- B) Partial or complete loss of a body part.
- C) The result of a sharp object driven through soft tissue.
- D) A scrape or rubbing away of the epidermis.

Correct Answer: D (A scrape or rubbing away of the epidermis.)

Q2. A soft tissue injury resulting from the impact of a blunt object is called:

- A) laceration.
- B) An avulsion.
- C) A contusion.
- D) A concussion.

Correct Answer: C (A contusion.)

Q3. The acronym SHARP refers to:

- A) The assessment of the severity of a wound injury.
- B) Signs and symptoms used to determine the need to transport.
- C) The depth of a penetrating wound and potential damage.
- D) Signs of infection as assessed in wound injuries.

Correct Answer: D (Signs of infection as assessed in wound injuries.)

Q4. Responders involved in wound management should consider:

- A) Wounds should be thoroughly flushed to prevent infection.
- B) All open wounds are contaminated to some degree.
- C) The risk of infection continues until the wound is dressed and bandaged.
- D) Gloves should be worn if you suspect the patient may be ill.

Correct Answer: B (All open wounds are contaminated to some degree.)

Q5. Tetanus is a condition:

- A) That involves specific infection in the lower jaw.
- B) That only occurs in the third world.
- C) That is caused by a virus
- D) That can be prevented by immunization.

Correct Answer: D (That can be prevented by immunization.)

Q6. Which one of the following factors increases the risk of life -threatening injury:

- A) High velocity.
- B) Low velocity.
- C) High density.
- D) Low energy.

Correct Answer: A (High velocity.)



Q7. Which action is part of first aid for a nosebleed?

- A) Place an ice-pack on the back of the neck.
- B) Plug the nose with gauze.
- C) Lean forward and firmly pinch the soft parts of the nose.
- D) Lean the casualty backwards in a sitting position.

Correct Answer: C (Lean forward and firmly pinch the soft parts of the nose.)

Q8. To prevent further contamination and infection of a wound, you should cleanse the surrounding skin by:

- A) Swabbing in circular motions around the wound.
- B) Wiping lightly over the edges of the wound.
- C) Swabbing from one side of the wound to the other.
- D) Wiping away from the edges of the wound.

Correct Answer: D (Wiping away from the edges of the wound. 17 Heat and Cold Illness and Injury)

Q9. The body loses heat in a variety of ways including:

- A) radiation, conduction, evaporation and respiration.
- B) radiation, refrigeration, evaporation and conduction.
- C) conduction, convection, evaporation and refrigeration.
- D) submersion, convection, radiation and respiration.

Correct Answer: A (radiation, conduction, evaporation and respiration.)

Q10. If it becomes necessary to thaw a hand with deep frostbite you should ensure that:

- A) The patient sees a doctor no later than 48 hrs after thawing
- B) There is no danger of refreezing
- C) Adequate pain relief medication is available
- D) Capillary refill takes no longer than 15 seconds

Correct Answer: B (There is no danger of refreezing)

Q11. You have been called for an 81 year old unconscious male patient. He is in his living room. Despite the hot day, he is overly dressed. He is flushed and dry, extremely hot to touch. You suspect:

- A) Heat exhaustion.
- B) Stroke.
- C) Hyperglycemia.
- D) Heatstroke. Lifting and Carrying

Correct Answer: D (Heatstroke. Lifting and Carrying)

Q12. Body mechanics refers to:

- A) Positioning and movement of the body
- B) Number of rescuers needed to move a patient
- C) A type of rescue carry
- D) A special device to lift heavy loads

Correct Answer: A (Positioning and movement of the body)



You should

- A) Grab his wrists and drag him lengthwise.
- B) Keep his body rigid, support his head and neck and roll him away from the scene.
- C) Tie his legs together and drag him feet first.
- D) Grasp his clothing under his shoulders, support his head and neck, and drag him lengthwise.

Correct Answer: D (Grasp his clothing under his shoulders, support his head and neck, and drag him lengthwise.)

Q14. To safely perform a chair carry you need

- A) 2 rescuers
 - B) 3 rescuers
 - C) 4 rescuers
 - D) 5 rescuers
- 19 Musculoskeletal - Upper Limbs

Correct Answer: B (3 rescuers)

Q15. Crepitus is:

- A) . Poor distal circulation
- B) A grating sound caused by bone ends rubbing together
- C) A type of splint
- D) A type of fracture

Correct Answer: B (A grating sound caused by bone ends rubbing together)

Q16. After discovering a possible fracture you should assess which of the following before and after splinting:

- A) Sensation, morbidity, protrusion.
- B) Capillary refill, pulse, sensation.
- C) Pulse, motor function, sensation.
- D) Crepitus, resistance, pain.

Correct Answer: C (Pulse, motor function, sensation.)

Q17. A worker has had his hand caught in a car door. You suspect several fractured bones in the hand. The hand should be splinted:

- A) Flat against the splint.
- B) In a tight fist position.
- C) With the fingers taped together.
- D) In the position of function.

Correct Answer: D (In the position of function.)

Q18. In the case of suspected fracture of the clavicle, emergency responders should use:

- A) An arm sling supported with broad bandages.
- B) A St. John tubular sling tied on the injured side.
- C) A St. John tubular sling tied on the uninjured side.
- D) Rigid splints to support the arm and support the shoulder.

Correct Answer: C (A St. John tubular sling tied on the uninjured side.)



- A) Sprain
- B) Fracture
- C) Dislocation
- D) Strain

Correct Answer: C (Dislocation)

Q20. After an extremity fracture has been immobilized, the responder should check for circulation:

- A) In the injured limb only.
- B) Distal to the injury.
- C) Proximal to the injury.
- D) At the site of the fracture.

Correct Answer: B (Distal to the injury.)

Q21. The longest, strongest bone in the body is the:

- A) Humerus.
- B) Fibula.
- C) Tibia.
- D) Femur.

Correct Answer: D (Femur.)

Q22. Effective immobilization of the tibia includes immobilization of the:

- A) Knee, tibia, fibula and ankle.
- B) Femur, knee, tibia and ankle.
- C) Femur and ankle.
- D) Hip, femur, knee, tibia and ankle.

Correct Answer: A (Knee, tibia, fibula and ankle.)

Q23. You can best control the swelling and pain of an ankle sprain by:

- A) Tight bandages and a heating pad.
- B) Rest, immobilization, application of cold, and elevation.
- C) Rigid splints and bandages
- D) Compression, elevation and application of heat.

Correct Answer: B (Rest, immobilization, application of cold, and elevation.)

Q24. If the patient has an open fracture responders should :

- A) Attempt to push bones back into the wound.
- B) Use bulky dressings to pad around the protruding bones ends.
- C) Apply pressure directly over the fracture to control bleeding.
- D) Apply a tourniquet above the fracture site.

Correct Answer: B (Use bulky dressings to pad around the protruding bones ends.)



- A) fractured pelvis.
- B) An injured knee joint.
- C) A dislocated hip.
- D) A mid-shaft femur fracture.

Correct Answer: D (A mid-shaft femur fracture.)

Q26. A concussion is best described as:

- A) Bruising or swelling of the spinal cord.
- B) Tearing of brain tissue.
- C) Pooling of blood in the brain.
- D) Temporary loss of brain function

Correct Answer: D (Temporary loss of brain function)

Q27. You are examining the head of an infant who has been involved in a car crash. You need to be aware of:

- A) Soft spots in the infant's skull.
- B) Infant's pupils react differently than adults.
- C) Whether or not the infant can cry forcefully.
- D) The startle reflex.

Correct Answer: A (Soft spots in the infant's skull.)

Q28. When immobilizing a patient on a spine board, which part of the body is the first to be strapped?

- A) Chest
- B) Pelvis
- C) Legs
- D) Head

Correct Answer: A (Chest)

Q29. Contusions are:

- A) Usually controlled with direct pressure
- B) Very serious for the patient due to increased pressure in the brain.
- C) Usually associated with scalp lacerations
- D) Almost always seen in children.

Correct Answer: B (Very serious for the patient due to increased pressure in the brain.)

Q30. The Glasgow Coma Scale measures three basic functions. They are:

- A) Eye, verbal and motor responses.
- B) Pulse rate, speech, involuntary movement.
- C) Pulse rate, respiration rate, eye response.
- D) Respiration rate, eye response, voluntary movement.

Correct Answer: A (Eye, verbal and motor responses.)



Q1. Which one of the following changes in vital signs is characteristic of brain injury:

- A) Increase in pulse rate.
- B) Constant respiratory rate.
- C) increase in blood pressure.
- D) Decrease in blood pressure.

Correct Answer: C (increase in blood pressure.)

Q2. "Raccoon eyes" indicates:

- A) Possible eye injury
- B) Possible fracture of the jaw
- C) Possible scalp laceration
- D) Possible head injury

Correct Answer: D (Possible head injury)

Q3. You are called to the scene of a motorcycle crash. The rider is dazed and walking around. You should:

- A) Suspect spinal injury and manage accordingly.
- B) Not worry about spinal injury because the person is mobile.
- C) Advise the person to lie down in case of fainting.
- D) Determine the chief complaint and provide first aid for it.

Correct Answer: A (Suspect spinal injury and manage accordingly.)

Q4. When managing a possible spinal injury:

- A) The cervical immobilization device is applied by the police.
- B) Transport the patient in the position of greatest comfort.
- C) Apply a cervical immobilization device before assessing the patient.
- D) Responders must provide initial stabilization by supporting the head and ensuring neutral alignment

Correct Answer: D (Responders must provide initial stabilization by supporting the head and ensuring neutral alignment)

Q5. A helmet must be removed from a patient:

- A) If there are no airway or breathing problems.
- B) If the patient will be immobilized to a long spinal immobilization device.
- C) When the helmet has a face mask that interferes with the responder's ability to assist with ventilations
- D) It is a full- face helmet.

Correct Answer: C (When the helmet has a face mask that interferes with the responder's ability to assist with ventilations)

Q6. When dealing with pelvic injuries you must always consider the possibility of:

- A) Ruptured bladder.
- B) Ruptured spleen.
- C) Spinal injuries.
- D) Rib fractures.

Correct Answer: C (Spinal injuries.)



- A) Measure the distance from the ear lobe to the shoulder.
- B) Measure the distance from the corner of the mouth to the ear lobe.
- C) Measure the distance from the cheekbone to the shoulder blade.
- D) Measure the distance from the trapezius muscle to the angle of the jaw

22 Poisons, Bites and Stings

Correct Answer: D (Measure the distance from the trapezius muscle to the angle of the jaw)

Q8. Contact dermatitis occurs when:

- A) The skin comes in contact with a poisonous substance.
- B) A poison enters the eye of a contact lens wearer.
- C) Hot gases are inhaled.
- D) A poison is injected under the skin.

Correct Answer: A (The skin comes in contact with a poisonous substance.)

Q9. Carbon monoxide:

- A) May result from fires or automobile exhaust.
- B) Has a very distinct odour
- C) Is not life-threatening.
- D) Requires the administration of low concentration O2.

Correct Answer: A (May result from fires or automobile exhaust.)

Q10. A child has swallowed an unknown poisonous substance. You should:

- A) Dilute the poison with several glasses of cool water.
- B) Call your local Poison Control Centre and follow directions.
- C) Give a solution of mild liquid dish detergent and water.
- D) Use activated charcoal.

14 23 Anatomy and Physiology

Correct Answer: B (Call your local Poison Control Centre and follow directions.)

Q11. The external layer of the skin is called::

- A) Cutaneous tissue.
- B) Dermis.
- C) Adipose tissue.
- D) Epidermis.

Correct Answer: D (Epidermis.)

Q12. Muscles are attached to the bones by:

- A) Ligaments.
- B) Meninges.
- C) Cartilage.
- D) Tendons.

Correct Answer: D (Tendons.)

Q13. The appendix is located in which quadrant of the abdomen:



- A) Lower right.
- B) Upper right.
- C) Upper left.
- D) Lower left.

Correct Answer: A (Lower right.)

Q14. The white exterior portion of the eye is called the:

- A) Pupil.
- B) Sclera.
- C) Cornea.
- D) Iris.

Correct Answer: B (Sclera.)

Q15. A skeletal injury to the lower back could result from direct trauma to the:

- A) Cervical vertebrae.
- B) Thoracic vertebrae.
- C) Lumbar vertebrae.
- D) Coccygeal vertebrae.

Correct Answer: C (Lumbar vertebrae.)

Q16. The pulse point located in the upper portion of the thigh is called the:

- A) Popliteal.
- B) Femoral.
- C) Brachial.
- D) Temporal.

Correct Answer: B (Femoral.)

Q17. Which one of the following is a function of the skin?

- A) Regulation of oxygen levels.
- B) Production of proteins.
- C) Regulation of body temperature.
- D) Removal of carbon dioxide.

Correct Answer: C (Regulation of body temperature.)

Q18. Urine is expelled from the body through the:

- A) Kidneys.
- B) Bladder.
- C) Ureters.
- D) Urethra.

Correct Answer: D (Urethra.)



- A) Spleen.
- B) Liver.
- C) Pancreas.
- D) Gall bladder.

Correct Answer: C (Pancreas.)

Q20. When communicating with patients:

- A) Stay at eye level & maintain eye contact.
- B) Stay as close to the patient as possible.
- C) Use medical terminology to enhance credibility.
- D) Hide facts if the situation is serious.

Correct Answer: A (Stay at eye level & maintain eye contact.)

Q21. In cases of behavioural disorders, the responder should:

- A) Always use restraints.
- B) Give the patient a choice in the treatment
- C) Play along with auditory or visual hallucinations.
- D) Force the patient to make decisions.

Correct Answer: B (Give the patient a choice in the treatment)

Q22. The best way to prevent a Critical Incident Stress situation from becoming worse is:

- A) Ignore it.
- B) Be familiar with the signs and symptoms.
- C) Ensure early use of proper medication.
- D) Force the responder to talk about the situation.

Correct Answer: B (Be familiar with the signs and symptoms.)

Q23. Remove a contact lense only if:

- A) There is a chemical burn to the eye.
- B) The eyeball is injured.
- C) Transport time is short.
- D) The casualty is wearing hard lenses.

Correct Answer: A (There is a chemical burn to the eye.)

Q24. You discover a particle embedded in the casualty's cornea. You should:

- A) Remove the particle with a moist corner of a facial tissue or clean cloth.
- B) Flush the eye with water for 10 minutes.
- C) Use a splinter forcep to remove the particle.
- D) Cover the eye and transport the casualty to medical help.

Correct Answer: D (Cover the eye and transport the casualty to medical help.)



- A) Bleeding from the conjunctiva.
- B) Gritty feeling in the eyes.
- C) Lacerations to the globe.
- D) Discolouration of the orbits.

Correct Answer: B (Gritty feeling in the eyes.)

Q26. Eye injuries that include lacerations:

- A) Should be covered with clean, moist dressings.
- B) Indicate possible skull fracture.
- C) Usually involve a great deal of bleeding.
- D) Require moderate pressure on the eye to control bleeding.

Correct Answer: C (Usually involve a great deal of bleeding.)

Q27. Irrigate eyes with:

- A) Diluted vinegar.
- B) Saline solution.
- C) An appropriate chemical antidote.
- D) Sodium bicarbonate.

Correct Answer: B (Saline solution.)

Q28. Signs and symptoms of acute abdomen include:

- A) Excessive hunger or thirst.
- B) Hypertension and brachycardia.
- C) Distension with rebound tenderness.
- D) Fever with deep, shallow breathing.

Correct Answer: C (Distension with rebound tenderness.)

Q29. Signs that a hernia is serious include:

- A) Pain and tenderness at site.
- B) Pale, ashen skin colour at site.
- C) Hernia can be pushed back into body.
- D) Hernia is above the diaphragm.

Correct Answer: A (Pain and tenderness at site.)

Q30. Which one of the following is not crucial in cases of abdominal distress:

- A) Establish the cause of the pain.
- B) Give oxygen by non-rebreather mask.
- C) Be prepared to deal with vomiting.
- D) Monitor the patient for signs of hypovolemic shock.

Correct Answer: A (Establish the cause of the pain.)



Q1. In the case of abdominal evisceration, the emergency responder should:

- A) Irrigate the wound before dressing it.
- B) Give oxygen by nasal canula.
- C) Dress the wound with a moist, sterile dressing
- D) Gently reposition the organs in the abdominal cavity.

Correct Answer: C (Dress the wound with a moist, sterile dressing)

Q2. "Referred pain":

- A) Occurs when the sensation of pain is delayed.
- B) Results when the two nervous systems come into contact.
- C) Is the symptomatic description of pain by the patient.
- D) Is the description of pain as recorded on the patient care report.

Correct Answer: B (Results when the two nervous systems come into contact.)

Q3. Trauma to male and female genitalia:

- A) Often involves significant bleeding.
- B) Is not usually painful because of protected nerve endings.
- C) Usually results in sterility.
- D) Often results in post-traumatic infection.

Correct Answer: A (Often involves significant bleeding.)

Q4. The priorities of MCIs include all but one of the following.

- A) Overestimating the resources that may be required.
- B) Comprehensive care of all patients.
- C) Planning for the positioning of all vehicles.
- D) Arranging for advanced level care providers at the scene.

Correct Answer: B (Comprehensive care of all patients.)

Q5. At the scene of a MCI:

- A) All patients are assessed quickly and triaged.
- B) Only conscious patients are assessed.
- C) Patients without a detectable pulse are given the highest priority.
- D) All patients are transported immediately.

Correct Answer: A (All patients are assessed quickly and triaged.)

Q6. Typically in a triage system:

- A) Those whose survival requires immediate care are classified level 3.
- B) Those with minor injuries but suffering extreme pain are classified level 1.
- C) Those in cardiac arrest or not breathing are classified level 2.
- D) Those who will survive if care is somewhat delayed are classified level 2.

Correct Answer: D (Those who will survive if care is somewhat delayed are classified level 2.)



Q7. The most knowledgeable responder in the first ambulance is generally assigned to role of:

- A) Incident manager.
- B) Triage officer.
- C) Staging officer.
- D) Communications officer.

Correct Answer: A (Incident manager.)

Q8. Of the following casualties, which would you care for first? The casualty who is:

- A) In shock without apparent injuries, but conscious.
- B) Unconscious and lying on his back.
- C) Bleeding from the forehead, but conscious.
- D) Unconscious and lying on his stomach.

Correct Answer: B (Unconscious and lying on his back.)

Q9. The factor which indicates a drug related emergency is life-threatening is:

- A) Altered mental status.
- B) Dilated pupils slow to respond to light.
- C) Lack of coordination and slurred speech.
- D) High, low or irregular pulse.

Correct Answer: D (High, low or irregular pulse.)

Q10. Constant monitoring of the patient in a drug- related emergency is important because:

- A) The patient will likely try to take more of the drug.
- B) These patients are more likely to sue.
- C) The patient's condition may change quickly.
- D) The patient is more likely to become violent. 29 Communications

Correct Answer: C (The patient's condition may change quickly.)

Q11. The most accurate source of time for reporting purposes is:

- A) The responder's watch.
- B) The dispatch unit.
- C) The local radio station.
- D) The patient's watch.

Correct Answer: B (The dispatch unit.)

Q12. An essential element of a responder's verbal report includes:

- A) The name of the patient.
- B) The patient's telephone number.
- C) The time of the incident.
- D) The patient's response to the care given

Correct Answer: D (The patient's response to the care given)



- A) Check the mucous membranes in the eyes and mouth.
- B) Check skin condition.
- C) Take an oral temperature.
- D) Assess the pulse rate.

Correct Answer: A (Check the mucous membranes in the eyes and mouth.)

Q14. An eighty-five year old lady has fallen down a flight of stairs. You suspect spinal injury and decide to put the patient on a spineboard. You should be aware that:

- A) It may be difficult to fit a cervical collar.
- B) Spinal curvature may make it difficult and uncomfortable to put the patient on the board.
- C) A K.E.D. is better for this patient.
- D) Because of her age, she may be more prone to vomiting.

Correct Answer: B (Spinal curvature may make it difficult and uncomfortable to put the patient on the board.)

Q15. Which of the following is NOT a special consideration when assessing elderly patients:

- A) Geriatric patients often have a reduced gag reflex.
- B) Radial pulses may be weakened by peripheral vascular disease.
- C) Patients may not show signs of fever even in cases of serious infection.
- D) Mental status often reflects a fear of emergency responders. 31 Emergency Childbirth

Correct Answer: D (Mental status often reflects a fear of emergency responders. 31 Emergency Childbirth)

Q16. Crowning occurs:

- A) In the first stage of labour.
- B) In the second stage of labour.
- C) In the third stage of labour.
- D) In the fourth stage of pregnancy.

Correct Answer: B (In the second stage of labour.)

Q17. When suctioning a newborn you should use:

- A) bulb syringe.
- B) A V-Vac suction device.
- C) A straw.
- D) A battery powered suction device.

Correct Answer: A (bulb syringe.)

Q18. To help control bleeding after the baby is born:

- A) Pack the vagina with pads.
- B) Massage the mother's uterus.
- C) Elevate the mother's legs above heart level.
- D) Place the mother on her right side. 32 Pediatric Emergencies

Correct Answer: B (Massage the mother's uterus.)



- A) Blood pressure.
- B) Pain.
- C) Adequate perfusion.
- D) Level of consciousness.

Correct Answer: B (Pain.)

Q20. The onset of shock in infants and children:

- A) May be sudden and severe.
- B) Is no different than adults.
- C) Usually progresses slowly.
- D) Usually results from cardiac problems.

Correct Answer: A (May be sudden and severe.)

Q21. Assessments of children should take into consideration:

- A) No splints should be used since all are designed for adult patients.
- B) Abdominal injuries are less serious since muscles are less developed.
- C) The skull has not completely fused, so head injuries are uncommon.
- D) Chest injury is likely to involve organs without damage to ribs.

Correct Answer: D (Chest injury is likely to involve organs without damage to ribs.)

Q22. The responder who suspects a child to be the victim of abuse should:

- A) Question parents immediately.
- B) Focus on management of injuries to the child.
- C) Gather evidence and question bystanders.
- D) Immediately remove the child from parents or caregivers.

Correct Answer: B (Focus on management of injuries to the child.)

Q23. A child with history of a sore throat, fever and painful swallowing, has breathing difficulties. You should:

- A) Arrange for immediate transport to a medical facility.
- B) Start mouth-to-mouth artificial respiration.
- C) Stand by and encourage coughing.
- D) Begin abdominal thrusts.

Correct Answer: A (Arrange for immediate transport to a medical facility.)

Q24. The mammalian reflex refers to:

- A) An increase in heart rate and respiration.
- B) A decrease in heart rate and dilation of blood vessels.
- C) An increase in heart rate and constriction of pupils.
- D) A decrease in heart rate and constriction of blood vessels.

Correct Answer: D (A decrease in heart rate and constriction of blood vessels.)



- A) Air embolism, decompression sickness and mammalian reflex.
- B) Air embolism, decompression sickness and barotrauma.
- C) Decompression sickness, the bends and barotrauma.
- D) Air embolism, barotrauma and the squeeze.

Correct Answer: B (Air embolism, decompression sickness and barotrauma.)

Q26. In cases of near drowning:

- A) The lungs are often filled with water and require deep suctioning.
- B) Laryngospasm prevents water from entering the lungs.
- C) Abdominal thrusts should always be performed before ventilating the patient.
- D) Manual suction devices are not powerful enough to remove water and fluids from the airway.

Correct Answer: B (Laryngospasm prevents water from entering the lungs.)

Q27. You have been helping at a water rescue when one of the divers begins to have problems. You suspect air embolism. This patient should receive:

- A) Oxygen at 15 lpm.
- B) Oxygen at 6 lpm.
- C) Epinephrine.
- D) Inhaled steroids.

Correct Answer: A (Oxygen at 15 lpm.)

Q28. To reduce the risk of respiratory infection and pneumonia where medical care is delayed, responders should:

- A) Have the patient breathe deeply and cough.
- B) Provide the patient with adequate hydration.
- C) Position the patient in the position of comfort.
- D) Ensure the patient is kept warm and at rest.

Correct Answer: A (Have the patient breathe deeply and cough.)

Q29. When positioning a patient, responders should consider that:

- A) Raising arms and legs will produce swelling.
- B) Semi-sitting will improve blood flow to the heart.
- C) Knees raised will reduce pressure on the abdomen.
- D) Recovery position will often improve breathing.

Correct Answer: C (Knees raised will reduce pressure on the abdomen.)

Q30. When caring for wounds over extended periods, always:

- A) Remove dressings and bandages regularly to check the wound.
- B) Clean wounds with antiseptic before applying dressings.
- C) Apply a tourniquet to prevent excessive bleeding.
- D) Check regularly for signs of infection or reduced circulation.

Correct Answer: D (Check regularly for signs of infection or reduced circulation.)



Practice Set 7

Q1. The maintenance of fluids and nutrition is key to extended patient care and can be provided by:

- A) Giving fluids and food to all patients regardless of MOI.
- B) Use small amounts of liquids and foods to determine whether the patient can tolerate the intake.
- C) Allow the patient to consume any foods they would like to make them feel more comfortable.
- D) Even if the patient is not thirsty or hungry the responder must insist on intake.

Correct Answer: B (Use small amounts of liquids and foods to determine whether the patient can tolerate the intake.)

Q2. When administering medication, responders should consider that:

- A) Right medication, right amount, right person, right time, right method.
- B) Right nutrition, , right method, right time, right person, right amount
- C) Right medication, right action, right person, right time, right responsibility
- D) Right medication, right amount, right person, right responsibility, right action

Correct Answer: A (Right medication, right amount, right person, right time, right method.)

Q3. To give ear drops to an adult, the ear should be pulled:

- A) Downward and backward.
- B) Downward and forward.
- C) Upward and backward.
- D) Upward and forward.

Correct Answer: C (Upward and backward.)

Q4. Provincial first aid legislation prescribes the contents of:

- A) First aid boxes, first aid rooms and first aid manuals.
- B) First aid stations, training programs and equipment manufacturers.
- C) First aid boxes, first aid reports and record keeping.
- D) First aid boxes, first aid rooms and training.

Correct Answer: D (First aid boxes, first aid rooms and training.)

Q5. First aid stations must:

- A) Be easily accessible to workers
- B) Have a shower
- C) Be at least 6 square meters in size
- D) Have oxygen available Ambulance Operation/Maintenance

Correct Answer: A (Be easily accessible to workers)

Q6. Footprints can best be described as:

- A) The area of contact between the road surface and the tires.
- B) The type of tread on the tire.
- C) Footsteps used to get into the back of the ambulance.
- D) Skid patterns on dry pavement.

Correct Answer: A (The area of contact between the road surface and the tires.)



Q7. Ground guides or spotters should be situated:

- A) Behind the vehicle on the driver side.
- B) On the front left side of the vehicle.
- C) Front and centre of the vehicle.
- D) On the front right side of the vehicle.

Correct Answer: A (Behind the vehicle on the driver side.)

Q8. When completing the patient narrative:

- A) Try to establish an accurate diagnosis.
- B) Record the signs and symptoms as accurately as possible.
- C) Do not include negative data.
- D) Always use medical terminology.

Correct Answer: B (Record the signs and symptoms as accurately as possible.)

Q9. A patient refuses your care. You should:

- A) Fill out and personally sign a "Refusal of Care" form.
- B) Proceed with patient assessment and management and ignore the refusal.
- C) Find a relative or family member to give you consent.
- D) Have the patient and a witness sign the "Refusal of Care" form.

Correct Answer: D (Have the patient and a witness sign the "Refusal of Care" form.)

Q10. Sublingual medication is given:

- A) By injection using an auto-injector.
- B) Under the patient's tongue.
- C) By inhalation using a metered dose inhaler.
- D) As a tablet to be swallowed.

Correct Answer: B (Under the patient's tongue.)

Q11. Contraindications of a medication tell responders:

- A) When a medication should not be given to the patient.
- B) When the medication is commonly used for patients.
- C) The expected results of the medication.
- D) Actions that might not be desirable yet occur along with desired effects.

Correct Answer: A (When a medication should not be given to the patient.)

Q12. The ACTION of the drug refers to:

- A) The harmful effects.
- B) The side effects.
- C) The expected effects.
- D) The method of administration.

Correct Answer: C (The expected effects.)



Q13. To manage severe toothaches, responders should:

- A) Advise the patient to chew on the other side of the mouth.
- B) Advise the patient to drink hot liquids to sooth discomfort.
- C) Advise the patient to suck on ice cubes to dull the pain.
- D) Extract the tooth with the assistance of a second responder.

Correct Answer: A (Advise the patient to chew on the other side of the mouth.)

Q14. When handling a knocked-out tooth you should:

- A) Place the tooth back in the patient's mouth.
- B) Place the tooth in a cup of milk.
- C) Wrap the tooth in dry gauze.
- D) Place the tooth in a mild acidic solution using vinegar.

Correct Answer: B (Place the tooth in a cup of milk.)

Q15. Lividity refers to:

- A) Stiffening of the joints post mortem.
- B) Cooling of the body post mortem.
- C) Settling of blood in the body due to gravity.
- D) The beginning signs of decomposition.

Correct Answer: C (Settling of blood in the body due to gravity.)

Q16. Later signs of death include:

- A) Staining, milky corneas and flushed skin.
- B) Staining, lividity and decomposition.
- C) Rigor mortis, lividity and milky corneas.
- D) Rigor mortis, flushed skin and decomposition.

Correct Answer: C (Rigor mortis, lividity and milky corneas.)