

Measure Theory

Name: AZMAIN BISWAS

Roll No.: 2022MAM008

Supervisor's Name: Dr. UJJAL DEBNATH

Mini Project for 1st semester course in

M.Sc. in Applied Mathematics



IEST, Shibpur
आई आई ई एस टि, शिवपुर
Erstwhile B E College (Estd 1856)

Department of Mathematics

**INDIAN INSTITUTE OF ENGINEERING SCIENCE AND
TECHNOLOGY, Shibpur**

DECLARATION: I affirm that I have identified all my sources and that no part of my dissertation paper uses unacknowledged materials.

ACKNOWLEDGEMENT

I would like to take this opportunity to express my profound gratitude to my supervisor Dr. Ujjal Debnath for his exemplary guidance and constant encouragement throughout the course of this project.

I am grateful to IEST, Shibpur for providing me with the opportunity to complete my dissertation. In this project I have tried to give a short overview of **Measure Theory**.

I would like to extend my deepest gratitude and thanks to my parents without whose constant support and encouragement this project would never have been completed. I would also like to thank my friends for their valuable suggestions and support.

Azmain Biswas

Abstract

Measure Theory is the study of Measure. Here we try extend the notion of size of interval to more general sets. We know the size of a interval in \mathbb{R} is its length. Then by instinct we say in \mathbb{R}^2 it will be area and in \mathbb{R}^3 it will be volume of sets. Finding more general notion of size of interval leads us to concept of measure.

We will start with outer measure even though it looks promising but it fails to have crucial property. This unfortunate result leads us to σ -algebra and measurable space. From there we construct measure in an abstract context. Taking restriction on outer measure we construct Lebesgue Measure which leads to more general theory of integration then Riemann integration.

Contents

1	Outer Measure	4
2	Measurable Space	8
3	Topology Space	8
4	Borel set	9
5	Measurable Function	10
5.1	Borel measurable function	13
6	Measure	16
7	Lebesgue Measure	20
8	Cantor Set	26
9	Vitali Set	28

1 Outer Measure

Definition 1.1 (length of open interval). The length $\ell(I)$ of an open interval is define by

$$\ell(I) = \begin{cases} b - a & \text{if } I = (a, b) \text{ for some } a, b \in \mathbb{R} \text{ with } a < b, \\ 0 & \text{if } I = \emptyset, \\ \infty & \text{if } I = (-\infty, a) \text{ or } I = (a, \infty) \text{ for some } a \in \mathbb{R} \\ \infty & \text{if } I = (-\infty, \infty). \end{cases}$$

Definition 1.2 (Outer Measure). The outer measure $\mu^*(A)$ of a set $A \subset \mathbb{R}$ is define by

$$\mu^*(A) = \inf \left\{ \sum_{k=1}^{\infty} \ell(I_k) \right\}$$

where, I_1, I_2, \dots are open intervals such that $A \subset \bigcup_{k=1}^{\infty} I_k$.

Example 1.1 (finite sets have outer measure 0). Suppose $A = \{a_1, a_2, \dots, a_n\}$ is a finite set of real numbers. Suppose $\epsilon > 0$. Define a sequence I_1, I_2, \dots of open intervals by

$$I_k = \begin{cases} (a_k - \epsilon, a_k + \epsilon) & \text{if } k \leq n; \\ \emptyset & \text{if } k > n. \end{cases}$$

Then I_1, I_2, \dots is a sequence of open intervals whose union contain A .

Clearly, $\sum_{k=1}^{\infty} \ell(I_k) = 2\epsilon n$.

Hence, $\mu^*(A) \leq 2\epsilon n$.

Since ϵ is arbitrary small positive number, $\mu^*(A) = 0$.

Properties of Outer Measure

Properties of outer measure are followings,

1. Every countable subset of \mathbb{R} has outer measure 0.
2. Suppose A and B are subset of \mathbb{R} with $A \subset B$. Then $\mu^*(A) \leq \mu^*(B)$.
3. Suppose $t \in \mathbb{R}$ and $A \subset \mathbb{R}$. Then $\mu^*(t + A) = \mu^*(A)$.
4. Suppose A_k 's are a sequence of subset of \mathbb{R} . Then,

$$\mu^* \left(\bigcup_{k=1}^{\infty} A_k \right) \leq \sum_{k=1}^{\infty} \mu^*(A_k).$$

5. Suppose $a, b \in \mathbb{R}$, with $a < b$. Then $\mu^*([a, b]) = b - a$.

Theorem 1.1 (outer measure is not additive). *There exist disjoint subsets A and B of \mathbb{R} such that,*

$$\mu^*(A \cup B) \neq \mu^*(A) + \mu^*(B).$$

Proof. For $a \in [-1, 1]$, let \tilde{a} be the set of numbers in $[-1, 1]$ such that.

$$\tilde{a} = \{c \in [-1, 1] : a - c \in \mathbb{Q}\}.$$

If $a, b \in [-1, 1]$ and $\tilde{a} \cap \tilde{b} \neq \emptyset$, Then $\tilde{a} = \tilde{b}$.

(*Proof:* Suppose There exist $d \in \tilde{a} \cap \tilde{b}$. Then $a - b$ and $b - d$ are rational number; subtracting, we conclude that $a - d$ is rational number. The equitation $a - c = (a - b) + (b - c)$ now implies that if $c \in [-1, 1]$, then $a - c$ is rational iff $b - c$ is rational number. In other word $\tilde{a} = \tilde{b}$.)

Clearly, $a \in \tilde{a}$ for each $a \in [-1, 1]$. Thus $[-1, 1] = \cup_{a \in [-1, 1]} \tilde{a}$.

Let V be a set that contains exactly one element in each of the distinct sets in $\{\tilde{a} : a \in [-1, 1]\}$

In other words, for every $a \in [-1, 1]$, the set $V \cap \tilde{a}$ has exactly one element.

Let r_1, r_2, \dots be a sequence of distinct rational numbers such that

$$[-2, 2] \cap \mathbb{Q} = r_1, r_2, \dots$$

Then

$$[-1, 1] \subset \bigcup_{k=1}^{\infty} (r_k + V),$$

Where the set inclusion above holds because if $a \in [-1, 1]$ then v is the unique element of $V \cap \tilde{a}$, we have $a - v \in \mathbb{Q}$, which implies that $a = r_k + v \in r_k + V$ for some $k \in \mathbb{Z}^+$.

Then by order-preserving property of outer measure, and countable subadditivity of outer measure imply

$$\mu^*([-1, 1]) \leq \sum_{k=1}^{\infty} \mu^*(r_k + V).$$

Now $\mu^*([-1, 1]) = 2$ and by translation invariance property of outer measure we can say,

$$2 \leq \sum_{k=1}^{\infty} \mu^*(V).$$

Thus $\mu^*(V) > 0$.

Note that the sets $r_k + V$'s are disjoint.

(*Proof:* Suppose there exists $t \in (r_j + V) \cap (r_k + V)$. Then $t = r_j + v_1 = r_k + v_2$ for some $v_1, v_2 \in V$, which implies that $v_1 - v_2 = r_k - r_j \in \mathbb{Q}$ from the construction of V now implies that $v_1 = v_2 \implies r_j = r_k \implies j = k$.) Let $n \in \mathbb{Z}^+$. Clearly,

$$\bigcup_{k=1}^n (r_k + V) \subset [-3, 3]$$

Because $V \subset [-1, 1]$ and each $r_k \in [-2, 2]$. The set inclusion above implies that

$$\mu^*\left(\bigcup_{k=1}^n (r_k + V)\right) \leq 6 \tag{1}$$

However,

$$\sum_{k=1}^n \mu^*(r_k + V) = \sum_{k=1}^n \mu^*V = n\mu^*(V). \tag{2}$$

Now (1) and (2) suggest that we choose $n \in \mathbb{Z}^+$ such that $n\mu^*(V) > 6$. Thus,

$$\mu^* \left(\bigcup_{k=1}^n (r_k + V) \right) < \sum_{k=1}^n \mu^*(r_k + V). \quad (3)$$

If we had $\mu^*(A \cup B) = \mu^*(A) + \mu^*(B)$ for all disjoint subset A, B of \mathbb{R} , then by induction on n we would have $\mu^*(\bigcup_{k=1}^n A_k) = \sum_{k=1}^n \mu^*(r_k + V)$. For all disjoint subset A_1, A_2, \dots, A_n of \mathbb{R} .

However, from (3) we see no such result holds. Thus, there exist disjoint subsets on A, B of \mathbb{R} such that $\mu^*(A \cup B) \neq \mu^*(A) + \mu^*(B)$. \square

Now we have seen in above theory that outer measure is not additive. So does there exist other notion than outer measure for size of subsets of \mathbb{R} that follow additive property?

Theorem 1.2 (nonexistence of extension of length to all subset of \mathbb{R}). *There does not exist a function μ with all the following properties:*

1. $\mu : P(\mathbb{R}) \rightarrow [0, \infty]$.
2. $\mu(I) = \ell(I)$ for every open interval $I \subset \mathbb{R}$.
3. $\mu(\bigcup_{k=1}^{\infty} A_k) = \sum_{k=1}^{\infty} \mu(A_k)$ for every disjoint sequence $\{A_k\}$ of subsets of \mathbb{R} .
4. $\mu(t + A) = \mu(A)$ for every $A \subset \mathbb{R}$ and for every $t \in \mathbb{R}$.

Proof. Suppose there exist a function μ that follow all the property listed above.

Observed that $\mu(\emptyset) = 0$, as follows from (2) because the empty set is a open interval with length 0.

If $A \subset B \subset \mathbb{R}$, then $\mu(A) \leq \mu(B)$, as follows from (3) because we can write B as the union of the disjoint sequence $A, B \setminus A, \emptyset, \emptyset, \dots$;

Thus,

$$\mu(B) = \mu(A) + \mu(B \setminus A) + 0 + 0 + \dots = \mu(A) + \mu(B \setminus A) \geq \mu(A).$$

If $a, b \in \mathbb{R}$ with $a < b$, then $(a, b) \subset [a, b] \subset (a - \epsilon, b + \epsilon)$ for every $\epsilon > 0$. Thus $b - a \leq \mu([a, b]) \leq b - a + 2\epsilon$ for every $\epsilon > 0$. Hence $\mu([a, b]) = b - a$.

If A_k is sequence of subset of \mathbb{R} , then $A_1, A_2 \setminus A_1, A_3 \setminus (A_1 \cup A_2), \dots$ is a disjoint sequence of \mathbb{R} whose union is $\bigcup_{k=1}^{\infty} A_k$. Thus,

$$\begin{aligned} \mu \left(\bigcup_{k=1}^{\infty} A_k \right) &= \mu(A_1 \cup (A_2 \setminus A_1) \cup (A_3 \setminus (A_1 \cup A_2)) \cup \dots) \\ &= \mu(A_1) + \mu(A_2 \setminus A_1) + \mu(A_3 \setminus (A_1 \cup A_2)) + \dots \\ &\leq \sum_{k=1}^{\infty} \mu(A_k), \end{aligned}$$

We have shown that μ has all the properties of outer measure that were used in the proof of (1.1). Repeating the proof of 1.1, we see that there exist disjoint subset A, B of \mathbb{R} such that $\mu(A \cup B) \neq \mu(A) + \mu(B)$. Which is a contradiction of (3). This contradiction completes the proof. \square

So the above result shows us we have to give up one of the described properties in our goal of extending the notion of size from intervals to more general sets. We can not give up property (2) because the size of an interval needs to be its length. We cannot give up property (3) because additivity is needed. We can not give up (4) either because length have to be translation invariant.

So we are force to relax the property (1) that size is defined for all subset of \mathbb{R}

2 Measurable Space

Definition 2.1 (σ -algebra). Suppose X is a non empty set and \mathcal{A} is a set of subset of X . Then \mathcal{A} is called σ -algebra on X if the following three condition are satisfied:

1. $\emptyset \in \mathcal{A}$,
2. if $E \in \mathcal{A}$, than $E^c \in \mathcal{A}$. (*close under complementation*)
3. if $\{E_k\}$ is a sequence of element of \mathcal{A} , then $\bigcup_{k=1}^{\infty} E_k \in \mathcal{A}$. (*closed under countable union*)

Theorem 2.1. Suppose \mathcal{A} is a σ -algebra on a set X . Then

1. $X \in \mathcal{A}$
2. If $\{E_k\}$ is a sequence of element of \mathcal{A} , then $\bigcap_{k=1}^{\infty} E_k \in \mathcal{A}$. (*closed under countable intersection*)

Proof. Because $\emptyset \in \mathcal{A}$ and \mathcal{A} is closed under complementation so, $X \in \mathcal{A}$.

Now, E_1, E_2, \dots is in \mathcal{A}

then, $E_k^c \in \mathcal{A}$ for all $k = 1, 2, \dots$

Then,

$$\begin{aligned} \bigcup_{k=1}^{\infty} E_k^c &\in \mathcal{A}. \\ \implies \left(\bigcap_{k=1}^{\infty} E_k \right)^c &\in \mathcal{A} \\ \implies \bigcap_{k=1}^{\infty} E_k &\in \mathcal{A}. \end{aligned}$$

This proves the result. □

Definition 2.2 (measurable space). A measurable space is an ordered pair (X, \mathcal{A}) , where X is a set and \mathcal{A} is a σ -algebra on X .

Definition 2.3 (measurable set). An element of \mathcal{A} is called an \mathcal{A} -measurable set or just measurable set.

3 Topology Space

Definition 3.1 (Topology). A collection τ of subsets of non-empty set X is said to be topology in X if τ satisfied the following conditions

1. $\emptyset \in \tau$ and $X \in \tau$.
2. if $V_i \in \tau$ for all $i = 1, 2, \dots, n$, then $\bigcap_{i=1}^n V_i \in \tau$.
3. if $\{V_\alpha\}$ is an arbitrary collection of members of τ (finite, countable, uncountable), then $\bigcup_\alpha V_\alpha \in \tau$.

Definition 3.2 (topology space). If τ is a topology in X , Then an ordered pair (X, τ) is called topology space.

Definition 3.3 (Open set). The member of τ is called Open set in X .

Definition 3.4 (close set). E is said to be closed set if $E^c \in \tau$.

4 Borel set

Theorem 4.1. If \mathcal{F} is any collection of subsets of X , there exists a smallest σ -algebra \mathcal{B} in X such that $\mathcal{F} \in \mathcal{B}$

This \mathcal{B} is sometime called the σ -algebra generated by \mathcal{F} .

Proof. Let Ω be the family σ -algebras \mathcal{B}^* in X which contain \mathcal{F} . Since the collection of all subsets of X is such σ -algebra (i.e. $P(X)$ is a σ -algebra), Ω is non empty.

Let \mathcal{B} is intersection of all $\mathcal{B}^* \in \Omega$. It is clear that $\mathcal{F} \subset \mathcal{B}$ and \mathcal{B} lies in every σ -algebra in X which contain \mathcal{F} .

To Complete the proof, we have to show that \mathcal{B} is a σ -algebra.

If $A_n \in \mathcal{B}$ for $n = 1, 2, \dots$ and if $\mathcal{B}^* \in \Omega$, then $A_n \in \mathcal{B}^*$ so, $\bigcup_{n=1}^{\infty} A_n \in \mathcal{B}^*$, since \mathcal{B}^* is a σ -algebra

Since $\bigcup_{n=1}^{\infty} A_n \in \mathcal{B}^*$ for every $\mathcal{B}^* \in \Omega$, we conclude that $\bigcup_{n=1}^{\infty} A_n \in \mathcal{B}$. The other two properties of σ -algebra are verified in the same manner. \square

Definition 4.1 (Borel set). Let X be a topology space. By theorem 4.1 there exist a smallest σ -algebra \mathcal{B} in X such that every open set in X belongs to \mathcal{B} . The member of \mathcal{B} is called Borel sets of X .

In particular: The σ -algebra on \mathbb{R} containing all open subset of \mathbb{R} is called the collection of Borel subset of \mathbb{R} . An element of this σ -algebra is called a *Borel Set*.

Example 4.1.

- Every closed subset of \mathbb{R} is a Borel set because every closed set are complement of an open subset of \mathbb{R} .
- Every countable subset of \mathbb{R} is a Borel set because countable sets are closed set.
- Every half-open interval $[a, b)$ (where $a, b \in \mathbb{R}$) is a Borel sets because

$$[a, b) = \bigcap_{n=1}^{\infty} \left(a - \frac{a}{n}, b \right).$$

- All countable union of closed sets and all countable intersection of open set are Borel set. This sets are called F_σ and G_δ respectively.

5 Measurable Function

Definition 5.1 (Measurable Function). Let (X, \mathcal{A}) and (Y, \mathcal{S}) be two measurable spaces and a function $f : X \rightarrow Y$ is said to be \mathcal{A} -measurable function (or just measurable function) provided that $f^{-1}(V)$ is a measurable set in X for every measurable set V in Y .

i.e.

$$f^{-1}(V) \in \mathcal{A} \quad \text{for every } V \in \mathcal{S}$$

In particular: Suppose (X, \mathcal{A}) is a measurable space. A function $f : X \rightarrow \mathbb{R}$ is called \mathcal{A} -measurable if $f^{-1}(B) \in \mathcal{A}$ for every Borel set $B \subset \mathbb{R}$.

Example 5.1.

- If \mathcal{A} is a set of all subset of X , then every function from $X \rightarrow \mathbb{R}$ is a \mathcal{A} -measurable.
- If $\mathcal{A} = \{\emptyset, (-\infty, 0), [0, \infty), \mathbb{R}\}$ (which is a σ -algebra of \mathbb{R}), then constant function $f : \mathbb{R} \rightarrow \mathbb{R}$ is an \mathcal{A} -measurable function.

Definition 5.2 (characteristic function). Suppose E is a subset of X . The characteristic function of E is a function $\chi_E : X \rightarrow \mathbb{R}$ define by

$$\chi_E(x) = \begin{cases} 1 & \text{if } x \in E \\ 0 & \text{if } x \notin E \end{cases}$$

Observation: Suppose (X, \mathcal{A}) is a measurable space, $E \subset X$, and $B \subset \mathbb{R}$. Then,

$$\chi_E^{-1}(B) = \begin{cases} E & \text{if } 0 \notin B \text{ and } 1 \in B. \\ X \setminus E & \text{if } 0 \in B \text{ and } 1 \notin B. \\ X & \text{if } 0 \in B \text{ and } 1 \in B. \\ \emptyset & \text{if } 0 \notin B \text{ and } 1 \notin B. \end{cases}$$

thus we see that χ_E is an \mathcal{A} -measurable function iff $E \in \mathcal{A}$.

Definition 5.3 (Simple Function). Let ψ be real-valued function defined on X . If the range of ψ is finite, we say that ψ is a simple function.

Suppose the range of ψ consists of the distinct number c_1, \dots, c_n . Let,

$$E_i = \{x : \psi(x) = c_i\} \quad (i = 1, \dots, n)$$

Then,

$$\psi(x) = \sum_{k=1}^n c_k \chi_{E_k}.$$

i.e. every simple function is a finite linear combination of characteristic functions. Hence it is clear that ψ is measurable iff the sets E_1, \dots, E_n are measurable.

It is of interest that every function can be approximated by simple function.

Theorem 5.1 (Approximation by simple function). $f : (X, \mathcal{A}) \rightarrow \mathbb{R}^+$ is a non-negative measurable function. Then there exists a sequence of simple function $\{\psi_n\}$ such that.

1. ψ is monotone increasing.

2. $\lim \psi_n = f$.

Proof. Let k be any integer. We write,

$$E_{p,k} = \left\{ x : \frac{p-1}{2^k} \leq f(x) \leq \frac{p}{2^k} \right\} \text{ where } p = 1, 2, \dots, 2^{2k}.$$

and

$$E_{0,k} = X \setminus \bigcup_{p=1}^{2^{2k}} E_{p,k} = \left\{ x : f(x) \geq 2^k \right\}.$$

Then,

$$E_{p,k} = \left\{ x : f(x) < \frac{p}{2^k} \right\} \cap \left\{ x : f(x) \geq \frac{p-1}{2^k} \right\}.$$

is measurable, Since f is measurable. Also, $E_{p,k}$, $p = 0, 1, 2, \dots, 2^{2k}$ are mutually disjoint. we construct,

$$\psi_k(x) = \begin{cases} \frac{p-1}{2^k} & \text{for } x \in E_{p,k} \\ 2^k & \text{for } x \in E_{0,p}. \end{cases}$$

the ψ_k is a step function.

By construction $0 \leq \psi_k \leq f$.

If $x \in E_{p,k}$, $p \neq 0$ then we have the following cases.

Case:1

$x \in E_{2p-1,k+1}$ in that cases $\psi_k(x) = \psi_{k+1}$

Case:2

$x \in E_{2p,k+1}$ is that case $\psi_k(x) + \frac{1}{2^{k+1}} = \psi_{k+1}(x)$.

Thus for $x \in E_{p,k}$. Then $\psi_n(x) \leq \psi_{n+1}(x)$.

Also, if $x \in E_{0,p}$. Then $\psi_k(x) = 2^k \leq 2^{k+1} = \psi_{k+1}(x)$. Combining we get,

$$\psi_k(x) \leq \psi_{k+1}(x) \quad \forall x \in X$$

i.e. ψ_k is monotone increasing.

Next we show $\psi_k \rightarrow f$.

Case:1 f is finite.

then $f(x) < 2^k$ for some $k \in \mathbb{Z}^+$, in that case,

$$0 \leq f(x) - \psi_k(x) < 2^k - \frac{2^{2k} - 1}{2^k} = \frac{1}{2^k}.$$

which show that $\psi_k(x) \rightarrow f(x)$ as $k \rightarrow \infty$.

Case:2 $f = +\infty$

then $\psi_k(x) = 2^k$ and $\psi_k \rightarrow f = \infty$ as $k \rightarrow \infty$.

Combination both we get,

$$\psi_n \rightarrow f \text{ as } n \rightarrow \infty.$$

□

Theorem 5.2. Suppose (X, \mathcal{S}) is a measurable space and $f : X \rightarrow \mathbb{R}$ is a measurable

function iff

$$f^{-1}((a, \infty)) = \{x \in X : f(x) > a\} \in \mathcal{S} \quad \forall a \in \mathbb{R}$$

Proof. Let f is a measurable function.

Since $\forall a \in \mathbb{R}$, (a, ∞) is a Borel set of \mathbb{R} . $f^{-1}((a, \infty)) \in \mathcal{S}$

Conversely, let $f^{-1}((a, \infty)) \in \mathcal{S}$

We have to prove that f is a measurable.

let, $\tau = \{A \subset \mathbb{R} : f^{-1}(A) \in \mathcal{S}\}$

We have to show every Borel subset of \mathbb{R} is in τ . To do this first we prove τ is a σ -algebra on \mathbb{R} .

Clearly, $\emptyset \in \tau$, because $f^{-1}(\emptyset) = \emptyset \in \mathcal{S}$.

If $A \in \tau$, then $f^{-1}(A) \in \mathcal{S}$; hence,

$$f^{-1}(\mathbb{R} \setminus A) = X \setminus f^{-1}(A) \in \mathcal{S}.$$

Thus, $\mathbb{R} \setminus A \in \tau$. In other word τ is closed under complementation.

If $A_1, A_2, \dots \in \tau$

then, $f^{-1}(A_1), f^{-1}(A_2), \dots \in \mathcal{S}$.

hence,

$$f^{-1}\left(\bigcup_{k=1}^{\infty} A_k\right) = \bigcup_{k=1}^{\infty} f^{-1}(A_k) \in \mathcal{S}.$$

Thus, $\bigcup_{k=1}^{\infty} A_k \in \tau$. In other word τ is closed under countable unions. Thus τ is σ -algebra on \mathbb{R} .

By hypothesis, τ contains $\{(a, \infty) : a \in \mathbb{R}\}$. Since τ is closed under complementation, τ contains $\{(-\infty, b] : b \in \mathbb{R}\}$. τ also contain $\{(a, b] : a, b \in \mathbb{R}\}$, as τ is closed under countable intersection. Now $(a, b) = \bigcup_{k=1}^{\infty} (a, b - \frac{1}{k}]$. and $(-\infty, b) = \bigcup_{k=1}^{\infty} (-k, b - \frac{1}{k}]$. Hence, τ contain every open set of \mathbb{R} .

Thus the σ -algebra τ contains the smallest σ -algebra on \mathbb{R} that contains all open subset of \mathbb{R} . Hence τ contains every borel set of \mathbb{R} . Thus f is a measurable function. \square

Theorem 5.3. If (X, \mathcal{A}) is a measurable space and a function $f : X \rightarrow \mathbb{R}$ is measurable if any of the following holds for each $a \in \mathbb{R}$,

1. $\{x : f(x) \geq a\} = f^{-1}[a, \infty) \in \mathcal{A}$.
2. $\{x : f(x) \leq a\} = f^{-1}(-\infty, a] \in \mathcal{A}$.
3. $\{x : f(x) < a\} = f^{-1}(-\infty, a) \in \mathcal{A}$.

Proof.

1. Let f is a measurable function from $X \rightarrow \mathbb{R}$.

Then, $f^{-1}((a, \infty)) \in \mathcal{A}$

i.e. $\{x : f(x) > a\} \in \mathcal{A} \quad \forall a \in \mathbb{R}$. now,

$$\left\{x : f(x) > a + \frac{1}{k}\right\} \in \mathcal{A}$$

Then,

$$\bigcap_{k=1}^{\infty} \left\{ x : f(x) > a + \frac{1}{k} \right\} = \{x : f(x) \geq a\} \in \mathcal{A}$$

As, \mathcal{A} is a σ -algebra.

Conversely, Let f satisfy (1). Then, $\{x : f(x) \geq a\} \in \mathcal{A} \quad \forall a \in \mathbb{R}$.

now,

$$\left\{ x : f(x) \geq a - \frac{1}{k} \right\} \in \mathcal{A}.$$

then,

$$\bigcup_{k=1}^{\infty} \left\{ a : f(x) \geq a - \frac{1}{k} \right\} = \{a : f(x) > a\} \in \mathcal{A}.$$

as, \mathcal{A} is σ -algebra. Hence, f is a measurable function.

2. $\{x : f(x) \leq a\} = X \setminus \{x : f(x) > a\} \in \mathcal{A}$ as, \mathcal{A} is a σ -algebra.

3. $\{x : f(x) < a\} = X \setminus \{x : f(x) \geq a\} \in \mathcal{A}$

□

5.1 Borel measurable function

Definition 5.4 (Borel measurable function). A function $f : X(\subset \mathbb{R}) \rightarrow \mathbb{R}$ is called Borel measurable if $f^{-1}(B)$ is a Borel set for every Borel set $B \subset \mathbb{R}$

Theorem 5.4 (Continuous function is Borel measurable). *Every continuous real-valued function defined on a Borel subset of \mathbb{R} is a Borel measurable function.*

Proof. Let $X \subset \mathbb{R}$ is a Borel set and $f : X \rightarrow \mathbb{R}$ is a continuous function.

To prove that f is a Borel measurable function.

Since, f is continuous in X , $f^{-1}(A)$ is a open set whenever $A \cap X$ is open set. Hence, if A is Borel set then $f^{-1}(A)$ is also Borel set. Hence, f is Borel measurable function. □

Theorem 5.5 (Increasing function is Borel measurable). *Every increasing function define on a Borel subset of \mathbb{R} is a Borel measurable function.*

Proof. Suppose $X \subset \mathbb{R}$ and $f : X \rightarrow \mathbb{R}$ is a increasing function. To prove f is a Borel measurable function.

we fix $a \in \mathbb{R}$. Let $b = \inf f^{-1}((a, \infty))$, Then it is easy to see that,

$$f^{-1}((a, \infty)) = (b, \infty) \cap X \text{ or } f^{-1}((a, \infty)) = [b, \infty) \cap X$$

Either way, we can conclude that $f^{-1}((a, \infty))$ is a Borel set.

Hence f is a Borel measurable set. □

Theorem 5.6 (composition of measurable functions). *Suppose (X, \mathcal{A}) , (Y, \mathcal{S}) , (Z, \mathcal{B}) are measurable spaces. And $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ are \mathcal{A} -measurable and \mathcal{S} -measurable function respectively. Then $g \circ f : X \rightarrow Z$ is a \mathcal{A} -measurable function.*

Proof. Suppose $B \in \mathcal{B}$. Then ,

$$(g \circ f)^{-1}(B) = f^{-1}(g^{-1}(B)).$$

Since g is a measurable function, $g^{-1}(B) \in \mathcal{S}$. Then $f^{-1}(g^{-1}(B)) \in \mathcal{A}$, as f is a \mathcal{A} -measurable function.

Thus $(g \circ f)^{-1}(B) \in \mathcal{A}$, implies that $g \circ f$ is an \mathcal{A} -measurable set. \square

Corollary 5.6.1. *If f is measurable then $-f, kf, |f|, f^2$ is measurable where $k \in \mathbb{R}$ is a constant.*

Proof. If we choose $g(x) = -x$ then $g(x) = kx$ then $g(x) = |x|$ and then $g(x) = x^2$. Clearly all the $g(x)$'s are continuous hence measurable.

Then $g \circ f$ is measurable. Hence the result. \square

Theorem 5.7. *Suppose (X, \mathcal{A}) is a measurable space and $f, g : X \rightarrow \mathbb{R}$ are \mathcal{A} -measurable. Then,*

1. $f + g, f - g$ and fg is \mathcal{A} -measurable function.
2. If $g(x) \neq 0$ for all $x \in X$, Then $\frac{f}{g}$ is an measurable function.

Proof. We first proof $f + g$ is a measurable.

Let r is a rational numbers. Then for $c \in \mathbb{R}$

$$\{x : f(x) + g(x) > c\} = \bigcup_{r \in \mathbb{Q}} \{x : f(x) > r\} \cap \{x : g(x) > c - r\}. \quad (4)$$

(Because: If $f(x) + g(x) > c$. Hence $(c - g(x), f(x))$ is non empty and thus it contain some ration number r .

This implies that $r < f(x)$ and $c - g(x) < r$.

i.e. $f(x) > r$ and $g(x) > c - r$.

i.e. $x \in \{x : f(x) > r\} \cap \{x : g(x) > c - r\}$ for some $r \in \mathbb{Q}$.)

Now both f, g are measurable function. Then $\{x : f(x) > r\}$ and $\{x : g(x) > c - r\}$ are measurable.

Then $\{x : f(x) > r\} \cap \{x : g(x) > c - r\}$ is also measurable.

Thus $\{x : f(x) + g(x) > c\} = \bigcup_{r \in \mathbb{Q}} \{x : f(x) > r\} \cap \{x : g(x) > c - r\}$ is measurable.

Hence $f + g$ is also measurable.

Now if g is measurable then by 5.6.1 $-g$ is also measurable. Thuse $f + (-g) = f - g$ is also measurable function.

Now

$$fg = \frac{(f + g)^2 - f^2 - g^2}{2}.$$

Then by 5.6.1 fg is also measurable.

If $g(x) \neq 0 \forall x \in \mathbb{R}$. Then the function $h(x) = \frac{1}{x}$ define on $\mathbb{R} \setminus \{0\} \rightarrow \mathbb{R}$ is continuous. Hence $\frac{1}{g}$ is a measurable function. Then $f \times \frac{1}{g} = \frac{f}{g}$ is a measurable function. \square

Corollary 5.7.1. *If f and g is measurable function. Then $\max(f, g)$ and $\min(f, g)$ and measurable functions.*

In particular, $f^+ = \max(f, 0)$ and $f^- = -\min(f, 0)$ is also measurable function.

Proof. we know,

$$\max(f, g) = \frac{(f + g) + |f - g|}{2} \quad \min(f, g) = \frac{(f + g) - |f - g|}{2}. \quad (5)$$

then by 5.7 this result holds. \square

Theorem 5.8 (limits of measurable function). *Let $\{f_n\}$ be a sequence of measurable function (with the same domain of definition).*

Then $\sup\{f_1, \dots, f_n\}$, $\inf\{f_1, \dots, f_n\}$, $\sup_n f_n$, $\inf_n f_n$, $\limsup f_n$ and $\liminf f_n$ are measurable function.

Proof. let, $h(x) = \sup\{f_1, \dots, f_n\}$, then,

$$\{x : h(x) < a\} = \bigcap_{i=1}^n \{x : f_i(x) < a\}.$$

Since each f_i 's are measurable, $h(x)$ is also measurable.

Let, $g(x) = \inf\{f_1, \dots, f_n\}$, then

$$\{x : g(x) > a\} = \bigcap_{i=1}^n \{x : f_i(x) > a\}.$$

Since each f_i 's are measurable, $g(x)$ is also measurable.

Now,

$$\begin{aligned} \sup_n f_n &= \lim_{n \rightarrow \infty} \{x : h(x) < a\} = \bigcap_{i=1}^{\infty} \{x : f_i(x) < a\} \\ \inf_n f_n &= \lim_{n \rightarrow \infty} \{x : g(x) > a\} = \bigcap_{i=1}^{\infty} \{x : f_i(x) > a\} \end{aligned}$$

By measurability of f_i 's $\sup_n f_n$ and $\inf_n f_n$ is measurable.

Now,

$$\limsup f_n = \inf_n \left(\sup_{k \geq n} f_k \right) \quad \text{and} \quad \liminf f_n = \sup_n \left(\inf_{k \geq n} f_k \right)$$

Then $\limsup f_n$ and $\liminf f_n$ is also measurable. \square

6 Measure

Definition 6.1 (Measure). Suppose (X, \mathcal{A}) is a measurable space. A *measure* on (X, \mathcal{A}) is a function $\mu : \mathcal{A} \rightarrow [0, \infty]$ such that

1. $\mu(\emptyset) = 0$
2. For every disjoint sequence of sets $\{E_k\}$ in \mathcal{A} .

$$\mu \left(\bigcup_{k=1}^{\infty} E_k \right) = \sum_{k=1}^{\infty} \mu(E_k)$$

Example 6.1.

- If X is a set, then *counting measure* is a measure μ defined on the σ -algebra of all subsets of X by

$$\mu(E) = \begin{cases} n & \text{if } E \text{ is a finite set containing exactly } n \text{ element.} \\ \infty & \text{if } E \text{ is not a finite set.} \end{cases}$$

- (X, \mathcal{A}) is a measurable space and $c \in X$. Define the *Dirac measure* δ_c on (X, \mathcal{A}) by

$$\delta_c(E) = \begin{cases} 1 & \text{if } c \in E, \\ 0 & \text{if } c \notin E. \end{cases}$$

- Let, Ω is sample space and Σ is σ -algebra on Ω . Then $P : \Sigma \rightarrow [0, 1]$ is a measure on (Ω, Σ) such that $P(\Omega) = 1$ then, P is called probability measure and (Ω, Σ, P) is called probability space.

Definition 6.2 (measure space). A measure space is an ordered triple (X, \mathcal{A}, μ) , where X is a nonempty set, \mathcal{A} is a σ -algebra on X , and μ is a measure on (X, \mathcal{A}) .

Properties of measure

Theorem 6.1 (measure preserves order or monotonic property). Suppose (X, \mathcal{A}, μ) is a measure space and $D, E \in \mathcal{A}$ are such that $D \subset E$. Then,

1. $\mu(D) \leq \mu(E)$,
2. $\mu(E \setminus D) = \mu(E) - \mu(D)$ provided that $\mu(D) < \infty$.

Proof. We have $E = D \cup (E \setminus D)$ and this is a disjoint union

$$\mu(E) = \mu(D) + \mu(E \setminus D) \geq \mu(D). \quad (6)$$

which proves (1).

If $\mu(D) < \infty$, Then subtracting $\mu(D)$ from both side of equation (6) proves (2) also. \square

Theorem 6.2. Suppose (X, \mathcal{A}, μ) is a measure space and $\{E_k\} \in \mathcal{A} \forall k \in \mathbb{N}$. Then,

$$\mu \left(\bigcup_{k=1}^{\infty} E_k \right) \leq \sum_{k=1}^{\infty} \mu(E_k).$$

Proof. Let, $D_1 = \emptyset$ and $D_k = E_1 \cup \dots \cup E_k$ for $k \geq 2$. Then,

$$E_1 \setminus D_1, E_2 \setminus D_2, E_3 \setminus D_3, \dots$$

is a disjoint sequence of subsets of X whose union equals $\bigcup_{k=1}^{\infty} E_k$. Thus


$$\begin{aligned} \mu \left(\bigcup_{k=1}^{\infty} E_k \right) &= \mu \left(\bigcup_{k=1}^{\infty} (E_k \setminus D_k) \right) \\ &= \sum_{k=1}^{\infty} \mu(E_k \setminus D_k) \\ &\leq \sum_{k=1}^{\infty} \mu(E_k) \quad (\text{By theorem 6.1}) \end{aligned}$$

This proves the result. □

Theorem 6.3. Suppose (X, \mathcal{A}, μ) is a measure space and $E_1 \subset E_2 \subset \dots$ is an increasing sequence of sets in \mathcal{A} . Then

$$\mu \left(\bigcup_{k=1}^{\infty} E_k \right) = \lim_{n \rightarrow \infty} \mu(E_n).$$

Proof.



If $\mu(E_k) = \infty$ for some $k \in \mathbb{N}$, then the equation above holds because both sides equal ∞ . Hence we can consider only the case where $\mu(E_k) < \infty \forall k \in \mathbb{N}$.

For convenience, let $E_0 = \emptyset$. Then,

$$\bigcup_{k=1}^{\infty} E_k = \bigcup_{j=1}^{\infty} (E_j \setminus E_{j-1})$$

Where union on the right side is a disjoint union. Thus,

$$\begin{aligned}
\mu\left(\bigcup_{k=1}^{\infty} E_k\right) &= \sum_{j=1}^{\infty} \mu(E_j \setminus E_{j-1}) \\
&= \lim_{k \rightarrow \infty} \sum_{j=1}^k \mu(E_j \setminus E_{j-1}) \\
&= \lim_{k \rightarrow \infty} \sum_{j=1}^k (\mu(E_j) - \mu(E_{j-1})) \\
&= \lim_{k \rightarrow \infty} \mu(E_k).
\end{aligned}$$

Hence proved. \square

Theorem 6.4. Suppose (X, \mathcal{A}, μ) is a measure space and $E_1 \supset E_2 \supset \dots$ is a decreasing sequence of sets in \mathcal{A} , with $\mu(E_1) < \infty$. Then

$$\mu\left(\bigcap_{k=1}^{\infty} E_k\right) = \lim_{k \rightarrow \infty} \mu(E_k).$$

Proof. By De Morgan's law we have,

$$E_1 \setminus \bigcap_{k=1}^{\infty} E_k = \bigcup_{k=1}^{\infty} (E_1 \setminus E_k).$$


Now $E_1 \setminus E_1, E_1 \setminus E_2, E_1 \setminus E_3$ is an increasing sequence of sets in \mathcal{A} . Thus, by 6.3,

$$\begin{aligned}
\mu\left(E_1 \setminus \bigcap_{k=1}^{\infty} E_k\right) &= \lim_{k \rightarrow \infty} \mu(E_1 \setminus E_k) \\
\mu\left(E_1 \setminus \bigcap_{k=1}^{\infty} E_k\right) &= \lim_{k \rightarrow \infty} (\mu(E_1) - \mu(E_k)) \\
\mu(E_1) - \mu\left(\bigcap_{k=1}^{\infty} E_k\right) &= \mu(E_1) - \lim_{k \rightarrow \infty} \mu(E_k) \\
\mu\left(\bigcap_{k=1}^{\infty} E_k\right) &= \lim_{k \rightarrow \infty} \mu(E_k)
\end{aligned}$$

\square

Theorem 6.5 (measure of union). Suppose (X, \mathcal{A}, μ) is a measure space and $D, E \in \mathcal{A}$, with $\mu(D \cap E) < \infty$. Then

$$\mu(D \cup E) = \mu(D) + \mu(E) - \mu(D \cap E).$$



Proof. We have

$$D \cup E = (D \setminus (D \cap E)) \cup (E \setminus (D \cap E)) \cup (D \cap E)$$

where every sets in right side are disjoint so from properties of measure,

$$\begin{aligned} \mu(D \cup E) &= \mu(D \setminus (D \cap E)) + \mu(E \setminus (D \cap E)) + \mu(D \cap E) \\ &= \mu(D) - \mu(D \cap E) + \mu(E) - \mu(D \cap E) + \mu(D \cap E) \\ &= \mu(D) + \mu(E) - \mu(D \cap E). \end{aligned}$$

Hence proved. □

7 Lebesgue Measure

While outer measure has the advantage that it is defined for all sets but we see in 1.2 that it is not additive. However outer Measure become measure if we suitably reduce the family of sets on which it is defined.

Definition 7.1 (Lebesgue Measurable set). A set E is said to be *Lebesgue measurable set* or simply *measurable set* if for each set A we have

$$\mu^*(A) = \mu^*(A \setminus E^c) + \mu^*(A \cap E)$$

Note: Outer measure is countable subadditive so it always holds $\mu^*(A) \leq \mu^*(A \setminus E^c) + \mu^*(A \cap E)$ so to proving E is measurable showing $\mu^*(A) \geq \mu^*(A \setminus E^c) + \mu^*(A \cap E)$ is enough.

Note: From definition we see that if E is measurable then E^c is also measurable. Clearly \emptyset is a Lebesgue measurable set then \mathbb{R} is also measurable.

Theorem 7.1. If $\mu^*(E) = 0$, then E is Lebesgue measurable set.

Proof. Let A is any set. Then $A \cap E \subset E$, So,

$$\mu^*(A \cap E) \leq \mu^*(E) = 0$$

But $\mu^*(A \cap E) \geq 0$ so, $\mu^*(A \cap E) = 0$.

Also $A \subset A \cap E^c$, then,

$$\mu^*(A) \geq \mu^*(A \cap E^c) = \mu^*(A \cap E^c) + \mu^*(A \cap E)$$

Hence E is Lebesgue measurable set. □

Theorem 7.2. If E_1 and E_2 are Lebesgue measurable set. Then $E_1 \cup E_2$ is Lebesgue measurable.

Proof. Let A be any set. Since E_1 is measurable,

$$\mu^*(A) = \mu^*(A \cap E_1) + \mu^*(A \cap E_1^c).$$

Again E_2 is measurable then,

$$\mu^*(A \cap E_1^c) = \mu^*((A \cap E_1^c) \cap E_2) + \mu^*((A \cap E_1^c) \cap E_2^c)$$

Now, $(A \cap E_1^c) \cap E_2^c = A \cap (E_1^c \cap E_2^c) = A \cap (E_1 \cup E_2)^c$.

Now,

$$\mu^*(A) = \mu^*(A \cap E_1) + \mu^*(A \cap E_1^c \cap E_2) + \mu^*(A \cap (E_1 \cup E_2)^c)$$

We see, $A \cap (E_1 \cup E_2) = (A \cap E_1) \cup (A \cap E_2 \cap E_1^c)$

So by property of outer measure, $\mu^*(A \cap (E_1 \cup E_2)) \leq \mu^*(A \cap E_1) + \mu^*(A \cap E_2 \cap E_1^c)$

Then,

$$\mu^*(A) \geq \mu^*(A \cap (E_1 \cup E_2)) + \mu^*(A \cap (E_1 \cup E_2)^c)$$

Hence, $E_1 \cup E_2$ is also Lebesgue measurable. □

Theorem 7.3. *The collection \mathcal{L} of Lebesgue Measurable sets is a σ -algebra.*

Proof. Let A be any set.

Now, $A \cap \emptyset = \emptyset$ and $A \cap \emptyset^c = A$ So,

$$\mu^*(A) = \mu^*(A \cap \emptyset) + \mu^*(A \cap \emptyset^c).$$

Hence, $\emptyset \in \mathcal{L}$ and $X \in \mathcal{L}$ (say, $\emptyset^c = X$).

Now from definition we say if $E \in \mathcal{L}$ then $E^c \in \mathcal{L}$.

Hence, \mathcal{L} is closed under complementation.

Now from 7.2 we say if $E_1, E_2 \in \mathcal{L}$ then $E_1 \cup E_2 \in \mathcal{L}$.

Now from previous $E_1^c, E_2^c \in \mathcal{L}$. Then $(E_1^c \cup E_2^c) \in \mathcal{L}$.

Then, $(E_1 \cap E_2)^c \in \mathcal{L} \implies (E_1 \cap E_2) \in \mathcal{L}$.

Now using induction on 7.2 we get, $F_n = \cup_{i=1}^n E_k \in \mathcal{L}$ where each $E_i \in \mathcal{L}$.

(if $\{E_n\}$ is sequence of sets in \mathcal{L} . Then we can construct another sequence of disjoint sets

E'_n in \mathcal{L} . Where $E'_1 = E_1$, $E'_n = E_n \setminus \cup_{i=1}^{n-1} E_k$

and $\cup_{i=1}^n E_k = \cup_{i=1}^n E'_i$)

Without lose of generality we choose $\{E_n\}$ is sequence of disjoint set.

Now F_n is measurable Then,

$$\mu^*(A) = \mu^*(A \cap F_n) + \mu^*(A \cap F_n^c). \quad (7)$$

Let, for all A and for some $n \in \mathbb{Z}^+$

$$\mu^*(A \cap F_n) = \sum_{i=1}^n \mu^*(A \cap E_k).$$

Then,

$$\begin{aligned} \mu^*(A \cap F_{n+1}) &= \mu^*((A \cap F_{n+1}) \cap F_n) + \mu^*((A \cap F_{n+1}) \cap F_n^c) \quad (\because F_n \text{ is measurable.}) \\ &= \mu^*(A \cap F_n) + \mu^*(A \cap E_{n+1}) \quad (\because F_{n+1} \cap F_n^c = E_{n+1}). \\ &= \sum_{i=1}^n \mu^*(A \cap E_k) + \mu^*(A \cap E_{n+1}) \\ &= \sum_{i=1}^{n+1} \mu^*(A \cap E_k). \end{aligned}$$

Hence the proposition is true for $n + 1$. And it is true for $n = 1$.

Hence for all n

$$\mu^*(A \cap F_n) = \sum_{i=1}^n \mu^*(A \cap E_k). \quad (8)$$

Let, $E = \cup_{i=1}^n E_k$. Then clearly $F_n \subset E$ for all n .

Then,

$$\begin{aligned} \mu^*(A \cap E) &\geq \mu^*(A \cap F_n) \quad \text{By monotone property} \\ &= \sum_{i=1}^n \mu^*(A \cap E_k). \end{aligned}$$

making $n \rightarrow \infty$, We have,

$$\mu^*(A \cap E) \geq \sum_{i=1}^{\infty} \mu^*(A \cap E_k).$$

by countable subadditivity, the converse inequality holds. Then,

$$\mu^*(A \cap E) = \sum_{i=1}^{\infty} \mu^*(A \cap E_k) \quad (9)$$

From 7 and 8 we have.

$$\mu^*(A) = \sum_{i=1}^n \mu^*(A \cap E_k) + \mu^*(A \cap F_n^c) \quad (10)$$

now clearly $(A \cap F_n^c) \supset (A \cap E^c) \implies \mu^*(A \cap F_n^c) \geq \mu^*(A \cap E^c)$ Then,

$$\mu^*(A) \geq \sum_{i=1}^n \mu^*(A \cap E_k) + \mu^*(A \cap E^c) \quad (11)$$

Making $n \rightarrow \infty$ in 11 We get,

$$\begin{aligned} \mu^*(A) &\geq \sum_{i=1}^{\infty} \mu^*(A \cap E_k) + \mu^*(A \cap E^c) \\ &\geq \mu^*(A \cap E) + \mu^*(A \cap E^c) \text{ (By 9)} \end{aligned}$$

Hence $E = \cup_{i=1}^{\infty} E_k$ is measurable. In other word \mathcal{L} is closed under countable union.

Hence \mathcal{L} is a σ -algebra. \square

Hence outer measure become measure (i.e. additive) when it is restricted on σ -algebra.

Definition 7.2 (Lebesgue Measure). Let μ^* is outer measure on \mathbb{R} (set of all real number) and \mathcal{L} is a collection of Lebesgue measurable set. Then μ^* is *Lebesgue Measure* on the measure space $(\mathbb{R}, \mathcal{L}, \mu^*)$.

Since Lebesgue Measure is a measure so it follows all the property of measure.

Theorem 7.4. *The interval (a, ∞) is Lebesgue measurable.*

Proof. To prove (a, ∞) is measurable we show, for any set A ,

$$\begin{aligned} \mu^*(A) &= \mu^*(A \cap (a, \infty)) + \mu^*(A \cap (a, \infty)^c) \\ \text{i.e. } \mu^*(A) &\geq \mu^*(A \cap (a, \infty)) + \mu^*(A \cap (-\infty, a]). \end{aligned}$$

Let, $A_1 = A \cap (a, \infty)$ and $A_2 = A \cap (-\infty, a]$.

If $\mu^*(A) = \infty$ then we are done.

If $\mu^*(A) < \infty$, then given $\epsilon > 0$ there exist countable collection $\{I_n\}$ of open intervals which cover A and for which,

$$\sum_{k=1}^{\infty} \ell(I_k) \leq \mu^*(A) + \epsilon.$$

(Because, $\mu^*(A) = \inf \{\sum_{k=1}^{\infty} I_k\}$ where, $A \subset \cup_{k=1}^{\infty} I_n$)

Let, $I'_n = I_n \cap (a, \infty)$ and $I''_n = I_n \cap (-\infty, a)$.

Then I'_n and I''_n are interval(can be empty) and,

$$\begin{aligned}\ell(I_n) &= \ell(I'_n) + \ell(I''_n) \\ &= \mu^*(I'_n) + \mu^*(I''_n).\end{aligned}$$

Since, $A_1 \subset \cup_{i=1}^{\infty} I'_k$ we have,

$$\mu^*(A_1) \leq \mu^*\left(\bigcup_{i=1}^{\infty} I'_k\right) \leq \sum_{i=1}^{\infty} \mu^*(I'_i).$$

Since, $A_2 \subset \cup_{i=1}^{\infty} I''_k$ we have,

$$\mu^*(A_2) \leq \mu^*\left(\bigcup_{i=1}^{\infty} I''_k\right) \leq \sum_{i=1}^{\infty} \mu^*(I''_i).$$

Then,

$$\begin{aligned}\mu^*(A_1) + \mu^*(A_2) &\leq \sum_{i=1}^{\infty} \mu^*(I'_k) + \sum_{i=1}^{\infty} \mu^*(I''_i) \\ &= \sum_{i=1}^{\infty} (\mu^*(I'_n) + \mu^*(I''_n)) \\ &= \sum_{i=1}^{\infty} \ell(I_k) \\ &\leq \mu^*(A) + \epsilon.\end{aligned}$$

Since ϵ is a arbitrary positive real number so we have,

$$\mu^*(A) \geq \mu^*(A_1) + \mu^*(A_2).$$

Hence, (a, ∞) is a measurable set. □

Theorem 7.5 (Borel Set are measurable). *Every Borel set is Lebesgue measurable set.*

Proof. Let \mathcal{L} is a collection of Lebesgue measurable sets. Then by (7.3) \mathcal{L} is a σ -algebra and by (7.4) $(a, \infty) \in \mathcal{L}$ for all $a \in \mathbb{R}$

Then, $(a, \infty)^c = (-\infty, a] \in \mathcal{L}$.

Since,

$$(-\infty, b) = \bigcup_{n=1}^{\infty} \left(-\infty, b - \frac{1}{n}\right]$$

then, $(-\infty, b) \in \mathcal{L}$

Again, $(a, b) = (\infty, b) \cap (a, \infty)$. Then, $(a, b) \in \mathcal{L}$.

Now, every open set is union of countable number of open intervals. So, every open set must belong to \mathcal{L} .

Therefor, \mathcal{L} is a σ -algebra containing every open set.

Therefor, \mathcal{L} contain the family \mathcal{B} of Borel Set. As \mathcal{B} is the smallest σ -algebra containing

every open set.

Hence, Borel sets are Lebesgue measurable. \square

Theorem 7.6 (Alternative Definition of Lebesgue Measure). *For any set $E \subset \mathbb{R}$ the followings are equivalent.*

1. E is measurable.
2. Given $\epsilon > 0$, there is an open set $O \supset E$ with $\mu^*(O \setminus E) < \epsilon$.
3. Given $\epsilon > 0$, there is a closed set $F \subset E$ with $\mu^*(E \setminus F) < \epsilon$.
4. There exist close sets F_1, F_2, \dots contained in E such that $\mu^*(E \setminus \bigcup_{k=1}^{\infty} F_k) = 0$.
5. There exist a Borel set $B \subset E$ such that $\mu^*(E \setminus B) = 0$.
6. There exist open sets G_1, G_2, \dots containing E such that $\mu^*(\bigcap_{k=1}^{\infty} G_k \setminus E) = 0$.
7. There exist a Borel set $B \supset E$ such that $\mu^*(B \setminus E) = 0$.

Proof. (1) \implies (2)

if $\mu^*(E) = \infty$ then the result is trivial. Let $\mu^*(E) < \infty$, Then given $\epsilon > 0$ there exist countable collection $\{I_n\}$ of open intervals such that,

$$\sum \ell(I_n) < \mu^*(E) + \epsilon.$$

Now we know every open set is union of countable open intervals, Then there exist a open set O such that.

$$\begin{aligned} \mu^*(O) &= \sum \ell(I_n) \\ \text{i.e. } \mu^*(O) &< \mu^*(E) + \epsilon \\ \text{i.e. } \mu^*(O) - \mu^*(E) &< \epsilon \\ \text{i.e. } \mu^*(O \setminus E) &< \epsilon \quad (\because O, E \text{ both are measurable}) \end{aligned}$$

(2) \implies (3)

Suppose (2) holds. Let $F \subset E$ where F is close set.

Then, $E^c \subset F^c$ here F^c is open set.

Then,

$$\begin{aligned} \mu^*(F^c \setminus E^c) &< \epsilon \\ \mu^*(E \setminus F) &< \epsilon. \quad (\because (F^c \setminus E^c) = (E \setminus F)). \end{aligned}$$

(3) \implies (4)

Suppose (3) holds. Thus for each $n \in \mathbb{Z}^+$, there exist a close set $F_n \subset E$ such that $\mu^*(E \setminus F_n) < \frac{1}{n}$. Now,

$$E \setminus \bigcup_{k=1}^{\infty} F_k \subset E \setminus F_n.$$

Thus, $\mu^*(E \setminus \bigcup_{k=1}^{\infty} F_k) \leq \mu^*(E \setminus F_n) \leq \frac{1}{n}$ for each $n \in \mathbb{Z}^+$.

Hence, $\mu^*(E \setminus \bigcup_{k=1}^{\infty} F_k) = 0$, completing the proof.

(4) \implies (5)

Because every countable union of closed sets is Borel set, Hence the result.

(2) \implies (6)

Suppose (2) holds.

Thus for each $n \in \mathbb{Z}^+$, there exist an open set $G_n \supset E$ such that $\mu^*(G_n \setminus E) < \frac{1}{n}$. Now,

$$\left(\bigcap_{k=1}^{\infty} G_k \right) \setminus A \subset G_n \setminus A.$$

Thus $\mu^* \left(\left(\bigcap_{k=1}^{\infty} G_k \right) \setminus A \right) \leq \mu^*(G_n \setminus A) \leq \frac{1}{n}$ for each $n \in \mathbb{Z}^+$.

Hence $\mu^*(G_n \setminus A) = 0$, completing the proof. (6) \implies (7)

Because every countable intersection of open set is a borel set. Hence the result is proved.

(5) \implies (1)

Suppose (5) holds. Thus there exist a borel set $B \subset E$ such that $\mu^*(E \setminus B) = 0$. Then, $E \setminus B \in \mathcal{L}$ by 7.1. Now,

$$E = B \cup (E \setminus B).$$

Since $B \in \mathcal{L}$, $(E \setminus B) \in \mathcal{L}$ and \mathcal{L} is σ -algebra, $E \in \mathcal{L}$.

Hence E is measurable.

(7) \implies (1)

Suppose (7) Holds. Thus there exist a borel set $B \supset E$ such that $\mu^*(B \setminus E) = 0$ Now, $B \setminus E$ measurable implies $R \setminus (B \setminus E)$ is also measurable and B is measurable (AS Borel Sets are measurable)

$$E = B \cap (R \setminus (B \setminus E)).$$

Thus, E is also measurable. □

8 Cantor Set

Every countable set has outer measure 0. So the natural question is the converse holds. In other word, is every set with outer measure 0 countable?

To construct a cantor set we start with interval $[0, 1]$ and removing the middle-third open interval at each step of remaining intervals from previous step.

At first step we remove $G_1 = (\frac{1}{3}, \frac{2}{3})$ then we have $[0, 1] \setminus G_1 = [0, \frac{1}{3}] \cup [\frac{2}{3}, 1]$. In the second step we remove $G_1 \cup G_2 = (\frac{1}{3}, \frac{2}{3}) \cup (\frac{1}{9}, \frac{2}{9}) \cup (\frac{7}{9}, \frac{8}{9})$. Then we left with

$$[0, 1] \setminus (G_1 \cup G_2) = \left[0, \frac{1}{9}\right] \cup \left[\frac{2}{9}, \frac{1}{3}\right] \cup \left[\frac{2}{3}, \frac{7}{9}\right] \cup \left[\frac{8}{9}, 1\right]$$

repeating this presses we construct cantor set.

Definition 8.1 (Cantor Set). The cantor set C is $[0, 1] \setminus (\bigcup_{k=1}^{\infty} G_k)$, where $G_1 = (\frac{1}{3}, \frac{2}{3})$ and G_n for $n > 1$ is the union of the middle-third open intervals in the interval of $[0, 1] \setminus (\bigcup_{k=1}^{n-1} G_k)$.




Figure 1: Cantor Set

Theorem 8.1. *The Cantor set is nonempty.*

Proof. Each trisection G_n to form G_{n+1} leaves exactly two endpoints. For example removing $(\frac{1}{3}, \frac{2}{3})$ from $[0, 1]$ leaves the points $p_0 = \frac{1}{3}$ and $p_1 = \frac{2}{3}$. In fact, since the Cantor set is the infinite intersection of each G_n , C contain the endpoints of each such subinterval, and is clearly non-empty. In fact, it is infinite. \square

Theorem 8.2.

1. *The Cantor set is a closed subset of \mathbb{R} .*
2. *The Cantor set has Lebesgue measure 0.*
3. *The Cantor contains no interval with more than one element.*

Proof. Each set G_n used in the definition of cantor set is a union of open intervals. Thus G_n is open. Thus $\bigcup_{n=1}^{\infty} G_n$ is open set, and hence its complement is closed. The Cantor set equals $[0, 1] \cap (\mathbb{R} \setminus \bigcup_{n=1}^{\infty} G_n)$, which is the intersection of two closed sets. Thus the Cantor set is a closed, completing the proof of (1).

By induction on n , each G_n is the union of 2^{n-1} disjoint open interval, each of which length $\frac{1}{3^n}$. Thus $\mu^*(G_n) = \frac{2^{n-1}}{3^n}$. The set G_1, G_2, \dots are disjoint. Hence,

$$\begin{aligned}\mu^*\left(\bigcup_{n=1}^{\infty} G_n\right) &= \frac{1}{3} + \frac{2}{9} + \frac{4}{27} + \dots \\ &= \frac{1}{3} \left(1 + \frac{2}{3} + \frac{4}{9} + \dots\right) \\ &= \frac{1}{3} \cdot \frac{1}{1 - \frac{2}{3}} \\ &= 1.\end{aligned}$$

Thus the Cantor set, which is equal to $[0, 1] \setminus \bigcup_{n=1}^{\infty} G_n$, has Lebesgue measure $1 - 1$. In other word the Cantor has Lebesgue measure 0. This proves (2).

A set with Lebesgue measure 0 cannot contain an interval that has more than one element. Thus (2) \implies (3). \square

Now first element of Cantor set is 0. If it countable then we can tell its second elements but we cann't and it contain infinitely many point. Hence it is uncountable. In fact C contain as many element as \mathbb{R} . Although canto set is has measure 0.

9 Vitali Set

We see in 1.2 That every subset of \mathbb{R} is not measurable. So, There exist such subset of \mathbb{R} which is not measurable. Vitali set is such nonmeasurable set.

We define a relation on $(0, 1]$ by $x \sim y$ iff $x - y \in \mathbb{Q}$. Then,

1. $x \sim x$ as $x - x = 0 \in \mathbb{Q}$.
2. if $x \sim y$ then $y \sim x$ as $x - y = y - x \in \mathbb{Q}$.
3. if $x \sim y$ and $y \sim z$.

Then, $x - y \in \mathbb{Q}$ and $y - z \in \mathbb{Q}$

then, $(x - y) + (y - z) = x - z \in \mathbb{Q}$ i.e. $x \sim z$.

Hence the above relation is equivalence. Choosing a single point from each equivalence class we construct a set V . This construction is possible because of *axiom of choice* even though it is very controversial topic in mathematics. Without *axiom of choice* it impossible to construct a non measurable set.

Definition 9.1 (Vitali Set). A set $V \subset (0, 1]$ is called Vitali set if V contain a single point of $(x + \mathbb{Q}) \cap (0, 1]$ for each $x + \mathbb{Q} \in \mathbb{R}/\mathbb{Q}$.

Theorem 9.1. *Vitali set is not Lebesgue measurable.*

Proof. Let r_1, r_2, \dots be the sequence of all rational number in $[0, 1]$. To prove this theory we first proof.

1. if $i \neq j$ then $V + r_i \cap V + r_j = \emptyset$.
2. $(0, 1] = \bigcup_{i=1}^{\infty} V + r_i$.

Where $V + r_i$ is define by $\{a + x : a \in V\}$ and define

$$x + y = \begin{cases} x + y, & x + y \leq 1 \\ x + y - 1, & \text{Otherwise} \end{cases}$$

Proof(1): Suppose $x \in V + r_i \cap V + r_j$.

Then, $x = v_i + r_i = v_j + r_j$ for some $v_i, v_j \in V$

This implies that $v_i - v_j \in \mathbb{Q}$. So, v_i and v_j belong to same class.

Since V contain exactly one element of each class. Hence $s_i = s_j$.

Since $0 < r_i, r_j \leq 1$ we would also have $r_i = r_j \implies i = j$, completing the prove of (1).

Proof(2): If $x \in (0, 1]$ then, $x \sim v$ for some $v \in V$.

Since x must be in some equivalence class and representative of each class appears in V .

Then x differs from y by some rational number in $(0, 1]$ so $x \in V + r_i$ for some i .

Hence $(0, 1] = \bigcup_{i=1}^{\infty} V + r_i$ proving (2).

Then we get each $V + r_i$ are disjoint and $(0, 1] = \bigcup_{i=1}^{\infty} V + r_i$.

let V is measurable and $\mu^*(V) = a$. Then $\mu^*(V + r_i) = a$ as measure is translation invariant.

Then we have,

$$\begin{aligned} 1 = \mu^*((0, 1]) &= \mu^*\left(\bigcup_{i=1}^{\infty} V + r_i\right) \\ &= \sum_{i=1}^{\infty} \mu^*(V + r_i) \\ &= a + a + a + \dots \end{aligned}$$

Now write side is either 0 or ∞ . A contradiction.

Then V is non measurable set.

□

References

- [1] MTH 535 Fall 2004. A non-measurable set, 2004. <https://www.math.kth.se/matstat/gru/godis/nonmeas.pdf>.
- [2] Sheldon Axler. *Measure, integration & real analysis*. Springer Nature, 2020.
- [3] University of California DYLAN R. NELSON. The cantor set - a brief introduction. <https://www.math.uwaterloo.ca/~xzliu/cantor-set.pdf>.
- [4] H. L. Royden. *Real analysis*. Macmillan New York, 3 edition, 1988.
- [5] Walter Rudin. *Real and Complex Analysis*. McGraw-hill New York, 1974.
- [6] Walter Rudin et al. *Principles of mathematical analysis*, volume 3. McGraw-hill New York, 1976.