

Markov Chain

Azmain Biswas

April 2023

Contents

1	Introduction	2
2	Markov Chain	3
2.1	Definition	3
2.2	Homogeneous Markov Chain	3
2.3	Chapman-Kolmogorov Equations	4
2.4	Classification of states	7

Chapter 1

Introduction

Chapter 2

Markov Chain

2.1 Definition

Definition 2.1.1 (Markov Chain). A discrete time stochastic process $\{X_n, n = 1, 2, 3, \dots\}$ is defined to be *Discrete Time Markov Chain* or simply *Markov Chain* if it takes value the state space \mathbf{S} , and for every $n \geq 0$ it satisfy the property

$$\mathbf{P}(X_{n+1} = j | X_n = i, X_{n-1} = i_{n-1}, \dots, X_0 = i_0) = \mathbf{P}(X_{n+1} = j | X_n = i) \quad (2.1)$$

Unless otherwise mentioned we take the state space \mathbf{S} to be $\{0, 1, 2, 3, \dots\}$. If $X_n = i$ we say that the process is in i th state at time n . In the definition eq. (2.1) may be interpreted as for Markov Chain, the conditional distribution of any future state X_{n+1} , given the past states X_0, X_1, \dots, X_{n-1} and the present state X_n , is independent of the past and only depend on the present state. This property is called *Markovian Property*. In other word for markov chain predicting the future we only need information about the present state.

Note: Notice that Assumption we are making for markov chain to forget the past as long as present in known is very strong assumption.

2.2 Homogeneous Markov Chain

Definition 2.2.1 (Homogeneous Markov Chain). We say a markov chain $\{X_n, n \geq 0\}$ is homogeneous if $\mathbf{P}(X_{n+1} = j | X_n = i) = \mathbf{P}(X_2 = j | X_1 = i) \forall n > 0$.

The quantity $\mathbf{P}(X_{n+1} = j | X_n = i)$ is called the *transition probability* from state i to state j . For homogeneous Markov Chain we can specify the transition probabilities $\mathbf{P}(X_{n+1} = j | X_n = i)$ by a sequence of value $p_{xy} = \mathbf{P}(X_{n+1} = y | X_n = x)$.

For the case of finite state Markov chain, say the state space is $\{1, 2, 3, \dots\}$. Then the transition probabilities are p_{ij} , $1 \leq i, j \leq N$ for transition from state i to state j . The $N \times$

N matrix $P = (p_{ij})_{N \times N}$ is called The *Transition Matrix* of chain. Since probabilities are nonnegative and since the process must make a transition into some state, we have

$$P_{ij} \geq 0, \quad i, j \geq 0, \quad \text{and} \quad \sum_{j=0}^N P_{ij} = 1, \quad \forall i = 0, 1, 2, \dots$$

For, infinite state markov chain the probability transition matrix will be infinite order. Then,

$$P = \begin{bmatrix} p_{00} & p_{01} & p_{02} & \dots \\ p_{11} & p_{12} & p_{13} & \dots \\ \vdots & \vdots & \vdots & \dots \\ p_{i0} & \dots & p_{ij} & \dots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

Example 2.2.1 (Rain and sunny). Suppose that whether it rains today depends on previous weather conditions only from the last two days. Specifically, suppose that if it has rained for the past two days, then it will rain tomorrow with probability 0.7; if it rained today but not yesterday, then it will rain tomorrow with probability 0.5; if it rained yesterday but not today, then it will rain tomorrow with probability 0.4; if it has not rained in the past two days, then it will rain tomorrow with probability 0.2.

We can transform it into a Markov chain by letting the state on any day be determined by the weather conditions during both that day and the preceding one. For instance, we can say that the process is in

State 0: if it rained both today and yesterday

State 1: if it rained today but not yesterday

State 2: if it rained yesterday but not today

State 3: if it rained neither today nor yesterday

The it will be 4-state Markov chain whose transition matrix will be,

$$P = \begin{bmatrix} 0.7 & 0 & 0.3 & 0 \\ 0.5 & 0 & 0.5 & 0 \\ 0 & 0.4 & 0 & 0.6 \\ 0 & 0.2 & 0 & 0.8 \end{bmatrix}$$

2.3 Chapman-Kolmogorov Equations

We have already defined the one step transition probabilities p_{ij} . We now define the n -step transition probabilities p_{ij}^n to be the probability that a process in state i will be in state j after

n additional transitions. i.e.

$$p_{ij}^n = \mathbf{P}(X_{n+m} = j | X_m = i), \quad n \geq 0, \quad i, j \geq 0.$$

By definition of markov chain we get,

$$\begin{aligned} p_{ij}^{n+m} &= \mathbf{P}(X_{n+m} = j | X_0 = i) \\ &= \sum_{k=0}^{\infty} \mathbf{P}(X_{n+m} = j, X_n = k | X_0 = i) \quad (\text{By theorem of total probability}) \\ &= \sum_{k=0}^{\infty} \mathbf{P}(X_{n+m} = j | X_n = k, X_0 = i) \mathbf{P}(X_n = k | X_0 = i) \\ &= \sum_{k=0}^{\infty} \mathbf{P}(X_{n+m} = j | X_n = k) \mathbf{P}(X_n = k | X_0 = i) \\ &= \sum_{k=0}^{\infty} p_{kj}^m p_{ik}^n. \end{aligned} \tag{2.2}$$

If we take $n = m = 1$. Then

$$p_{ij}^2 = \sum_{k=0}^{\infty} p_{kj} p_{ik} \tag{2.3}$$

the above expression is (i, j) element of P^2 matrix then we see eq. (2.3) in matrix form,

$$P^2 = P * P$$

Hence, eq. (2.2) can also written in matrix form,

$$P^{n+m} = P^n * P^m$$

Where P^n & P^m are the n-step and m-step transition matrix respectively.

Example 2.3.1 (Transition matrix of 4-state Markov chain). Consider the 4-state Markov chain depicted in fig. 2.1 When no probabilities are written over the arrows, as in this case, it means all arrows originating from a given state are equally likely. For example, there are 3 arrows originating from state 1, so the transitions $1 \rightarrow 3$, $1 \rightarrow 2$, and $1 \rightarrow 1$ all have probability $1/3$. Therefore the transition matrix of the chain is.

$$P = \begin{bmatrix} \frac{1}{3} & \frac{1}{3} & \frac{1}{3} & 0 \\ 0 & 0 & \frac{1}{2} & \frac{1}{2} \\ 0 & 1 & 0 & 0 \\ \frac{1}{2} & 0 & 0 & \frac{1}{2} \end{bmatrix}$$




Figure 2.1: Example of A 4 state markov chain

To compute the probability that the chain is in state 3 after 5 steps, starting at state 1, we would look at the (1,3) entry of P^5 .

$$P^5 = \begin{bmatrix} \frac{853}{3888} & \frac{509}{1944} & \frac{52}{243} & \frac{395}{1296} \\ \frac{173}{864} & \frac{85}{432} & \frac{31}{108} & \frac{91}{288} \\ \frac{37}{144} & \frac{29}{72} & \frac{1}{9} & \frac{11}{48} \\ \frac{499}{2592} & \frac{395}{1296} & \frac{71}{324} & \frac{245}{864} \end{bmatrix}$$

so, $p_{13}^5 = \mathbf{P}(X_5 = 3 | X_0 = 1) = \frac{52}{243}$.

To get the marginal distributions of X_0, X_1, \dots , we need to specify not just the transition matrix, but also the initial conditions of the chain. This can be done by setting the initial state X_0 to be a particular state, or by randomly choosing X_0 according to some distribution. Let (t_1, t_2, \dots, t_N) be the PMF of X_0 displayed as a vector, that is, $t_i = \mathbf{P}(X_0 = i)$. Then the marginal distribution of the chain at any time can be computed from the transition matrix.

Proposition 2.3.1 (Marginal Distribution of X_n). *Define $\mathbf{t} = (t_1, t_2, \dots, t_n)$ by $t_i = \mathbf{P}(X_0 = i)$, and view \mathbf{t} as a row vector. Then the marginal distribution of X_n is given by the vector $\mathbf{t}P^n$. That is the j -th component of $\mathbf{t}P^n$ is $\mathbf{P}(X_n = j)$.*

Proof. By the law of total probability we get,

$$\begin{aligned}
\mathbf{P}(X_n = j) &= \sum_{i=0}^N \mathbf{P}(X_n, X_0 = i) \\
&= \sum_{i=0}^N \mathbf{P}(X_0 = i) \mathbf{P}(X_n = j | X_0 = i) \\
&= \sum_{i=0}^N t_i p_{ij}^n
\end{aligned}$$

Which is the j 0th component of $\mathbf{t}P^n$. □

Example 2.3.2 (Marginal distribution of 4-state Markov chain). Again consider the 4-state Markov chain in fig. 2.1. Suppose the initial conditions are $\mathbf{t} = (1/4, 1/4, 1/4, 1/4)$, Let X_n be the position of the chain at time n . Then distribution of X_1 is

$$\begin{aligned}
\mathbf{t}P &= \begin{bmatrix} \frac{1}{4} & \frac{1}{4} & \frac{1}{4} & \frac{1}{4} \end{bmatrix} \begin{bmatrix} \frac{1}{3} & \frac{1}{3} & \frac{1}{3} & 0 \\ 0 & 0 & \frac{1}{2} & \frac{1}{2} \\ 0 & 1 & 0 & 0 \\ \frac{1}{2} & 0 & 0 & \frac{1}{2} \end{bmatrix} \\
&= \begin{bmatrix} \frac{5}{24} & \frac{1}{3} & \frac{5}{24} & \frac{1}{4} \end{bmatrix}
\end{aligned}$$

The marginal distribution of X_5 is

$$\begin{aligned}
\mathbf{t}P^5 &= \begin{bmatrix} \frac{1}{4} & \frac{1}{4} & \frac{1}{4} & \frac{1}{4} \end{bmatrix} \begin{bmatrix} \frac{853}{3888} & \frac{509}{1944} & \frac{52}{243} & \frac{395}{1296} \\ \frac{173}{864} & \frac{85}{432} & \frac{31}{108} & \frac{91}{288} \\ \frac{37}{144} & \frac{29}{72} & \frac{1}{9} & \frac{11}{48} \\ \frac{499}{2592} & \frac{395}{1296} & \frac{71}{324} & \frac{245}{864} \end{bmatrix} \\
&= \begin{bmatrix} \frac{707}{3254} & \frac{472}{1619} & \frac{101}{486} & \frac{1469}{5184} \end{bmatrix}
\end{aligned}$$

2.4 Classification of states