

$$1. \textcircled{a} \quad A = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 4 & 16 \\ 1 & -1 & 1 \\ 1 & 1 & 1 \end{bmatrix}, x = \begin{bmatrix} a_0 \\ a_1 \\ a_2 \end{bmatrix}, b = \begin{bmatrix} 3 \\ -2 \\ 2 \\ 1 \end{bmatrix}$$

$$\begin{aligned} \textcircled{b} \quad A^T A &= \begin{bmatrix} 1 & 0 & 0 \\ 1 & 4 & 16 \\ 1 & -1 & 1 \\ 1 & 1 & 1 \end{bmatrix}^T \begin{bmatrix} 1 & 0 & 0 \\ 1 & 4 & 16 \\ 1 & -1 & 1 \\ 1 & 1 & 1 \end{bmatrix} \\ &= \begin{bmatrix} 1 & 1 & 1 & 1 \\ 0 & 4 & -1 & 1 \\ 0 & 16 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 1 & 4 & 16 \\ 1 & -1 & 1 \\ 1 & 1 & 1 \end{bmatrix} \\ &= \begin{bmatrix} 4 & 4 & 18 \\ 4 & 18 & 64 \\ 18 & 64 & 258 \end{bmatrix} \end{aligned}$$

$$\det(A^T A) = 1448$$

© Using least square approximation

$$(A^T A)x = A^T b$$

$$\begin{bmatrix} 4 & 4 & 18 \\ 4 & 18 & 64 \\ 18 & 64 & 258 \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \\ a_2 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 4 & -1 \\ 0 & 16 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ -2 \\ 2 \end{bmatrix}$$

$$\begin{bmatrix} 4 & 4 & 18 \\ 4 & 18 & 64 \\ 18 & 64 & 258 \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \\ a_2 \end{bmatrix} = \begin{bmatrix} 4 \\ -9 \\ -29 \end{bmatrix}$$

$$\begin{bmatrix} a_0 \\ a_1 \\ a_2 \end{bmatrix} = \begin{bmatrix} 4 & 4 & 18 \\ 4 & 18 & 64 \\ 18 & 64 & 258 \end{bmatrix}^{-1} \begin{bmatrix} 4 \\ -9 \\ -29 \end{bmatrix}$$

$$\begin{bmatrix} a_0 \\ a_1 \\ a_2 \end{bmatrix} = \begin{bmatrix} 2.129 \\ -0.383 \\ -0.166 \end{bmatrix}$$

$$P_2(x) = 2.129 - 0.383x - 0.166x^2$$

3. Consider the coordinates: $(x, f(x)) = (0, 1), (0.5, 1.4), (1, 1.7), (1.5, 2)$. In the following, you are asked to construct the **best-fit linear polynomial** by using the **QR-decomposition** method as follows:

- (2 marks) Construct the matrices A , b and x .
- (3 marks) Evaluate the orthonormal vectors q_1 and q_2 , and construct the matrix Q .
- (2 marks) Compute the matrix R .
- (3 marks) Using Q and R , evaluate the matrix x , and hence compute the best-fit linear polynomial.

$$\textcircled{3} \quad \textcircled{a)} \quad A = \begin{bmatrix} 1 & 0 \\ 1 & 0.5 \\ 1 & 1 \\ 1 & 1.5 \end{bmatrix} \quad x = \begin{bmatrix} a_0 \\ a_1 \end{bmatrix} \quad b = \begin{bmatrix} 1 \\ 1.4 \\ 1.7 \\ 2 \end{bmatrix}$$

$$\textcircled{b)} \quad \text{Hence, } u_1 = (1, 1, 1, 1)^T, \quad u_2 = (0, 0.5, 1, 1.5)^T$$

$$p_1 = u_1 = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}$$

$$q_1 = \frac{p_1}{|p_1|} = \frac{(1, 1, 1, 1)^T}{\sqrt{4}} = \frac{1}{2} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}$$

$$\begin{aligned} p_2 &= u_2 - (u_2^T q_1) q_1 \\ &= u_2 - \left([0 \ 0.5 \ 1 \ 1.5] \times \begin{bmatrix} 0.5 \\ 0.5 \\ 0.5 \\ 0.5 \end{bmatrix} \right) \times \begin{bmatrix} 0.5 \\ 0.5 \\ 0.5 \\ 0.5 \end{bmatrix} \end{aligned}$$

$$= u_2 - 1.5 \times \begin{bmatrix} 0.5 \\ 0.5 \\ 0.5 \\ 0.5 \end{bmatrix}$$

$$= \begin{bmatrix} 0 \\ 0.5 \\ 1 \\ 1.5 \end{bmatrix} - \begin{bmatrix} 0.75 \\ 0.75 \\ 0.75 \\ 0.75 \end{bmatrix} = \begin{bmatrix} -0.75 \\ -0.25 \\ 0.25 \\ 0.75 \end{bmatrix}$$

$$q_2 = \frac{p_2}{|p_2|} = \frac{2}{\sqrt{5}} \begin{bmatrix} -0.75 \\ -0.25 \\ 0.25 \\ 0.75 \end{bmatrix} = \begin{bmatrix} -0.67 \\ -0.22 \\ 0.22 \\ 0.67 \end{bmatrix}$$

$$Q = [q_1 \ q_2] = \begin{bmatrix} 0.5 & -0.67 \\ 0.5 & -0.22 \\ 0.5 & 0.22 \\ 0.5 & 0.67 \end{bmatrix}$$

$$\textcircled{c} \quad R = Q^T A = \begin{bmatrix} 0.5 & -0.67 \\ 0.5 & -0.22 \\ 0.5 & 0.22 \\ 0.5 & 0.67 \end{bmatrix}^T \begin{bmatrix} 1 & 0 \\ 1 & 0.5 \\ 1 & 1 \\ 1 & 1.5 \end{bmatrix}$$

$$= \begin{bmatrix} 2 & 1.5 \\ 0 & 1.115 \end{bmatrix}$$

$$\textcircled{d} \quad Rx = Q^T b$$

$$\Rightarrow \begin{bmatrix} 2 & 1.5 \\ 0 & 1.115 \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \end{bmatrix} = \begin{bmatrix} 0.5 & -0.67 \\ 0.5 & -0.22 \\ 0.5 & 0.22 \\ 0.5 & 0.67 \end{bmatrix}^T \begin{bmatrix} 1 \\ 1.4 \\ 1.7 \\ 2 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 2 & 1.5 \\ 0 & 1.115 \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \end{bmatrix} = \begin{bmatrix} 3.05 \\ 0.736 \end{bmatrix}$$

$$\therefore a_1 = \frac{0.736}{1.115} = \frac{736}{1115} = 0.66$$

$$\therefore 2a_0 + 1.5a_1 = 3.05 \Rightarrow a_0 = \frac{3.05 - 1.5a_1}{2} = 1.03$$

$$P_1(x) = a_0 + a_1 x = 1.03 + 0.66x$$

4. A function is given by $f(x) = e^{0.5x} + \sin x$ which is to be integrated on the interval $[0, 2]$.

a. (2 marks) Evaluate the exact integral $I(f)$.

b. (2 marks) Compute the numerical integral by using the Newton-Cotes formula with $n = 2$.

c. (4 marks) Evaluate the numerical integral $C_{1,4}$ by using the Composite Newton-Cotes formula and also find the percentage relative error.

④

①

$$I(f) = \int_a^b f(x) dx$$

$$= \int_0^2 (e^{0.5x} + \sin x) dx$$

$$= 4.853$$

$$\begin{aligned}
 \textcircled{b} \quad I_2(f) &= \frac{b-a}{6} [f(a) + 4f(m) + f(b)] \\
 &= \frac{2-0}{6} [f(0) + 4f(1) + f(2)] \\
 &= \frac{1}{3} [14.588] \\
 &= 4.863
 \end{aligned}$$

$$\begin{aligned}
 a &= 0 \\
 b &= 2 \\
 m &= \frac{a+b}{2} = 1
 \end{aligned}$$

$$\textcircled{c} \quad \text{Here, } h = \frac{b-a}{n} = \frac{2-0}{4} = \frac{1}{2}$$

$$\begin{aligned}
 \text{Now, } x_0 &= 0 \\
 x_1 &= 0 + \frac{1}{2} = \frac{1}{2} \\
 x_2 &= \frac{1}{2} + \frac{1}{2} = 1 \\
 x_3 &= 1 + \frac{1}{2} = \frac{3}{2} \\
 x_4 &= \frac{3}{2} + \frac{1}{2} = 2
 \end{aligned}$$

$$\begin{aligned}
 C_{1,1}(f) &= \frac{h}{2} [f(x_0) + 2f(x_1) + 2f(x_2) + 2f(x_3) + f(x_4)] \\
 &= \frac{1}{4} [f(0) + 2f(\frac{1}{2}) + 2f(1) + 2f(\frac{3}{2}) + f(2)] \\
 &= \frac{1}{4} [19.364] = 4.841
 \end{aligned}$$

$$\begin{aligned}
 \% \text{ relative error} &= \left| \frac{4.853 - 4.841}{4.841} \right| \times 100\% \\
 &= 0.248\%
 \end{aligned}$$