## Extension of Study TKT024 Evaluating Long-Term Safety and Clinical Outcomes in MPS II Patients Receiving Idursulfase

St. Joseph's Hospital, Phoenix, Arizona, United States

Pediatric Clinical Research Center, Children's Hospital Oakland, Oakland, California, United States

The Children's Hospital, Denver, Colorado, United States

Harbin Clinic, Rome, Georgia, United States

Mid-Illinois Hematology and Oncology Associates, Normal, Illinois, United States

University of Iowa Hospitals and Clinics, Iowa City, Iowa, United States

Children's Hospital Boston, Boston, Massachusetts, United States

Saint Louis University Cardinal Glennon Children's Hospital, St. Louis, Missouri, United States

University of Nebraska Medical Center, Omaha, Nebraska, United States

Comprehensive Cancer Centers of Nevada, Las Vegas, Nevada, United States

Upstate Medical University, State University of New York (SUNY), Syracuse, New York, United States

University of North Carolina at Chapel Hill, Chapel Hill, North Carolina, United States

Children's Hospital of Philadelphia, Philadelphia, Pennsylvania, United States

Baylor College of Medicine Texas Children's Hospital, Houston, Texas, United States

University of Utah Hospital, Salt Lake City, Utah, United States

Franciscan Skemp Healthcare, La Crosse, Wisconsin, United States

Fundacao Universidade de Ciencias da Saude de Alagoas Governador Lamenha Filho / UNCISAL, Maceio, AL, Brazil

Clinica Casa de Saude Sao Joao, Barreiras, BA, Brazil

c-HUPES/UFBA, Salvador, BA, Brazil

Hospital Universitario da Faculdade de Medicina da Universidade Federal de Mato Grosso do Sul, Campo Grande, MS,

Instituto de Puericultura e Pediatria Martagao Gesteira / Hospital Pediatrico, Rio de Janeiro, RJ, Brazil

Hospital de Clinicas de Porto Alegre, Servico de Genetica Medica, Porto Alegre, RS, Brazil

UNIFESP Instituto de Oncologia Pediatrica, Sao Paulo, SP, Brazil

Instituto da Crianca / Hospital das Clinicas da Faculdade de Medicina da Universidade de Sao Paulo, Sao Paulo, SP,

The Hospital for Sick Children Research Institute, Toronto, Ontario, Canada

University of Montreal / Hopital Ste-Justine, Montreal, Quebec, Canada

Hopital Edouard Herriot, Lyon, , France

Hopital de Hautepierre, Strasbourg Cedex, , France

Hospital Ducuing, Toulouse Cedex, , France

Universitatsklinikum Aachen Kinderklinik, Aachen, , Germany

Universitatsklinik Dusseldorf Kinderklinik, Dusseldorf, , Germany

Justus-Liebig Universitat, Giessen, , Germany

Universitatsklinikum Gottingen, Goettingen, , Germany

Universitatsklinikum Hamburg Eppendorf, Hamburg, , Germany

Children's University Hospital Mainz AG, Mainz, , Germany

Universita Milano Bicocca / Ospedale S. Gerardo, Monza (Milano), , Italy

Universita degli Studi di Napoli Federico II, Napoli, , Italy

Universita di Padova, Padova, , Italy

Ospedale S. S. Annunziata, Savigliano (Cuneo), , Italy

Spitalul Clinic de Copii, Cluj Napoca, Cluj, Romania

University Hospital Germans Trias i Pujol, Badalona, , Spain

Servicio de Pediatria, Linares (Jaen), , Spain

Drottning Silvias Barnsjukhus, Gothenberg, , Sweden

Karolinska University Hospital, Stockholm, , Sweden

Bath and NE Somerset Primary Care Trust, Bath, , United Kingdom

Addenbrooke's Hospital, Cambridge, , United Kingdom

Derbyshire Children's Hospital, Derby, , United Kingdom

Royal Hospital for Sick Children, Glasgow, , United Kingdom

Great Ormond Street Hospital for Sick Children, London, , United Kingdom

Royal Manchester Children's Hospital, Manchester, , United Kingdom

Royal Victoria Infirmary, Newcastle, , United Kingdom

Royal Surrey County Hospital, Surrey, , United Kingdom

Study TKT024EXT was a long-term, single-arm, open-label extension of Study TKT024, a one year Phase 2/Phase 3 registration study. The primary objective of this extension study was to

collect long-term safety and clinical outcome data in Mucopolysaccharidosis II (MPS II), also

known as Hunter Syndrome, from the Phase 2/Phase 3 Study TKT024. All patients enrolling into this study received weekly active treatment with idursulface, the primary dosing regimen