

# **A Double-Blind Study to Evaluate the Efficacy and Safety of BMN 110 in Patients With Mucopolysaccharidosis IVA (Morquio A Syndrome)**

Oakland, California, United States  
Wilmington, Delaware, United States  
Washington, District of Columbia, United States  
Chicago, Illinois, United States  
New York, New York, United States  
Seattle, Washington, United States  
Cordoba, , Argentina  
Campina Grande, , Brazil  
Porto Alegre, , Brazil  
Montreal, , Canada  
Sherbrooke, , Canada  
Toronto, , Canada  
Bogota, , Colombia  
Copenhagen, , Denmark  
Lyon, , France  
Paris, , France  
Mainz, , Germany  
Monza, , Italy  
Tokyo, , Japan  
Seoul, , Korea, Republic of  
Amsterdam, , Netherlands  
Coimbra, , Portugal  
Doha, , Qatar  
Riyadh, , Saudi Arabia  
Taipei, , Taiwan  
Birmingham, , United Kingdom  
London, , United Kingdom  
Manchester, , United Kingdom

This Phase 3 study will evaluate the efficacy and safety of 2.0 mg/kg/week BMN 110 and 2.0 mg/kg/every other week BMN 110 in patients with mucopolysaccharidosis IVA (Morquio A Syndrome).

There is currently no standard accepted treatment for MPS IVA other than supportive care. Enzyme replacement therapy (ERT) may be a potential new treatment option for MPS IVA patients. BMN 110 is administered to MPS IVA patients by IV infusion, allowing cellular uptake by the mannose-6-phosphate receptor and transportation to the lysosomes.

This enzyme uptake into the lysosomes is hypothesized to promote increased catabolism of keratan sulfate (KS) in tissue macrophages, hyaline cartilage, other connective tissues, and heart valve, and reduce the progressive accumulation of KS which is responsible for the clinical manifestations of the disorders.