

Real Analysis

Azure

2021

Contents

1	Day 1: Ordered Sets & Fields	4
1.1	Ordered Sets and Bounds	4
1.2	Least Upper Bound Property	5
1.3	Fields	6
1.4	Ordered Fields	7
2	Day 2: Complex Field & Euclidean Spaces	9
2.1	n th Root	9
2.2	Decimals	10
2.3	Extended Reals	10
2.4	Complex Numbers	10
2.5	Euclidean Spaces	12
2.6	Cauchy-Schwarz	12
3	Day 3: Existence of \mathbb{R}	14
4	Day 4: Cardinality	16
4.1	Cardinality	16
4.2	Set of Sets	17
5	Day 5: Metric Spaces and Closed/Open	20
5.1	Metric Spaces	20
5.2	Intervals and Balls	24
6	Day 6: Open Relative and Compactness	25
6.1	Closure	25
6.2	Open Relative	25
6.3	Compact Sets	26
7	Day 7: Perfect & Connected Sets	30
7.1	Perfect Sets	30
7.2	Connected Sets	31
8	Day 8: Convergence & Cauchy	32
8.1	Convergent Sequences	32
8.2	Subsequences	34
8.3	Cauchy Sequences	35

9 Day 9: Limits & Special Sequences	38
9.1 Upper and Lower Limits	38
9.2 Special Sequences	39
10 Day 10: Series & Convergence Tests	40
10.1 Series	40
10.2 Series of Nonnegative Terms	41
10.3 The Number e	42
10.4 Root and Ratio Tests	43
10.5 Power Series	44
10.6 Summation By Parts	44
10.7 Absolute Convergence	45
10.8 Addition & Multiplication	45
11 Day 11: Continuity	47
11.1 Limits of Functions	47
11.2 Continuous Functions	48
11.3 Continuity and Compactness	50
11.4 Continuity and Connectedness	51
11.5 Discontinuities	52
11.6 Monotonic Functions	52
11.7 Infinite Limits \ Limits at Infinity	53
12 Day 12: Differentiation	54
12.1 Derivative of a function	54
12.2 Mean Value Theorems	55
12.3 Continuity of Derivatives	56
12.4 L'Hospital's Rule	57
12.5 Derivative of Higher Order	57
12.6 Taylor's Theorem	58
12.7 Differentiation of Vector-Valued Functions	58

1 Ordered Sets and Fields

1.1 Ordered Sets and Bounds

Definition 1.1.1: Ordered Set

An order is:

- **Trichotomy**: For all $x, y \in S$, only one holds true:
 - $x < y$
 - $x = y$
 - $x > y$
- **Transitivity**: If $x < y$ and $y < z$, then $x < z$.

An ordered set is a set with an order.

Definition 1.1.2: Bounds

Let S be an ordered set and $E \subset S$.

An upper bound of E is a $\beta \in S$ such that for $x \leq \beta$ for all $x \in E$.

If such a β exists, then E is bounded from above.

A lower bound of E is a $\alpha \in S$ such that for $x \geq \alpha$ for all $x \in E$.

If such a α exists, then E is bounded from below.

Definition 1.1.3: Infimum & Supremum

Let S be an ordered set.

Let $E \subset S$ be bounded from above. Least upper bound $\beta \in S$ exists if:

- β is an upper bound for E
- If $\gamma < \beta$, then γ is not an upper bound for E .

Then $\beta = \sup(E)$.

Let $E \subset S$ be bounded from below. Greatest lower bound $\alpha \in S$ exists if:

- α is a lower bound for E
- If $\gamma > \alpha$, then γ is not a lower bound for E .

Then $\alpha = \inf(E)$.

Even if $\sup(E)$ exists, it may or may not exist at S .

If $\sup(E)$ exists, then $\sup(E)$ is unique. Statement also holds true for $\inf(E)$.

Example

Let $S = (1, 2) \cup [3, 4) \cup (5, 6)$ with the order $<$ from \mathbb{R} . For subsets E of S :

- $E = (1, 2)$ is bounded above with $\sup(E) = 2$ and not bounded below.
- $E = (5, 6)$ is not bounded above or below so $\inf(E), \sup(E) = \text{DNE}$.
- $E = [3, 4)$ is bounded below with $\inf(E) = 3$, but $\sup(E) = \text{DNE}$.

1.2 Least Upper Bound Property

Theorem 1.2.1: Least Upper Bound Property

An ordered set S has a least upper bound property if:

For every nonempty subset $E \subset S$ that is bounded from above:

$\sup(E)$ exists in S .

Proof

Let $z = y - \frac{y^2-2}{y+2} = \frac{2y+2}{y+2}$, then take $z^2 - 2 = \frac{2(y^2-2)}{(y+2)^2}$.

Let set $A = \{y > 0 \in \mathbb{Q} \text{ where } y^2 < 2\}$ and set $B = \{y > 0 \in \mathbb{Q} \text{ where } y^2 > 2\}$

- If $y^2 - 2 < 0$, then $z > y$ where $z \in A$. So, y is not an upper bound.
Since for any y , there is $z > y$ where $z \in A$, then $\sup(A)$ doesn't exist in \mathbb{Q} .
- If $y^2 - 2 > 0$, then $z < y$ where $z \in B$. So, y is an upper bound, but not $\sup(E)$.
Since for any y , there is $z < y$ where $z \in B$, then $\inf(B)$ doesn't exist in \mathbb{Q} .

Thus, \mathbb{Q} doesn't have the least upper bound or greatest lower bound property.

Example

\mathbb{Q} doesn't have a least upper bound property. Take for example, $\sqrt{2}$.

Let $x^2 = 2$. If x was rational, there is a rational $\frac{p}{q}$ where $x = \frac{p}{q}$ where both p and q are not even.

$$\left(\frac{p}{q}\right)^2 = 2 \quad \Rightarrow \quad p^2 = 2q^2$$

Since $2q^2$ is even, then p^2 is even so p is even. Thus, p is divisible by 2 so p^2 is divisible by 4 so q^2 is divisible by 2 so q is even. Thus, both p and q must be even which is a contradiction so $x = \sqrt{2}$ cannot be rational.

So if $\sqrt{2} < \frac{a}{b}$ for some rational $\frac{a}{b}$, there is always another rational $\frac{p}{q}$:

$$\sqrt{2} < \frac{p}{q} < \frac{a}{b}$$

and there will never be a rational $\frac{p}{q}$ such that $\sqrt{2} = \frac{p}{q}$ since $\sqrt{2}$ is not rational.

Theorem 1.2.2: Least Upper Bound + Lower Bound implies Greatest Lower Bound

Let S be an ordered set with the least upper bound property.

Let non-empty $B \subset S$ be bounded below.

Let L be the set of all lower bounds of B .

Then $\alpha = \sup(L)$ exists in S .

Proof


L is non-empty since B is bounded from below.

Thus, by the least upper bound property of S , $\alpha = \sup(L)$ exists in S .

We claim that $\alpha = \inf(B)$.

If $\gamma < \alpha$, then γ is not an upper bound for L so $\gamma \notin B$ since all upper bounds for L are in B . Thus, for every $x \in B$, $\alpha \leq x$.

If $\gamma \geq \alpha$, then γ is an upper bound of L so $\gamma \in B$. Thus, $\inf(B) = \alpha$.



1.3 Fields

Definition 1.3.1: Fields Axioms

- (a) Addition Axioms
- If $x, y \in F$, then $x+y \in F$
 - $x+y = y+x$ for all $x, y \in F$
 - $(x+y)+z = x+(y+z)$ for all $x, y, z \in F$
 - There exists $0 \in F$ such that $0+x = x$ for all $x \in F$
 - For every $x \in F$, there is $-x \in F$ where $x+(-x) = 0$
- (b) Multiplicative Axioms
- If $x, y \in F$, then $xy \in F$
 - $yx = xy$ for all $x, y \in F$
 - $(xy)z = x(yz)$ for all $x, y, z \in F$
 - There exists $1 \neq 0 \in F$ such that $1x = x$ for all $x \in F$
 - If $x \neq 0 \in F$, there is $\frac{1}{x} \in F$ where $x(\frac{1}{x}) = 1$
- (c) Distributive Law
- $x(y+z) = xy + xz$ hold for all $x, y, z \in F$

Theorem 1.3.2: Consequences of the Field Axioms

- (a) If $x+y = x+z$, then $y = z$

Proof

$$y = 0+y = (-x)+x+y = (-x)+x+z = 0+z = z$$

- (b) If $x+y = x$, then $y = 0$

Proof

$$\text{From (a), let } z = 0$$

- (c) If $x+y = 0$, then $y = -x$

Proof

$$\text{From (a), let } z = -x$$

- (d) $-(-x) = x$

Proof

$$\text{From (c), let } x = -x \text{ and } y = x$$

- (e) If $x \neq 0$ and $xy = xz$, then $y = z$

Proof

$$y = 1y = \frac{1}{x}xy = \frac{1}{x}xz = 1z = z$$

- (f) If $x \neq 0$ and $xy = x$, then $y = 1$

Proof

$$\text{From (e), let } z = 1$$

- (g) If $x \neq 0$ and $xy = 1$, then $y = \frac{1}{x}$

Proof

$$\text{From (e), let } z = \frac{1}{x}$$

- (h) If
- $x \neq 0$
- , then
- $\frac{1}{1/x} = x$

Proof

From (g), let $x = \frac{1}{x}$ and $y = x$

- (i)
- $0x = 0$

Proof

Since $0x + 0x = (0+0)x = 0x = 0x + 0$, then $0x = 0$
--

- (j) If
- $x, y \neq 0$
- , then
- $xy \neq 0$

Proof

Suppose $xy = 0$, then $1 = \frac{1}{y} \frac{1}{x} xy = \frac{1}{y} \frac{1}{x} 0 = 0$. $0 = 1$ is a contradiction.

- (k)
- $(-x)y = -(xy) = x(-y)$

Proof

$xy + (-x)y = (x+(-x))y = 0y = 0$. Then by part (c), $(-x)y = -(xy)$. $xy + x(-y) = x(y+(-y)) = x0 = 0$. Then by part (c), $x(-y) = -(xy)$.
--

- (l)
- $(-x)(-y) = xy$

Proof

By part (k), then $(-x)(-y) = -[x(-y)] = -[-(xy)]$. By part (d), $-[-(xy)] = xy$.

1.4 Ordered Fields

Definition 1.4.1: Ordered Field

An ordered field F is a field F which is also an ordered set for all $x, y, z \in F$.

- If $y < z$, then $y+x < z+x$
- If $x, y > 0$, then $xy > 0$

\mathbb{Q}, \mathbb{R} are ordered fields, but \mathbb{C} is not an ordered field since $i^2 = -1 \not> 0$.

Theorem 1.4.2: Properties of the Ordered Field

- (a) If
- $x > 0$
- , then
- $-x < 0$
- and vice versa

Proof

$-x = -x + 0 < -x + x = 0$

- (b) If
- $x > 0$
- and
- $y < z$
- , then
- $xy < xz$

Proof

Since $z-y > 0$, then $0 < x(z-y) = xz - xy$

- (c) If
- $x < 0$
- and
- $y < z$
- , then
- $xy > xz$

Proof

Since $-x > 0$ and $z-y > 0$, then $0 < -x(z-y) = xy - xz$

- (d) If
- $x \neq 0$
- ,
- $x^2 > 0$

Proof

If $x > 0 \Rightarrow x^2 = x \cdot x > 0$. If $x < 0 \Rightarrow (-x)^2 = (-x) \cdot (-x) = x \cdot x = x^2 > 0$
--

- (e) If
- $0 < x < y$
- , then
- $0 < 1/y < 1/x$

Proof

$(\frac{1}{y})y = 1 > 0$ so $\frac{1}{y} > 0$. Since $x < y$, then $\frac{1}{y} = (\frac{1}{y})(\frac{1}{x})x < (\frac{1}{y})(\frac{1}{x})y = \frac{1}{x}$.
--

Theorem 1.4.3: \mathbb{R} is an ordered field

There exists a unique ordered field \mathbb{R} with the least upper bound property.

Also, $\mathbb{Q} \subset \mathbb{R}$ so \mathbb{Q} is also an ordered field.

Proof

The proof in Day 5 is a construction of \mathbb{R} by defining a specific order $<$.

Theorem 1.4.4: \mathbb{Q} is dense in \mathbb{R}

- (a) **Archimedean Property:** For $x, y \in \mathbb{R}$, if $x > 0$, there is $n \in \mathbb{Z}$ where $nx > y$.

Proof

Fix $x > 0$. Let $A = \{ nx : n = 1, 2, \dots \}$. Suppose there is a y where $nx \leq y$. Then, A is nonempty and bounded from above by y . By the least upper bound property of \mathbb{R} , $\alpha = \sup(A)$ exists in \mathbb{R} . Since $x > 0$, then $-x < 0$ so $\alpha - x < \alpha - 0 = \alpha$. So $\alpha - x$ is not an upper bound of A . So there is a $mx \in A$ such that $mx > \alpha - x$. Then $\alpha < (m+1)x$, but $(m+1)x \in A$ contradicting α is an upper bound for A .

- (b) **\mathbb{Q} is dense in \mathbb{R} :** For $x, y \in \mathbb{R}$, if $x < y$, there is a $p \in \mathbb{Q}$ where $x < p < y$.

Proof

Since $x < y$, then $y - x > 0$. Then by the Archimedean Property, there exists $n \in \mathbb{Z}$ such that $n(y - x) > 1$. Thus, $ny > nx + 1 > nx$. Since there is a smallest $m \in \mathbb{Z}_+$ such that $m > nx$, then $m > nx \geq m - 1$ so $nx + 1 \geq m > nx$. Since $ny > nx + 1 \geq m > nx$, then $y > m/n > x$.

2 Roots, Complex Field, & Euclidean Spaces

2.1 nth Root

Theorem 2.1.1: nth Root

(a) If $0 < t \leq 1$, then $t^n \leq t$.

Proof

Since $t > 0$ and $t \leq 1$, then $t^2 \leq t$.
 Since $t^2 \leq t$, then $t^3 \leq t^2$ so $t^3 \leq t^2 \leq t$.
 Applying the process n times, then $t^n \leq t$.

(b) If $t \geq 1$, $t^n \geq t$.

Proof

Since $0 < 1 \leq t$, then $t \leq t^2$.
 Since $t \leq t^2$, then $t^2 \leq t^3$ so $t \leq t^2 \leq t^3$.
 Applying the process n times, $t \leq t^n$.

(c) If $0 < s < t$, then $s^n < t^n$.

Proof

$$\underbrace{s \cdot s \cdot \dots \cdot s}_n < t \cdot s \cdot \dots \cdot s < t \cdot t \cdot \dots \cdot s < \dots < \underbrace{t \cdot \dots \cdot t}_n$$

Theorem 2.1.2: $y^n = x$ has a unique y

Fix $n \in \mathbb{Z}_+$. For every $x > 0$, there exists a unique $y \in \mathbb{R}$ such that $y^n = x$.

Also, such a y is written as $y = \sqrt[n]{x} = x^{\frac{1}{n}}$.

Proof

Uniqueness:

y is unique since if $y_1 < y_2$, then $x = y_1^n < y_2^n \neq x$.

Existence:

Let set $A = \{ t > 0 : t^n < x \}$.

$A \neq \emptyset$ since let $t_1 = \frac{x}{x+1} < 1$ so $t_1 < x$ and thus, $0 < t_1^n < t_1 < x$ so $t_1 \in A$.

A is bounded above since if $t \geq x+1$, then $t > 1$ so $t^n \geq t \geq x+1 > x$ so $t \notin A$.

So $x+1$ is an upper bound of A .

Thus by the least upper bound property, $y = \sup(A)$ exists.

For $y^n = x$, show $y^n < x$ and $y^n > x$ cannot hold true.

*** (Not an upper bound of A if $<$ and not a least upper bound of A if $>$)***

For $0 < \alpha < \beta$:

$$\beta^n - \alpha^n = (\beta - \alpha) \underbrace{(\beta^{n-1} + \beta^{n-2}\alpha^1 + \dots + \alpha^{n-1})}_{\substack{\beta^{n-1} < \beta^{n-1} < \beta^{n-1}}} < (\beta - \alpha)n\beta^{n-1}$$

Suppose $y^n < x$. Pick $0 < h < 1$ and $h < \frac{x - y^n}{n(y+1)^{n-1}}$.

From inequality, let $\beta = y+h$ and $\alpha = y$

$$(y+h)^n - y^n < hn(y+h)^{n-1} < hn(y+1)^{n-1} < x - y^n$$

Thus, $(y+h)^n < x$ so $y+h \in A$ and thus, not an upper bound of A which is a contradiction since $y = \sup(A)$.

Suppose $y^n > x$. Pick $0 < k = \frac{y^n - x}{ny^{n-1}} < \frac{y^n}{ny^{n-1}} = \frac{1}{n}y < y$.

$$\text{Consider } t \geq y-k, \text{ then: } y^n - t^n \leq y^n - (y-k)^n < kny^{n-1} = y^n - x$$

Thus, $t^n > x$ so $t \notin A$.

Thus, $y-k$ is an upper bound of A which is a contradiction since $y = \sup(A)$.

Since $y^n < x$ and $y^n > x$, then $y^n = x$.

Corollary 2.1.3: n-th root of product = product of n-th root

If $a, b > 0$ and $n \in \mathbb{Z}_+$, then $(ab)^{\frac{1}{n}} = a^{\frac{1}{n}} b^{\frac{1}{n}}$

Proof

Let $A = a^{\frac{1}{n}}$, $B = b^{\frac{1}{n}}$. By **theorem 2.1.2**, since A is a root for $y_1^n = a$, then $A^n = a$. Similarly, B is a solution of $y_2^n = b$ so $B^n = b$. Thus:

$$\begin{aligned} ab &= A^n B^n = A_1 A_2 \dots A_n B_1 B_2 \dots B_n \\ &= A_1 A_2 \dots B_1 A_n B_2 \dots B_n = \dots = A_1 B_1 A_2 \dots A_{n-1} A_n B_2 \dots B_n \\ &= \dots = A_1 B_1 A_2 B_2 \dots A_n B_n = (AB)^n \end{aligned}$$

Then again by **theorem 2.1.2**, there is a unique $(ab)^{\frac{1}{n}} = AB = a^{\frac{1}{n}} b^{\frac{1}{n}}$.

2.2 Decimals**Definition 2.2.1: Decimals**

Let n_0 be the largest integer such that $n_0 \leq x$ for $x > 0 \in \mathbb{R}$.

Then let n_k be the largest integer such that $d_k = n_0 + \frac{n_1}{10} + \dots + \frac{n_k}{10^k} \leq x$

Let E be the set of d_k for $k = 0, 1, \dots, \infty$. Then, $x = \sup(E)$.

2.3 Extended Reals**Definition 2.3.1: Extended Reals**

The extended real number system consist of \mathbb{R} and $\pm\infty$ such that:

$$-\infty < x < \infty \quad \text{for every } x \in \mathbb{R}$$

with the properties:

- $x \pm \infty = \pm\infty$
- $x / \pm\infty = 0$
- If $x > 0$, then $x(\pm\infty) = \pm\infty$. If $x < 0$, then $x(\pm\infty) = \mp\infty$

2.4 Complex Numbers**Definition 2.4.1: Complex Number**

A complex number is an ordered pair (a, b) where $a, b \in \mathbb{R}$. For $x, y \in \mathbb{C}$

- $x + y = (a, b) + (c, d) = (a + c, b + d)$
- $xy = (a, b)(c, d) = (ac - bd, ad + bc)$
- $\frac{1}{x} = (a^2 + b^2)^{-1}(a, -b)$

Thus, the axioms form a field where $(0, 0) = 0$ and $(1, 0) = 1$ and $(0, 1) = i$.

Theorem 2.4.2: Imaginary i and Form a + bi

Let $i = (0, 1)$. Then, $i^2 = -1$.

Then, $(a, b) = a + bi$

Proof

$$\begin{aligned} i^2 &= (0, 1)(0, 1) = (0 - 1, 0 + 0) = (-1, 0) = -1 \\ (a, b) &= (a, 0) + (0, b) = (a, 0) + (b, 0)(0, 1) = a + bi \end{aligned}$$

Definition 2.4.3: Conjugate

Let conjugate: $\bar{z} = a - bi$ where $\text{Re}(z) = a$, $\text{Im}(z) = b$.

Let $z = (a,b)$ and $w = (c,d)$:

(a) $\overline{z+w} = \bar{z} + \bar{w}$

Proof

$$\overline{z+w} = \overline{(a+c, b+d)} = (a+c, -b-d) = (a, -b) + (c, -d) = \bar{z} + \bar{w}$$

(b) $\overline{zw} = \bar{z} \bar{w}$

Proof

$$\overline{zw} = \overline{(ac-bd, ad+bc)} = (ac-bd, -ad-bc) = (a, -b)(c, -d) = \bar{z} \bar{w}$$

(c) $z + \bar{z} = 2 \text{Re}(z)$ $z - \bar{z} = 2i \text{Im}(z)$

Proof

$$\begin{aligned} z + \bar{z} &= (a,b) + (a,-b) = (2a,0) = 2 \text{Re}(z) \\ z - \bar{z} &= (a,b) - (a,-b) = (0,2b) = (0,2)b = 2i \text{Im}(z) \end{aligned}$$

(d) $z\bar{z} \geq 0$

Proof

$$z\bar{z} = (a,b)(a,-b) = (a^2 + b^2, -ab+ab) = a^2 + b^2 \geq 0$$

Definition 2.4.4: Absolute Value

Let absolute value: $|z| = \sqrt{z\bar{z}}$

Let $z = (a,b)$ and $w = (c,d)$:

(a) If $z \neq 0$, then $|z| > 0$.

Proof

$$\sqrt{z\bar{z}} = \sqrt{a^2 + b^2} \geq 0 \text{ where } |z| = 0 \text{ only if } a,b = 0 \text{ so only if } z = (0,0).$$

(b) $|\bar{z}| = |z|$

Proof

$$|\bar{z}| = \sqrt{a^2 + (-b)^2} = \sqrt{a^2 + b^2} = |z|$$

(c) $|zw| = |z| |w|$

Proof

$$\begin{aligned} |zw| &= |(ac-bd, ad+bc)| = \sqrt{(ac-bd)^2 + (ad+bc)^2} \\ &= \sqrt{a^2c^2 + b^2d^2 + a^2d^2 + b^2c^2} = \sqrt{(a^2 + b^2)(c^2 + d^2)} \\ &= \sqrt{a^2 + b^2} \sqrt{c^2 + d^2} = |z| |w| \end{aligned}$$

(d) $|\text{Re}(z)| \leq |z|$

Proof

$$|\text{Re}(z)| = |a| = \sqrt{a^2} \leq \sqrt{a^2 + b^2} = |z|$$

(e) $|z+w| \leq |z| + |w|$

Proof

$$\begin{aligned} |z+w|^2 &= (z+w)\overline{(z+w)} = (z+w)(\bar{z} + \bar{w}) = z\bar{z} + z\bar{w} + w\bar{z} + w\bar{w} \\ &= |z|^2 + |w|^2 + 2 \text{Re}(z\bar{w}) \leq |z|^2 + |w|^2 + 2|z\bar{w}| \\ &= |z|^2 + |w|^2 + 2|z||w| = (|z| + |w|)^2 \end{aligned}$$

2.5 Euclidean Spaces

Definition 2.5.1: Euclidean Spaces

For each positive integer k , let \mathbb{R}^k be the set of all ordered k -tuples:

$$x = (x_1, \dots, x_k) \quad \text{for each } x_i \in \mathbb{R}$$

with the properties:

- $x+y = (x_1 + y_1, \dots, x_k + y_k) \in \mathbb{R}^k$
- $cx = (cx_1, \dots, cx_k) \in \mathbb{R}^k$

So, \mathbb{R}^n has a vector space structure. Similarly, for \mathbb{C}^n .

Definition 2.5.2: Inner Product for \mathbb{R}^k

$$x \cdot y = x_1 y_1 + \dots + x_k y_k \in \mathbb{R}$$

Definition 2.5.3: Norm

$$|x| = \sqrt{x \cdot x} = \sqrt{\sum_{i=1}^k x_i^2}$$

Definition 2.5.4: Extension to \mathbb{C}^k

For $z, w \in \mathbb{C}^n$

- $z \cdot w = z_1 \overline{w_1} + \dots + z_k \overline{w_k}$
- $z \cdot z = z_1 \overline{z_1} + \dots + z_k \overline{z_k} = |z_1|^2 + \dots + |z_k|^2 = |z|^2$

2.6 Cauchy-Schwarz

Theorem 2.6.1: Cauchy-Schwarz

If $\alpha_1, \dots, \alpha_n \in \mathbb{C}$ and $b_1, \dots, b_n \in \mathbb{C}$, then:

$$|\sum_{j=1}^n \alpha_j \overline{b_j}|^2 \leq \sum_{j=1}^n |\alpha_j|^2 \sum_{j=1}^n |b_j|^2$$

Proof

Let $A = \sum |a_j|^2$ and $B = \sum |b_j|^2$ and $C = \sum a_j \overline{b_j}$.

If $B = 0$, then $b_1 = \dots = b_n = 0$. Thus, $0 \leq A(0)$ holds true.

Suppose $B > 0$. Then:

$$\begin{aligned} \sum |Ba_j - Cb_j|^2 &= \sum (Ba_j - Cb_j) \overline{(Ba_j - Cb_j)} = \sum (Ba_j - Cb_j) (\overline{B} \overline{a_j} - \overline{C} \overline{b_j}) \\ &= \sum (Ba_j - Cb_j) (\overline{B} \overline{a_j} - \overline{C} \overline{b_j}) = \sum B^2 a_j \overline{a_j} - B \overline{C} a_j \overline{b_j} - B C \overline{a_j} b_j + C \overline{C} b_j \overline{b_j} \\ &= B^2 \sum |a_j|^2 - B \overline{C} \sum a_j \overline{b_j} - B C \sum \overline{a_j} b_j + |C|^2 \sum |b_j|^2 \\ &= B^2 A - B \overline{C} C - B C \overline{C} + |C|^2 B = B^2 A - 2|C|^2 B + |C|^2 B = B^2 A - |C|^2 B \\ &= B(AB - |C|^2) \end{aligned}$$

Since $|Ba_j - Cb_j|^2 \geq 0$, then $B(AB - |C|^2) \geq 0$.

Since $B > 0$, then $AB - |C|^2 \geq 0$ so $AB \geq |C|^2$.

Corollary 2.6.2: $|z \cdot w| \leq |z||w|$

Since $|z_i|^2 = z_i \overline{z_i}$, then $\sum z_i \overline{z_i} = \sum |z_i|^2 = |z|^2$. Thus:

$$|z \cdot w|^2 = |\sum z_i \overline{w_i}|^2 \leq \sum |z_i|^2 \sum |w_i|^2 = |z|^2 |w|^2$$

Thus, $|z \cdot w| \leq |z||w|$.

Theorem 2.6.3: Properties of \mathbb{R}^k

Let $x, y, z \in \mathbb{R}^k$ where $\alpha \in \mathbb{R}$:

- (a) $|x| \geq 0$ where $|x| = 0$ only if $x = 0$

Proof

$$|x| = \sqrt{\sum_{i=1}^k x_i^2} \geq 0 \text{ where } |x| = 0 \text{ only if } x_1 = \dots = x_k = 0$$

- (b) $|\alpha x| = |\alpha||x|$

Proof

$$|\alpha x| = \sqrt{\sum_{i=1}^k (\alpha x_i)^2} = \sqrt{\alpha^2} \sqrt{\sum_{i=1}^k x_i^2} = |\alpha||x|$$

- (c) $|x + y| \leq |x| + |y|$

Proof

$$\begin{aligned} |x + y|^2 &= (x + y) \cdot (x + y) = |x|^2 + 2(x \cdot y) + |y|^2 \\ &\leq |x|^2 + 2|x||y| + |y|^2 = (|x| + |y|)^2 \end{aligned}$$

- (d) $|x - y| \leq |x - z| + |y - z|$

Proof

$$|x - y| = |x - z + z - y| \leq |x - z| + |z - y| = |x - z| + |y - z|$$

3 Construction of \mathbb{R}

There exists an ordered field \mathbb{R} which has the least upper bound property.
Also, \mathbb{R} contains \mathbb{Q} as a subfield.

Definition 5.1: Cuts

Define a cut as any set $\alpha \subset \mathbb{Q}$ with the properties:

- α is not empty and $\alpha \neq \mathbb{Q}$
- If $p \in \alpha$ and $q \in \mathbb{Q} < p$, then $q \in \alpha$
- If $p \in \alpha$, then $p < r \in \mathbb{Q}$ for some $r \in \alpha$

Proposition 5.2: Order of $\mathbb{R} \rightarrow$ ordered set \mathbb{R}

Define $\alpha < \beta$ if α is a proper subset of β .

- If $\alpha \not\subseteq \beta$, then β is not a subset of α .
Then there is a $p \in \beta$ such that $p \notin \alpha$.
Then for any $q \in \alpha$, $q < p$ and thus, $q \in \beta$.
Thus, $\alpha \subset \beta$ and since $\alpha \neq \beta$, then $\alpha < \beta$.
- If $\alpha \not\subseteq \beta$ and $\alpha \not\supseteq \beta$, then either $\alpha = \beta$ or $\alpha \neq \beta$.
If $\alpha \neq \beta$, there are p, q such that $p \in \alpha$, but $p \notin \beta$ and $q \in \beta$, but $q \notin \alpha$.
But if $p \notin \beta$, then for any $b \in \beta$, $b < p$. Thus, $q < p$.
Similarly, if $q \notin \alpha$, then for any $a \in \alpha$, $a < q$. Thus, $p < q$.
Thus, there is a contradiction since $p > q$ and $p < q$ so $\alpha = \beta$.
- If $\alpha \not\subseteq \beta$, then α is not a subset of β .
Then there is a $p \in \alpha$ such that $p \notin \beta$.
Then for any $q \in \beta$, $q < p$ and thus, $q \in \alpha$.
Thus, $\beta \subset \alpha$ and since $\alpha \neq \beta$, then $\beta < \alpha$.
- If $\alpha < \beta$ and $\beta < \gamma$, then since α is a proper subset of β and β is a proper subset of γ , then α is a proper subset of γ . Thus, $\alpha < \gamma$.

Thus, \mathbb{R} is an ordered set with such an order $<$.

Proposition 5.3: Least Upper Bound of $\mathbb{R} \rightarrow$ Least Upper Bound Property

Let $A \subset \mathbb{R}$ and β be an upper bound for A . Let γ be the union of all $\alpha \in A$.
Thus, $p \in \gamma$ if and only if $p \in \alpha$ for some $\alpha \in A$.

γ defines a cut since:

- Since A is nonempty, there exists a $\alpha_0 \in A$ where α_0 is nonempty.
Since α_0 is nonempty, then γ is nonempty.
Since every $\alpha \in A$ is $\alpha < \beta$, then $\gamma < \beta$ so $\gamma \subset \beta$ and thus, $\gamma \neq \mathbb{Q}$.
- If $p \in \gamma$, then $p \in \alpha_1$ for some $\alpha_1 \in A$. If $q < p$, then $q \in \alpha_1$ so $q \in A$.
- If $p \in \gamma$, then $p \in \alpha_1$ for some $\alpha_1 \in A$. Thus, there is a $r \in \alpha_1$ such that $r > p$ so $r \in \gamma$. Thus, there is a $r \in \gamma$ where $r > p$.

Since γ defines a cut, then $\gamma \in \mathbb{R}$. Since every $\alpha \in A \subset \gamma$, then $\alpha \leq \gamma$ so γ is an upper bound for A .

Suppose $\delta < \gamma$. Then there is a $s \in \gamma$ such that $s \notin \delta$. Since $s \in \gamma$, then there is a $\alpha \in A$ such that $s \in \alpha$. Since $\delta < \alpha$, then δ is not an upper bound of A .

Thus, $\gamma = \sup(A)$.

Proposition 5.4: \mathbb{R} is a field

If $\alpha, \beta \in \mathbb{R}$, define $\alpha + \beta$ as the set of all sums $r + s$ where $r \in \alpha$ and $s \in \beta$. Also, let 0^* be the set of all negative rational numbers which is a cut since:

- 0^* is nonempty and $0^* \neq \mathbb{Q}$
- If $p \in 0^*$, then any $q \in \mathbb{Q} < p$ is a negative rational and thus, $q \in 0^*$.
- Since \mathbb{Q} is dense in \mathbb{R} , then for any $p \in 0^*$, there is a $r \in \mathbb{Q}$ where $p < r < 0$ so r is a negative rational so $r \in 0^*$.

$\alpha + \beta \in \mathbb{R}$ since $\alpha + \beta$ is a cut:

- $\alpha + \beta$ is non-empty since α, β are non-empty. Take $r' \notin \alpha, s' \notin \beta$, then $r' + s' > r + s$ for $r \in \alpha, s \in \beta$. Thus, $r' + s' \notin \alpha + \beta$ so $\alpha + \beta \neq \mathbb{Q}$.
- If $p \in \alpha + \beta$, then $p = r + s$ where $r \in \alpha$ and $s \in \beta$.
If $q < p$, then $q - s < p - s = (r + s) - s = r$ so $q - s \in \alpha$.
Since $q - s \in \alpha$ and $s \in \beta$, then $(q - s) + s = q \in \alpha + \beta$.
- If $r \in \alpha$, then there is a $t \in \alpha$ such that $t > r$. Let $s \in \beta$.
Thus, for any $p = r + s \in \alpha + \beta$, there is a $q = t + s \in \alpha + \beta$ such that $p = r + s < t + s = q$.

$\alpha + \beta = \beta + \alpha$

If $p = r + s \in \alpha + \beta$ where $r \in \alpha, s \in \beta$, then $s + r = r + s = p \in \beta + \alpha$.

$(\alpha + \beta) + \gamma = \alpha + (\beta + \gamma)$

If $r \in \alpha, s \in \beta, t \in \gamma$, then $r + s + t = (r + s) + t \in (\alpha + \beta) + \gamma$ and $r + s + t = r + (s + t) \in \alpha + (\beta + \gamma)$.

$\alpha + 0^* = \alpha$

If $r \in \alpha, s \in 0^*$, then $r + s < r$. Thus, $r + s \in \alpha$. Thus, $\alpha + 0^* \subset \alpha$.

If $p \in \alpha$, there is a $r \in \alpha$ where $r > p$. Thus, $p - r \in 0^*$.

Since $p = r + (p - r) \in \alpha + 0^*$, then $\alpha \subset \alpha + 0^*$. Thus, $\alpha + 0^* = \alpha$.

There is a $-\alpha$ such that $\alpha + (-\alpha) = 0^*$

Fix $\alpha \in \mathbb{R}$. Let set β be all p where there is $r > 0$ such that $-p - r \notin \alpha$.

$\beta \in \mathbb{R}$ since β is a cut:

- If $s \notin \alpha$ and $p = -s - 1$, then $-p - 1 \notin \alpha$. Thus, $p \in \beta$ so β is nonempty. If $q \in \alpha$, then $-q \notin \beta$ so $\beta \neq \mathbb{R}$.
- If $p \in \beta$, let $r > 0$ so $-p - r \notin \alpha$. If $q < p$, then $-q - r > -p - r$ and thus, $-q - r \notin \alpha$ so $q \in \beta$.
- If $p \in \beta$, let $t = p + (r/2)$. Then $-t - (r/2) = -p - r \notin \alpha$ and thus, $t \in \beta$ where $p < t$.

If $r \in \alpha, s \in \beta$, then $s \notin \alpha$. Thus, $r < -s$ so $r + s < 0$. Thus, $\alpha + \beta \subset 0^*$.

Let $v \in 0^*$ and let $w = -v/2$ so $w > 0$.

Thus, by the Archimedean property, there is an integer n such that $nw \in \alpha$, but $(n+1)w \notin \alpha$. Let $p = -(n+2)w$ so $-p - w = (n+1)w \notin \alpha$ so $p \in \beta$.

Then, $v = -2w = nw + -nw - 2w = nw + -(n+2)w = nw + p \in \alpha + \beta$.

Since $v \in 0^*$, then $0^* \subset \alpha + \beta$. Thus, $\alpha + \beta = 0^*$. Then, let $-\alpha = \beta$.

Thus, if $\alpha, \beta, \gamma \in \mathbb{R}$ and $\beta < \gamma$, then $\alpha + \beta < \alpha + \gamma$.

Thus, if $\alpha > 0^*$, then $-\alpha = -\alpha + 0^* < -\alpha + \alpha = 0^*$ so $-\alpha < 0^*$.

If $\alpha, \beta \in \mathbb{R}_+$, define $\alpha\beta$ as the set of all p such that $p \leq rs$ for $r \in \alpha, s \in \beta$.

Define 1^* as the set of all $q < 1$. Then all multiplication axioms holds with similar proofs as addition. Also, note since $\alpha, \beta > 0^*$, then $\alpha\beta > 0^*$.

Also, $\alpha(\beta + \gamma) = \alpha\beta + \alpha\gamma$ holds through cases were $\alpha, \beta, \gamma >, < 0^*$.

4 Cardinality


4.1 Cardinality

Definition 4.1.1: Onto and 1-1 Mapping

Suppose for every $x \in A$, there is an associated $f(x) \in B$.

Then f maps A into $B = f: A \rightarrow B$.

- If $f(A) = B$, then f maps A onto B .
- If for each $y \in B$, $f^{-1}(y)$ consist of at most one $x \in A$ where $f^{-1}(y_1) = x_1 \neq x_2 = f^{-1}(y_2)$ for $y_1 \neq y_2$, then f is a 1-1 mapping of A into B .



Definition 4.1.2: 1-1 Correspondence

Sets A and B are equivalent (have the same cardinality) if there is a 1-1 onto function $f: A \rightarrow B$. (1-1 correspondence between A and B) Then, $A \sim B$.

If $f: A \rightarrow B$ is 1-1 and onto, then there is a $f^{-1}: B \rightarrow A$ that is 1-1 and onto.

Definition 4.1.3: Countability

- A is **finite** if $A \sim J_n = \{0, 1, \dots, n\}$ for some $n \in \mathbb{N}$
- A is **infinite** if A is not finite
- A is **countably infinite** if $A \sim J = \mathbb{Z}_+$
- A is **uncountable** if A is not finite or countably infinite
- A is **at most countable** if A is finite or countably infinite

Example

\mathbb{Z} is countably infinite


Proof

$$\text{Let } f(n): \mathbb{Z}_+ \rightarrow \mathbb{Z} = \begin{cases} \frac{n}{2} & n \text{ is even} \\ -\frac{n-1}{2} & n \text{ is odd} \end{cases}$$

So $1 \mapsto 0$, $2 \mapsto 1$, $3 \mapsto -1$, $4 \mapsto 2$, $5 \mapsto -2$, etc. Thus, $\mathbb{Z} \sim \mathbb{Z}_+$.

Definition 4.1.4: Pigeonhole Principle

If A is finite, A is not equivalent to any proper set of A .



Theorem 4.1.5: Infinite subsets of countable sets are countable

An infinite subset E of a countably infinite set A is countably infinite

Proof

Let $E \subset A$ be an infinite subset. For every distinct $x_i \in A$, let $\{x_1, x_2, \dots\} \in A$.
 Let n_1 be smallest integer such that $x_{n_1} \in E$.
 Then let n_2 be the smallest integer where $n_2 > n_1$ such that $x_{n_2} \in E$.
 Repeat the process to create sequence $f(k) = \{x_{n_1}, x_{n_2}, \dots, x_{n_k}, \dots\}$.
 Thus, there is a 1-1 correspondence between E and \mathbb{Z}_+ so E is countably infinite.

**4.2 Set of Sets****Definition 4.2.1: Union and Intersection**

Let sets Ω, B be such that for each $x \in \Omega$, there is an associated $E_x \subset B$.

- $E = \cup_{x=1}^n E_x$ only if for every $x \in E$, $x \in E_x$ for at least one $x \in \Omega$.
- $P = \cap_{x=1}^n E_x$ only if for every $x \in P$, $x \in E_x$ for all $x \in \Omega$.

with properties:

- | | |
|---|---|
| (a) $A \cup B = B \cup A$ | (a) $A \cap B = B \cap A$ |
| (b) $(A \cup B) \cup C = A \cup (B \cup C)$ | (b) $(A \cap B) \cap C = A \cap (B \cap C)$ |
| (c) $A \subset A \cup B$ | (c) $(A \cap B) \subset A$ |
| (d) If $A \subset B$, then $A \cup B = B$ and $A \cap B = A$ | |

Proof

If $x \in A \cup B$, then $x \in A$ or/and $x \in B$.

- If $x \in A$, since $A \subset B$, then $x \in B$. Then, $(A \cup B) \subset B$.
- If $x \in B$, then immediately $(A \cup B) \subset B$.

If $x \in B$, then $x \in A \cup B$ so $B \subset (A \cup B)$. Thus, $A \cup B = B$.

If $x \in A \cap B$, then $x \in A$ and $x \in B$. Thus, $(A \cap B) \subset A$.

If $x \in A$, since $A \subset B$, then $x \in B$ so $x \in A \cap B$. Thus, $A \subset (A \cap B)$.

Thus, $A \cap B = A$.

- (e) $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$

Proof

If $x \in A \cap (B \cup C)$, then $x \in A$ and ($x \in B$ or/and $x \in C$).

- If $x \in B$, then $x \in (A \cap B)$ so $x \in (A \cap B) \cup (A \cap C)$.
- If $x \in C$, then $x \in (A \cap C)$ so $x \in (A \cap B) \cup (A \cap C)$.

Thus, $A \cap (B \cup C) \subset (A \cap B) \cup (A \cap C)$.

If $x \in (A \cap B) \cup (A \cap C)$, then $x \in A$ and ($x \in B$ or/and $x \in C$).

Thus, $(A \cap B) \cup (A \cap C) \subset A \cap (B \cup C)$.

Thus, $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$.

$$(f) A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$$

Proof

If $x \in A \cup (B \cap C)$, then $x \in A$ or/and $(x \in B \text{ and } x \in C)$.

- If $x \in A$, then $x \in (A \cup B)$ and $x \in (A \cup C)$ so $A \cup (B \cap C) \subset (A \cup B) \cap (A \cup C)$.
- If $x \in B, C$, then $x \in (A \cup B)$ and $x \in (A \cup C)$ so $A \cup (B \cap C) \subset (A \cup B) \cap (A \cup C)$.

If $x \in (A \cup B) \cap (A \cup C)$, then $x \in A$ or/and $(x \in B \text{ and } x \in C)$.

Thus, $(A \cup B) \cap (A \cup C) \subset A \cup (B \cap C)$.

Thus, $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$.

Theorem 4.2.2: Union of countably infinite sets is countably infinite

If E_1, E_2, \dots are countably infinite sets, then $S = \bigcup_{n=1}^{\infty} E_n$ is countably infinite.

Proof

For each E_n , there is a sequence $\{x_{n1}, x_{n2}, \dots\}$. Then construct an array as such:

$$\begin{pmatrix} x_{11} & x_{12} & \dots \\ x_{21} & x_{22} & \dots \\ \vdots & \vdots & \ddots \end{pmatrix}$$

Take elements diagonally, then sequence $S^* = \{x_{11}; x_{21}, x_{12}; x_{31}, x_{22}, x_{13}; \dots\}$.

Since $S^* \sim S$ so S is at most countable and S is infinite since E_1, E_2, \dots are infinite, then S cannot be finite and thus, countably infinite.

Alternative Proof


For each E_n , let set $\widetilde{E}_n = E_n - \bigcup_{m=1}^{\infty} E_m$ where $m \neq n$. Thus, $S = \bigcup_{n=1}^{\infty} \widetilde{E}_n$.

Since each E_n is countably infinite, there exists a 1-1 mapping $\delta_n: E_n \rightarrow \mathbb{Z}_+$.

Thus, for each \widetilde{E}_n , there is a 1-1 mapping $\delta_n: \widetilde{E}_n \rightarrow A \subset \mathbb{Z}_+$.

Let p_1, p_2, \dots be distinct primes. Since for $s \in S$, there exists a unique \widetilde{E}_i such that $s \in \widetilde{E}_i$, then let $f(s) = p_1^{\delta_1(s)} p_2^{\delta_2(s)} \dots$ where $p_k^{\delta_k(s)} = 1$ if $k \neq i$.

Then, by the Fundamental theorem of arithmetic, f maps s to a unique $z \in \mathbb{Z}_+$ and thus, f is a 1-1 function so S is at most countable. Since any $E_n \subset S$ is countably infinite, then S cannot be finite and thus, S is countably infinite.



Theorem 4.2.3: The set of countable n-tuples are countable

Let A be a countably infinite set and B_n be the set of all n -tuples (a_1, \dots, a_n) where $a_k \in A$. Then B_n is countably infinite.

Proof

The base case B_1 is countably infinite since $B_1 = A$.

Suppose B_{n-1} is countably infinite. Then for every $x \in B$:

$$x = (b, a) \quad b \in B_{n-1} \text{ and } a \in A$$

Since for every fixed b , $(b, a) \sim A$ and thus, countably infinite.

Since B is a set of countably infinite sets, then B_n is countably infinite.

Theorem 4.2.4: \mathbb{Q} is countable

The set of rational numbers, \mathbb{Q} , is countably infinite

Proof

Since elements of \mathbb{Q} are of form $\frac{a}{b}$ which is a 2-tuple, then by the **theorem 4.2.3**, \mathbb{Q} is countably infinite.

Alternative Proof

For every $x \in \mathbb{Q}$, let $x = (-1)^i \frac{p}{q}$ where $p, q \in \mathbb{Z}_+$.
 Let $f(x) = 2^i 3^p 5^q$. Then by the Fundamental theorem of arithmetic, f is a 1-1 mapping of x to $E \subset \mathbb{Z}_+$.
 Thus, \mathbb{Q} is at most countable, but since $p, q \in \mathbb{Z}_+$, then \mathbb{Q} cannot be finite and thus, is countably infinite.

Example

Let A be the set of all sequences whose elements are digits 0 and 1. Then A is uncountable.

Proof: Cantor's Diagonalization Proof

Let set E be a countably infinite subset of A which consist of sequences s_1, s_2, \dots .
 Then construct a sequence s as follows:
 If the n -th digit in s_n is 1, then let the n -th digit of s be 0 and vice versa.
 Thus, s differs from every $s_n \in E$ so $s \notin E$.
 But, $s \in A$ so E is a proper subset of A .
 Thus, every countably infinite subset of A is a proper subset of A .
 If A is countably infinite, then A is a proper subset of A which is a contradiction.

5 Metric Spaces & Closed/Open

5.1 Metric Spaces

Definition 5.1.1: Metric Spaces

A set X is a metric space if for any $p, q \in X$, there is an associated $d(p, q) \in \mathbb{R}$ such that:

- $d(p, q) > 0$ if $p \neq q$
- $d(p, q) = 0$ if and only if $p = q$
- **Symmetry**: $d(p, q) = d(q, p)$
- **Triangle Inequality**: $d(p, q) \leq d(p, r) + d(r, q)$ for any $r \in X$.


For euclidean spaces \mathbb{R}^k , $d(x, y) = |x - y|$ where $x, y \in \mathbb{R}^k$.

Definition 5.1.2: Types of Points and Sets

For metric space X and set $E \subset X$:

(a) Neighborhood

For $p \in X$ and $r > 0$, $N_r(p)$ is the set of all $q \in X$ where $d(q, p) < r$



(b) Limit Points and Closed Sets

Closed set E contain all $p \in X$ where every $N_r(p)$ contain a $q \neq p \in E$

• Limit Points

For point $p \in X$, every $N_r(p)$ contains a $q \neq p \in E$


The set of all limit points of $E = E'$

• Isolated Points

If $p \in E$ is not a limit point of E

• Closed

If every limit point p of E is a $p \in E$



(c) Interior Points and Open Sets

Open set E contains all its p which has a $N_r(p) \subset E$


- Interior Point

For $p \in X$, there is a $N_r(p) \subset E$

The set of all interior points = E°

- Open

If every $p \in E$ is an interior point of E



(d) More about Sets

- Bounded

If there is $M \in \mathbb{R}$, $q \in X$ such that $d(p, q) < M$ for all $p \in E$

- Complement

From E , E^c is the set of all $p \in X$ such that $p \notin E$

- Perfect

If E is closed and if every $p \in E$ is a limit point of E

- Dense

If every $p \in X$ is a limit point of E or/and $p \in E$

- Boundary Point

For $p \in X$, if every $N_r(p)$ contains a $x \in E$ and $y \in E^c$

The set of all boundary points = ∂E


For a metric space X , $\{X, \emptyset\}$ are both open and closed.

Theorem 5.1.3: $N_r(p)$ is open

Every neighborhood is an open set

Proof

Let $q \in N_r(p)$. Then there is a $h > 0 \in \mathbb{R}$ such that $d(q, p) = r - h$.
 Then for any $s \in N_h(q)$, $d(s, p) \leq d(s, q) + d(q, p) = h + (r - h) = r$.
 Thus, for any $q \in N_r(p)$, there exists a $N_h(q) \subset N_r(p)$.




Theorem 5.1.4: If a set has a limit point, there are infinite $q \in E$ in $N_r(p)$

If p is a limit point of set E , then every $N_r(p)$ contains infinitely many $q \in E$.

Proof

Suppose there is $N_{r_1}(p)$ which contains finitely many $q = \{q_1, \dots, q_n\}$.
 Let $r = \min_{m \in [1, n]} d(p, q_m)$. Then $N_r(p)$ contains no $q \in E$ such that $q \neq p$.
 So, p is not a limit point of E which is a contradiction since p is a limit point of E .



Corollary 5.1.5: Limit points do not exist in finite sets

A finite set E has no limit points. Since $\emptyset \in E$, all finite set must be closed.

Proof


Let p be a limit point of finite set E . By **theorem 5.1.4**, then any $N_r(p)$ contain infinite $q \in E$ so E is an infinite set which is a contradiction since E is finite.
 So p cannot be limit point of E and thus, E has no limit points. Since finite set E contains all its limit points because there are no limit points, then E is closed.

Theorem 5.1.6: De Morgan's Laws

Let E_1, E_2, \dots be a collection of sets. Then, $(\cup E_x)^c = \cap (E_x^c)$.

Proof

If $p \in (\cup E_x)^c$, then $p \notin (\cup E_x)$.
 Thus, $p \notin E_x$ for any x so $p \in E_x^c$ for all x . Thus, $p \in \cap (E_x^c)$ so $(\cup E_x)^c \subset \cap (E_x^c)$.
 If $p \in \cap (E_x^c)$, then $p \in E_x^c$ for all x .
 Thus, $p \notin E_x$ for any x so $p \notin \cup E_x$. Thus, $p \in (\cup E_x)^c$ so $\cap (E_x^c) \subset (\cup E_x)^c$.
 Thus, $(\cup E_x)^c = \cap (E_x^c)$.



Theorem 5.1.7: Open set \rightarrow Closed complement

A set E is open if and only if E^c is closed

Proof

Suppose E is open. Let x be a limit point of E^c .

Then for every $r > 0$, $N_r(x)$ must contain a $p \in E^c$ such that $p \neq x$.

Then, $N_r(x) \not\subset E$ so x is not an interior point of E and thus, $x \notin E$ so $x \in E^c$.

Since any limit point x of E^c is a $x \in E^c$, then E^c is closed.

Suppose E^c is closed. Let $x \in E$.

Since $x \notin E^c$, x is not a limit point of E^c . Then there exists a $r > 0$ such that any $p \in N_r(x)$ is not in E^c . Thus, every $p \in N_r(x)$ is $p \in E$ so $N_r(x) \subset E$ and thus, x is an interior point of E . Since any $x \in E$ is an interior point of E , then E is open.

Corollary 5.1.8: Closed set \rightarrow Open complement

A set F is closed if and only if F^c is open.

Proof

From **theorem 5.1.7**, let $E = F^c$

Theorem 5.1.9: Union open \rightarrow open and Intersection closed \rightarrow closed

- (a) If $\{G_x\}$ is a finite or infinite collection of open sets, then $\cup G_x$ is open.

Proof

If $p \in \cup G_x$, then $p \in G_x$ for at least one x . Let \bar{x} be such an x .

Since $G_{\bar{x}}$ is open, then p is an interior point of $G_{\bar{x}}$ and thus, there is a $N_r(p)$ such that $N_r(p) \subset G_{\bar{x}} \subset \cup G_x$. So p is an interior point of $\cup G_x$.

Since any $p \in \cup G_x$ is an interior point, then $\cup G_x$ is open.

- (b) If $\{F_x\}$ is a finite or infinite collection of closed sets, then $\cap F_x$ is closed.

Proof

By **theorem 5.1.7**, any F_x^c is open. Since $\{F_x^c\}$ is a finite or infinite collection of open set, then by part (a), $\cup F_x^c$ is open.

Thus, again by **theorem 5.1.7**, $(\cup F_x^c)^c$ is closed.

By **theorem 5.1.6**, $(\cup F_x^c)^c = \cap (F_x^c)^c = \cap F_x$.

- (c) If G_1, \dots, G_n is a finite collection of open sets, then $\cap_{x=1}^n G_x$ is open.

Proof

If $p \in \cap_{x=1}^n G_x$, then $p \in G_x$ for all G_x for $x = \{1, 2, \dots, n\}$.

Since each G_x is open, then for any G_x , there is a $N_{r_x}(p) \subset G_x$.

Let $r = \min(r_1, r_2, \dots, r_n)$. Thus, $p \in N_r(p) \subset N_{r_x}(p)$ for all x .

So, $N_r(p) \subset \cap_{x=1}^n G_x$ and thus, p is an interior point of $\cap_{x=1}^n G_x$ so $\cap_{x=1}^n G_x$ is open.

Infinite + Closed: $G_i = (-1/i, 1/i)$

Infinite + Open: $G_i = (-i, i)$

- (d) If F_1, \dots, F_n is a finite collection of closed sets, then $\cup_{x=1}^n F_x$ is closed.

Proof

By **theorem 5.1.7**, any F_x^c is open. Since F_1^c, \dots, F_n^c is a finite collection of open set, then by part (c), $\cap_{x=1}^n F_x^c$ is open.

Thus, again by **theorem 5.1.7**, $(\cap_{x=1}^n F_x^c)^c$ is closed.

By **theorem 5.1.6**, $(\cap_{x=1}^n F_x^c)^c = \cup_{x=1}^n (F_x^c)^c = \cup_{x=1}^n F_x$.

Infinite + Closed: $F_i = [-1/i, 1/i]$

Infinite + Open: $F_i = [1/i, \infty)$

Theorem 5.1.10: E' is closed

Let $E \subset X$. Then, $(E')' \subset E'$. Thus, E' is closed.

Proof

If $x \in (E')'$, then for every $N_{r_1}(x)$, there is a $y \neq x$ where $y \in E'$.
 Since $y \in E'$, then for every $N_{r_2}(y)$, there is a $z \neq y$ where $z \in E$.
 Let $r = r_1 + r_2$.
 Then for every $N_r(x)$, there exists a $z \neq x$ where $z \in E$. Thus, $x \in E'$ so $(E')' \subset E'$.

Theorem 5.1.11: E° is open

Let $E \subset X$. Then, E° is open.

Proof

If $p \in E^\circ$, there is a $r > 0$ such that $N_r(p) \subset E$.
 Then for $0 < s < r$, $N_s(p) \subset N_r(p)$ so any $q \in N_s(p)$ is $q \in E^\circ$.
 Since any $p \in E^\circ$ have a $N_s(p) \subset E^\circ$, then E° is open.

5.2 Intervals and Balls**Definition 5.2.1: Segments and Intervals**

In \mathbb{R} , a **segment** is an open interval $(a,b) = \{ x \in \mathbb{R} : a < x < b \}$
 In \mathbb{R} , a **interval** is a closed interval $[a,b] = \{ x \in \mathbb{R} : a \leq x \leq b \}$

Definition 5.2.2: Open Balls

In \mathbb{R}^k , an **open ball** of radius $r > 0$ centered at p is:

$$N_r(p) = \{ x \in \mathbb{R}^k : |x - p| < r \} = \{ x \in \mathbb{R}^k : d(x,p) < r \}$$

A **closed ball** has $d(x,p) \leq r$.

Definition 5.2.3: Convex

$E \subset \mathbb{R}^k$ is **convex** if for all $x, y \in E$ and $t \in [0,1]$, $tx + (1-t)y \in E$.

Example

Balls in \mathbb{R}^k are convex

Let $x, y \in$ open ball $N_r(p)$. Let $z = tx + (1-t)y$ for $t \in [0,1]$.
 Since $|x - p| < r$ and $|y - p| < r$:

$$\begin{aligned} |z - p| &= |tx + (1-t)y - p| = |tx + (1-t)y - tp + (t-1)p| \\ &= |t(x-p) + (1-t)(y-p)| \leq t|(x-p)| + (1-t)|(y-p)| \\ &< tr + (1-t)r = r \end{aligned}$$

 Thus, $z \in N_r(p)$ so balls are convex. Same proof applies to closed balls.

Definition 5.2.4: Dense

$E \subset X$ is dense if every $x \in X$ is either in E or a limit point of E .

Example

Let $X = \mathbb{R}$. Then, $E = \mathbb{Q}$ is dense in \mathbb{R} .

Fix $x \in \mathbb{R}$ and $r > 0$. There is a $q \in \mathbb{Q}$ such that $x-r < q < x$. So for any $r > 0$ and $q \in \mathbb{Q}$, $q \neq x$ and $q \in N_r(x)$. Thus, every $x \in \mathbb{R}$ is a limit point of \mathbb{Q} .

6 Closure, Open Relative, & Compact

6.1 Closure

Definition 6.1.1: Closure

Let $E \subset$ metric space X and E' be the set of all limit points of E in X .

Then the closure of E : $\overline{E} = E \cup E'$

with the properties:

- (a) \overline{E} is closed

Proof

Suppose $x \in X$, but $x \notin \overline{E}$. Thus, $x \in \overline{E}^c$.

Thus, there is a $N_r(x) \subset \overline{E}^c$ since else there is always a $p \in N_r(x)$ where $p \in \overline{E}$ so x is a limit point of \overline{E} so $x \in \overline{E}$. Thus, \overline{E}^c is open so \overline{E} is closed by [theorem 5.1.7](#).

- (b) $E = \overline{E}$ if and only if E is closed

Proof

If $E = \overline{E}$, then by part (a), E is closed.

If E is closed, then $E' \subset E$ so $E = E \cup E' = \overline{E}$.

- (c) $\overline{E} \subset F$ for every closed $F \subset X$ such that $E \subset F$

Proof

If closed set F , then $F' \subset F$. Since $E \subset F$, then $E' \subset F' \subset F$ so $\overline{E} \subset F$.

Theorem 6.1.2: $\sup(E) \in \overline{E}$

Let non-empty set of real numbers, E , be bounded above. Let $y = \sup(E)$.

Then, $y \in \overline{E}$. Thus, $y \in E$ if E is closed and $y \notin E$ if E is open in \mathbb{R} .

Proof

If $y \in E$, then $y \in \overline{E}$. Suppose $y \notin E$.

For every $h > 0$, there exists a $x \in E$ such that $y-h < x < y$ otherwise $y-h$ is an upper bound for E which is a contradiction since $y = \sup(E)$.

Thus, y is a limit point of E so $y \in E'$.

If E is closed, then $y \in E$ since $y \in E'$. Also, $y \in \overline{E}$.

If E is open, then any $N_r(y) \not\subset E$ since $N_r(y)$ in \mathbb{R} must contain a $\gamma > y$ so $y \notin E^\circ$.

6.2 Open Relative

Definition 6.2.1: Open Relative

Suppose $E \subset Y \subset$ metric space X .

Then E is open relative to Y if for each $p \in E$:

There is an $r > 0$ such that for any $q \in Y$ where $d(q,p) < r$, then $q \in E$.

Theorem 6.2.2: E is open relative to $Y \subset X$ if $E = Y \cap G$ and G is open in X

Suppose $E \subset Y \subset X$.

E is open relative to Y if and only if $E = Y \cap G$ for some open $G \subset X$.

Proof

Suppose E is open relative to Y .

Then for each $p \in E$, there is a $r_p > 0$ such that for any $q \in Y$ where $d(p, q) < r_p$, then $q \in E$.

Since $Y \subset X$, let V_p be the set of all $q \in X$ such that $d(p, q) < r_p$ and define $G = \bigcup_{p \in E} V_p$. Since V_p is open by **theorem 5.1.3**, then by **theorem 5.1.9a**, open $G \subset X$.

Since $p \in V_p$ for all $p \in E$, then $E \subset G \cap Y$. Also, by construction, then $V_p \cap Y \subset E$ so $G \cap Y \subset E$. Thus, $E = Y \cap G$.

If G is open in X and $E = G \cap Y$, then every $p \in E$ has a $V_p \subset G$.

Then, $V_p \cap Y \subset G \cap Y = E$ so E is open relative to Y .

6.3 Compact Sets

Definition 6.3.1: Open Cover

An open cover of set $E \subset X$ is a collection of open $G_1, G_2, \dots \subset X$ such that $E \subset \bigcup G_i$.

Definition 6.3.2: Compact

$K \subset X$ is compact if every open cover of K contains a finite subcover.

If G_1, G_2, \dots is an open cover of K , then $K \subset \bigcup_{i=1}^n G_i$ for some n .

Theorem 6.3.3: A compact set is compact in every metric space

Suppose $K \subset Y \subset X$.

Then K is compact relative to X if and only if K is compact relative to Y .

Proof

Suppose K is compact relative to X .

Let V_1, V_2, \dots be sets open relative to Y such that $K \subset \bigcup V_x$. Then by **theorem 6.2.2** for each V_x , there is a G_x open relative to X where $V_x = Y \cap G_x$.

Since K is compact relative to X , then there is a n such that $K \subset G_{x_1} \cup \dots \cup G_{x_n}$.

Thus, $K = K \cap Y \subset (\bigcup_{i=1}^n G_{x_i}) \cap Y = (\bigcup_{i=1}^n (G_{x_i} \cap Y)) = \bigcup_{i=1}^n V_{x_i}$.


Since there are open V_{x_1}, \dots, V_{x_n} where $K \subset \bigcup_{i=1}^n V_{x_i}$ so K is compact relative to Y .

Suppose K is compact relative to Y .

Let open $G_1, G_2, \dots \subset X$ such that $X \subset \bigcup G_x$. For each G_x , let $V_x = Y \cap G_x \subset Y$.

Since K is compact relative to Y , there is a n such that $K \subset \bigcup_{i=1}^n V_{x_i}$.

Thus, $K \subset \bigcup_{i=1}^n V_{x_i} = \bigcup_{i=1}^n (Y \cap G_{x_i}) \subset \bigcup_{i=1}^n G_{x_i}$ so K is compact relative to X .




Theorem 6.3.4: A compact set is closed

Compact subsets of metric spaces are closed

Proof

Let compact $K \subset X$. Suppose $p \in X$, but $p \notin K$ so $p \in K^c$.
 If $q \in K$, let W_q be a neighborhood of q with $r < \frac{1}{2}d(p,q)$. Let $V_{p,q}$ be a neighborhood of p with $r < \frac{1}{2}d(p,q)$. Since K is compact, then there are finite points q_1, \dots, q_n such that $K \subset W$ where $W = W_{q_1} \cup \dots \cup W_{q_n}$.
 Let $V = V_{p,q_1} \cap \dots \cap V_{p,q_n}$, then $K \cap V \subset W \cap V = \emptyset$ so $V \subset K^c$.
 Since there is a neighborhood V for $p \in K^c$ where $V \subset K^c$, then every $p \in K^c$ is an interior point so K^c is open. Then by [theorem 5.1.7](#), K is closed.

**Theorem 6.3.5: If closed $E \subset$ compact set K , E is compact**

Closed subsets of compact sets are compact

Proof

Suppose $F \subset K \subset X$ where F is closed relative to X and K is compact.
 Let V_1, V_2, \dots be an open cover for F . Let open set F^c be all $k \in K$ where $k \notin F$.
 $K = F \cup F^c \subset V_1 \cup V_2 \cup \dots \cup F^c$
 Thus, $V_1 \cup V_2 \cup \dots \cup F^c$ is an open cover for K .
 Since K is compact, there is a finite subcover Ω that covers K and thus, finite subcover Ω covers $F \cup F^c$.
 Remove F^c from Ω . Since finite subcover $\Omega - F^c$ covers F , then F is compact.

Corollary 6.3.6: Closed $F \cap$ compact $K =$ compact

If F is closed and K is compact, then $F \cap K$ is compact

Proof

Since K is compact, then K is closed by [theorem 6.3.4](#).
 Then, by [5.1.9b](#), $F \cap K$ is closed.
 Since $F \cap K \subset K$, then by [theorem 6.3.5](#), $F \cap K$ is compact.

Theorem 6.3.7: Nonempty $\cap_{i=1}^n K_i \rightarrow$ nonempty $\cap K_i$

For compact sets $K_1, K_2, \dots \subset X$ where any intersection of finite K_i is nonempty, then $\cap K_i$ is nonempty

Proof

Fix K_1 . If there is a $k \in K_1$ where $k \in K_i$ for all i , then $k \in \cap K_i$ so $\cap K_i \neq \emptyset$.
 Suppose for every $k \in K_1$, $k \notin K_i$ for some i .
 Then for every $k \in K_1$, there is a K_i such that $k \notin K_i$ so $k \in K_i^c$.
 Thus, K_2^c, K_3^c, \dots form an open cover for K_1 . Since K_1 is compact, there is a n where $K_1 \subset K_{i_1}^c \cup \dots \cup K_{i_n}^c$. But then, $K_1 \cap K_{i_1} \cap \dots \cap K_{i_n} = \emptyset$ which is a contradiction.

Corollary 6.3.8: Nonempty K_i where $K_{i+1} \subset K_i \rightarrow$ nonempty $\cap K_i$

If K_1, K_2, \dots is a sequence of nonempty compact sets such that $K_{i+1} \subset K_i$, then $\cap K_i$ is nonempty

Proof

Since each K_i is nonempty and if $i_1 < \dots < i_n$, then $K_{i_1} \cap \dots \cap K_{i_n} = K_{i_n}$ is nonempty, then by **theorem 6.3.7**, $\cap K_i$ is nonempty.

Theorem 6.3.9: Nonempty intervals I_n where $I_{n+1} \subset I_n \rightarrow$ nonempty $\cap I_n$

If I_1, I_2, \dots is a sequence of intervals in \mathbb{R}^1 such that $I_{n+1} \subset I_n$, then $\cap I_n$ is nonempty.

Proof

Let $I_n = [a_n, b_n]$ and thus, each I_n is nonempty. If $n_1 < \dots < n_m$, then $I_{n_1} \cap \dots \cap I_{n_m} = [a_{n_m}, b_{n_m}]$ is nonempty. Thus, by **theorem 6.3.7**, $\cap I_n$ is nonempty.

Theorem 6.3.10: $p \in E'$ exists if infinite $E \subset$ compact K

If E is an infinite subset of compact set K , then E has a limit point in K

Proof

If no $p \in K$ is a $p \in E'$, then each p would have a neighborhood V_p contains at most $p \in E$ if $p \in E$. Thus, there is no finite subcover that covers E and thus, there is no finite subcover that covers K since $E \subset K$ which contradicts K is compact.

Definition 6.3.11: K-cells

The set of all $x = (x_1, \dots, x_k) \in \mathbb{R}^k$ where $x_i \in [a_i, b_i]$ for fixed $a_i, b_i \in \mathbb{R}$

Theorem 6.3.12: K-cells are compact

Every k-cell is compact

Proof

Let k-cell I consists of all $x = (x_1, \dots, x_k)$ where $x_i \in [a_i, b_i]$ for fixed $a_i, b_i \in \mathbb{R}$.
 Let $\delta = \sqrt{\sum_{i=1}^k (b_i - a_i)^2}$. Thus, $|x - y| \leq \delta$ for $x, y \in I$.
 Suppose there exists an open cover G_1, G_2, \dots of I which contain no finite subcover.
 Let $c_i = \frac{a_i + b_i}{2}$. Then each interval splits into $[a_i, c_i]$ and $[c_i, b_i]$ for $i \in [1, k]$ so there now exists 2^k k-cells Q_i whose union is I .
 At least one Q_i cannot be covered else I would be covered. Then subdivide Q_i as before and repeating the process so $Q_{i+1} \subset Q_i$ and each are not covered.
 However, there is a point $x^* \in Q_{i_j}$ for all j such that $N_r(x^*) \subset G$ so Q_{i_1} is covered which is a contradiction.

Theorem 6.3.13: Heine-Borel Theorem

If a set $E \subset \mathbb{R}^k$ has one of the three properties, then it has the other two:

- (a) E is closed and bounded
- (b) E is compact
- (c) Every infinite subset of E has a limit point in E

Proof

Suppose E is closed and bounded.

Then there exists a $M \in \mathbb{R}$ and $q \in \mathbb{R}^k$ such that $d(p, q) < M$ for all $p \in E$.

Thus, there is a k -cell $K = [-M + q_1, q_1 + M] \times \dots \times [-M + q_k, q_k + M]$ such that $E \subset K$. Then by [theorem 6.3.12](#), K is compact and thus by [theorem 6.3.5](#), E is compact so (a) \rightarrow (b).

Then by [theorem 6.3.10](#), any infinite subset of E has a limit point in E so (b) \rightarrow (c). Suppose E is not bounded.

Then there exists $p \in E$ such that $d(p, q) > M$ for any $M \in \mathbb{R}$ and $q \in \mathbb{R}^k$.

Let $S \subset E$ be such points p .

Then S is infinite else there is a maximal p and thus, p is bounded. Thus, S is infinite and contains no limit points in E since any $d(p_1, p_2) > M$ which contradicts that every infinite subset of E has a limit point in E . Thus, E is bounded.

Suppose E is not closed.

Then there exists a $p \in E'$, but $p \notin E$. Since p is a limit point, then there is a $q \in E$ such that $\frac{1}{n+1} < d(q, p) < \frac{1}{n}$ for $n = \{1, 2, \dots\}$.

Let $S \subset E$ be such points q .

Thus, p is the only limit point of S since for $r < \frac{1}{n}$, any $N_r(q_i)$ contains no points of S other than q_i since $d(q_i, q_j) > \frac{1}{n}$ for any $q_1, q_2 \in S$.

Thus, S is infinite, but the only $p \in S'$ is $p \notin E$ which contradicts that every infinite subset of E has a limit point in E . Thus, E is closed. So, (c) \rightarrow (a).

Theorem 6.3.14: Weierstrass Theorem

Every bounded infinite set $E \subset \mathbb{R}^k$ has a limit point in \mathbb{R}^k .

Proof

Since E is bounded, then there exists a k -cell K such that $E \subset K$. Since K is compact, then by [theorem 6.3.10](#), E has a limit point in K and thus, in \mathbb{R}^k .

7 Perfect and Connected Sets

7.1 Perfect Sets

Definition 7.1.1: Perfect Set

$E \subset X$ is perfect if E is closed and if every $p \in E$ is $p \in E'$

Theorem 7.1.2: Perfect sets are uncountable

Let P be a nonempty perfect set in \mathbb{R}^k . Then, P is uncountable.

Proof

Since P has limit points, then by [theorem 5.1.4](#), P is infinite.

Suppose P is countable. Then let $x_1, x_2, \dots \in P$.

Let V_i be a neighborhood of x_i where $y \in V_i$ for any $y \in \mathbb{R}^k$ such that $|y - x_i| < r$.

Thus, the $\overline{V_i}$ is the set of all $y \in \mathbb{R}^k$ such that $|y - x_i| \leq r$.


Since every x_i are limit points, then any $V_i \cap P$ is not empty where there is a V_{i+1}

(a) $\overline{V_{i+1}} \subset V_i$

(b) $x_i \notin \overline{V_{i+1}}$

(c) $V_{i+1} \cap P$ is nonempty

Let $K_i = \overline{V_i} \cap P$. Since $\overline{V_i}$ is closed and bounded, then by [theorem 6.3.11](#), $\overline{V_i}$ is compact. Since $x_i \notin K_{i+1}$, then no $x_i \in P$ is $x_i \in \cap K_i$. Since $K_n \subset P$, then $\cap K_i$ is empty which contradicts [corollary 6.3.8](#) since each K_i is nonempty and $K_{i+1} \subset K_i$.



Corollary 7.1.3: \mathbb{R} is not countable

Every interval $[a, b]$ is uncountable. Thus, \mathbb{R} is uncountable.

Proof

Since $[a, b]$ is closed and every $p \in [a, b]$ is a limit point, then nonempty set $[a, b]$ is perfect. Thus, by [theorem 7.1.2](#), $[a, b]$ is uncountable.

Definition 7.1.4: Cantor Set

There exists perfect segments in \mathbb{R}^1 which contain no segment.

Let $E_0 = [0,1]$.

For E_1 , remove $(\frac{1}{3}, \frac{2}{3})$. Thus, $E_1 = [0, \frac{1}{3}] \cup [\frac{2}{3}, 1]$.

For E_2 , remove $(\frac{1}{9}, \frac{2}{9})$ and $(\frac{7}{9}, \frac{8}{9})$. Thus, $E_2 = [0, \frac{1}{9}] \cup [\frac{2}{9}, \frac{3}{9}] \cup [\frac{6}{9}, \frac{7}{9}] \cup [\frac{8}{9}, 1]$.

Continuing such a sequence, the set of compact sets E_n are such that:

(a) $E_{n+1} \subset E_n$

(b) E_n is the union of 2^n intervals each of length 3^{-n} .

$P = \cap E_n$ is called the Cantor set. P is compact and nonempty.

Thus, any segment of form $(\frac{3k+1}{3^m}, \frac{3k+2}{3^m})$ where $k, m \in \mathbb{Z}_+$ has no points in common with P . Since any segment (a, b) contain a segment of such a form since $3^{-m} < \frac{b-a}{6}$, then P contains no segment.

Let $x \in P$ and segment S contain x . Let I_n be an interval of E_n containing x . Then choose a large enough n so $I_n \subset S$.

Let x_n be an endpoint of I_n where $x_n \neq x$ and thus, x is a limit point. Since P is closed and every $p \in P$ is $p \in P'$, then P is perfect.

7.2 Connected Sets**Definition 7.2.1: Connected Set**

$A, B \subset X$ are separated if both $A \cap \overline{B}$ and $\overline{A} \cap B$ are empty.

$E \subset X$ is connected if E is not the union of two nonempty separated sets.

Separated sets are disjoint, but disjoint sets need not be separated.

Theorem 7.2.2: All points between points in connected sets exists

$E \subset \mathbb{R}^1$ is connected if and only if:

If $x, y \in E$ and $x < z < y$, then $z \in E$.

Proof

If there exists $x, y \in E$ and $z \in (x, y)$ such that $z \notin E$, then $E = A_z \cup B_z$ where $A_z = E \cap (-\infty, z)$ and $B_z = E \cap (z, \infty)$.

Since $x \in A_z$ and $y \in B_z$, then A and B are nonempty. Since $A_z \subset (-\infty, z)$ and $B_z \subset (z, \infty)$, then A_z and B_z are separated. Thus, E is not connected.

Suppose E is not connected. Then, there are nonempty separated sets A and B such that $A \cup B = E$. Pick $x \in A$, $y \in B$ where $x < y$. Let $z = \sup(A \cap [x, y])$.

Since, $z \in \overline{A}$ so $z \notin B$, then $x \leq z < y$. If $z \notin A$, then $x < z < y$ so $z \notin E$.

If $z \in A$, then $z \notin \overline{B}$ and thus, there exists a z_1 such that $z < z_1 < y$ and $z_1 \notin B$. Then, $x < z_1 < y$ so $z_1 \notin E$.

8 Convergent and Cauchy Sequences

8.1 Convergent Sequences


Definition 8.1.1: Convergent Sequence

A sequence $\{x_n\}$ in metric space X converge if there is a $x \in X$ such that:

For every $\epsilon > 0$, there is a $N \in \mathbb{Z}$ such that for all $n \geq N$, $d(x_n, x) < \epsilon$

Then, $\{x_n\}$ converges to x : $\lim_{n \rightarrow \infty} x_n = x$

If $\{x_n\}$ does not converge, then it diverges.



Example

- (a) Let $x_n = \frac{1}{n}$ in \mathbb{R}^2 . Then, $\lim_{n \rightarrow \infty} x_n = 0$

Proof

For $\epsilon > 0$, there is a $\frac{1}{N} < \epsilon$. Then:

$$d(x_n, 0) = |x_n - 0| = \frac{1}{n} < \frac{1}{N} < \epsilon$$

- (b) Let $x_n = (-1)^n + \frac{1}{n}$ in \mathbb{R}^2 . Then, $\{x_n\}$ diverges.

Proof

$$\lim_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} (-1)^n + \lim_{n \rightarrow \infty} \frac{1}{n} = \lim_{n \rightarrow \infty} (-1)^n$$

Since $(-1)^n$ alternates between -1 and 1, then $\{x_n\}$ diverges.

Theorem 8.1.2: A convergent sequence is unique and bounded

- (a) $\{p_n\}$ converges to $p \in X$ if and only if every $N_r(p)$ contains p_n for all, but finitely many n .

Proof

Suppose $p_n \rightarrow p$. Then for $N_\epsilon(p)$, any $q \in X$ such that $d(q, p) < \epsilon$ is $q \in N_\epsilon(p)$. Since $p_n \rightarrow p$, there is a N such that for $n \geq N$, $d(p_n, p) < \epsilon$.

Thus, for $n \geq N$, $p_n \in N_\epsilon(p)$.

Suppose every $N_r(p)$ contains p_n for all, but finitely many n .

For $\epsilon > 0$, let $N_\epsilon(p)$ be the set of all $q \in X$ such that $d(p, q) < \epsilon$. Thus, there exists a N such that $p_n \in N_\epsilon(p)$ if $n \geq N$.

Thus, $d(p_n, p) < \epsilon$ so $p_n \rightarrow p$.

- (b) If $p, p' \in X$ and $\{p_n\}$ converges to p and p' , then $p = p'$.

Proof

For $\epsilon > 0$, there exists N, N' such that:

$$d(p_n, p) < \frac{\epsilon}{2} \text{ for } n \geq N \quad d(p_n, p') < \frac{\epsilon}{2} \text{ for } n \geq N'$$

Then for $n \geq \max(N, N')$, $d(p, p') \leq d(p, p_n) + d(p_n, p') < \epsilon$.

Thus, $p = p'$.

- (c) If
- $\{p_n\}$
- converges, then
- $\{p_n\}$
- is bounded.

Proof

If $\{p_n\} \rightarrow p$, there is a N such that for $n > N$, $d(p_n, p) < 1$.
 Let $r = \max(d(p_1, p), \dots, d(p_N, p), 1)$. Thus for all n , $d(p_n, p) \leq r$.

- (d) If
- $E \subset X$
- and
- $p \in E'$
- , there is a
- $\{p_n\}$
- in
- E
- such that
- $p = \lim_{n \rightarrow \infty} p_n$
- .

Proof

Since $p \in E'$, then for each $n \in \mathbb{Z}_+$, there is a $p_n \in E$ such that $d(p_n, p) < \frac{1}{n}$.
 For $\epsilon > 0$, there is a $\frac{1}{N} < \epsilon$ so for $n \geq N$, $d(p_n, p) < \frac{1}{n} \leq \frac{1}{N} < \epsilon$.
 Thus, $p = \lim_{n \rightarrow \infty} p_n$.

Theorem 8.1.3: Arithmetic Operations for sequences

Suppose $\{s_n\}, \{t_n\} \in \mathbb{C}$ where $\lim_{n \rightarrow \infty} s_n = s$ and $\lim_{n \rightarrow \infty} t_n = t$.

- (a)
- $\lim_{n \rightarrow \infty} s_n + t_n = s + t$

Proof

For $\epsilon > 0$, there exists N_1, N_2 such that
 $|s_n - s| < \frac{\epsilon}{2}$ for $n \geq N_1$ $|t_n - t| < \frac{\epsilon}{2}$ for $n \geq N_2$
 If $N = \max(N_1, N_2)$, then for $n \geq N$:
 $|s_n + t_n - s + t| \leq |s_n - s| + |t_n - t| < \epsilon$

- (b)
- $\lim_{n \rightarrow \infty} cs_n = cs$
- and
- $\lim_{n \rightarrow \infty} c + s_n = c + s$

Proof

For $\epsilon > 0$, there exists a N such that
 $|s_n - s| < \frac{\epsilon}{|c|}$ for $n \geq N$
 $|cs_n - cs| \leq |c| \cdot |s_n - s| < \epsilon$

- (c)
- $\lim_{n \rightarrow \infty} s_n t_n = st$

Proof

Note $s_n t_n - st = (s_n - s)(t_n - t) + t(s_n - s) + s(t_n - t)$.
 For $\epsilon > 0$, there exists N_1, N_2 such that
 $|s_n - s| < \sqrt{\epsilon}$ for $n \geq N_1$ $|t_n - t| < \sqrt{\epsilon}$ for $n \geq N_2$
 If $N = \max(N_1, N_2)$, then for $n \geq N$, $|(s_n - s)(t_n - t)| < \epsilon$.
 Thus, $\lim_{n \rightarrow \infty} (s_n - s)(t_n - t) = 0$.
 $\lim_{n \rightarrow \infty} (s_n t_n - st) = \lim_{n \rightarrow \infty} (s_n - s)(t_n - t) + t(s_n - s) + s(t_n - t)$
 $= 0 + t \cdot 0 + s \cdot 0 = 0$

- (d)
- $\lim_{n \rightarrow \infty} \frac{1}{s_n} = \frac{1}{s}$
- where
- $s_n, s \neq 0$

Proof

Choose m such that $|s_n - s| < \frac{1}{2}|s|$ if $n \geq m$ so $|s_n| > \frac{1}{2}|s|$ for $n \geq m$.
 For $\epsilon > 0$, there is a $N > m$ such that for $n \geq N$, $|s_n - s| < \frac{1}{2}|s|^2\epsilon$.
 Thus, for $n \geq N$, $|\frac{1}{s_n} - \frac{1}{s}| = \frac{|s_n - s|}{|s_n s|} < \frac{2}{|s|^2}|s_n - s| < \epsilon$.

Theorem 8.1.4: Extension to \mathbb{R}^k

- (a) Suppose $x_n \in \mathbb{R}^k$ and $x_n = (\alpha_{n_1}, \dots, \alpha_{n_k})$. Then $\{x_n\}$ converges to $x = (\alpha_1, \dots, \alpha_k)$ if and only if $\lim_{n \rightarrow \infty} \alpha_{n_i} = \alpha_i$ for $i \in [1, k]$.

Proof

Suppose $\{x_n\}$ converges to $x = (\alpha_1, \dots, \alpha_k)$.

Since for any $i \in [1, k]$:

$$|\alpha_{n_i} - \alpha_i| \leq \sqrt{|\alpha_{n_1} - \alpha_1|^2 + \dots + |\alpha_{n_k} - \alpha_k|^2} = |x_n - x| < \epsilon.$$

Then, $\lim_{n \rightarrow \infty} \alpha_{n_i} = \alpha_i$.

Suppose $\lim_{n \rightarrow \infty} \alpha_{n_i} = \alpha_i$ for $i \in [1, k]$.

Then for $\epsilon > 0$, there is an N such that for $n \geq N$:

$$|\alpha_{n_i} - \alpha_i| < \frac{\epsilon}{\sqrt{k}} \text{ for } i \in [1, k]$$

$$|x_n - x| = \sqrt{\sum_{i=1}^k |\alpha_{n_i} - \alpha_i|^2} < \sqrt{k \cdot \left(\frac{\epsilon}{\sqrt{k}}\right)^2} = \epsilon$$

- (b) Suppose $\{x_n\}, \{y_n\} \in \mathbb{R}^k$ and $\{\beta_n\} \in \mathbb{R}$ and $x_n \rightarrow x, y_n \rightarrow y, \beta_n \rightarrow \beta$.
 $\lim_{n \rightarrow \infty} x_n + y_n = x + y \quad \lim_{n \rightarrow \infty} x_n \cdot y_n = x \cdot y \quad \lim_{n \rightarrow \infty} \beta_n x_n = \beta x$

Proof

By part a, then $\lim_{n \rightarrow \infty} x_{n_i} + y_{n_i} = x_i + y_i$ so $\{x_n + y_n\} \rightarrow x + y$.

Also, $\lim_{n \rightarrow \infty} \sum_{i=1}^k x_{n_i} y_{n_i} = \sum_{i=1}^k x_i y_i$ so $\{x_n \cdot y_n\} \rightarrow x \cdot y$.

Also, $\lim_{n \rightarrow \infty} \beta_i x_{n_i} = \beta_i x_i$ so $\{\beta_n x_n\} \rightarrow \beta x$.

8.2 Subsequences

Definition 8.2.1: Subsequence

For sequence $\{p_n\}$, let $\{n_k\} \in \mathbb{Z}_+$ where $n_k < n_{k+1}$.

Then $\{p_{n_k}\}$ is a subsequence of $\{p_n\}$.

If $\{p_{n_k}\}$ converges, then its limit is called a subsequential limit.

Theorem 8.2.2: $\{p_n\} \rightarrow p \iff \text{Every } \{p_{n_k}\} \rightarrow p$

$\{p_n\}$ converges to p if and only if every subsequence converges to p

Proof


Suppose $\{p_n\}$ converges to p .

Then for $\epsilon > 0$, there is a N such that for $n \geq N$, $d(p_n, p) < \epsilon$.

Let $\{p_{n_k}\} \subset \{p_n\}$. Then for $n_k \geq N$, $|p_{n_k} - p| < \epsilon$. Thus, $\{p_{n_k}\} \rightarrow p$.

Suppose every subsequence converges to p .

Since $\{p_n\}$ is a subsequence of itself, then $\{p_n\}$ converges to p .



Theorem 8.2.3: $\{p_n\}$ in compact space have $\{p_{n_k}\} \rightarrow p$

- (a) If $\{p_n\}$ is a sequence in a compact metric space X , then some subsequence converges to $p \in X$.

Proof

Let E be the range of $\{p_n\}$.

If E is finite, there is a $p \in E$ and sequence $\{n_k\}$ with $n_k < n_{k+1}$ such that $p_{n_1} = p_{n_2} = \dots = p$. Thus, $\{p_{n_k}\} \rightarrow p$.

If E is infinite, then by **theorem 6.3.10**, then there exists a $p \in E'$.

Then there are n_k such that $d(p_{n_k}, p) < \frac{1}{k}$. Thus, $\{p_{n_k}\} \rightarrow p$.

- (b) Every bounded sequence in \mathbb{R}^k contains a convergent subsequence.

Proof

Let E be a bounded sequence in \mathbb{R}^k . Since $E \cup E'$ is bounded and closed, then by **theorem 6.3.13**, $E \cup E'$ is compact.

Thus by part a, E contains a convergent subsequence.

Theorem 8.2.4: The set of subsequential limits is closed

The subsequential limits of $\{p_n\}$ in metric space X form a closed subset of X

Proof

Let E be the range of the set of all subsequential limits of $\{p_n\}$.

If E is empty, then E is closed. If E is finite, then E' is empty so E is closed.

Suppose E is infinite. Then, let $q \in E'$.

Since $q \in E'$, there is a $x \in E$ where $d(x, q) < \frac{\epsilon}{2}$.

Since $x \in E$, there is a $\{p_{n_k}\} \rightarrow x$ so there is a N such that for $n \geq N$, $d(p_{n_k}, x) < \frac{\epsilon}{2}$.

Thus, $d(p_{n_k}, q) \leq d(p_{n_k}, x) + d(x, q) < \epsilon$ so q is a subsequential limit of $\{p_n\}$.

Thus, $q \in E$ so E is closed.



8.3 Cauchy Sequences

Definition 8.3.1: Metric Spaces

Sequence $\{p_n\} \in X$ is a Cauchy sequence if:

For every $\epsilon > 0$, there is a $N \in \mathbb{Z}$ such that for all $n, m \geq N$, $d(p_n, p_m) < \epsilon$

Let nonempty $E \subset X$ and $S \subset \mathbb{R}$ of $d(p, q)$ where $p, q \in E$.

Let $\sup(S) = \text{diam}(E)$. If $\{p_n\} \in X$, and $p_N, p_{N+1}, \dots \in E_N$, then $\{p_n\}$ is a Cauchy sequence if and only if $\lim_{N \rightarrow \infty} \text{diam}(E_N) = 0$.

Theorem 8.3.2: Cauchy sequences and its closure have the same diam

- (a) If
- $\overline{E} \subset X$
- , then
- $\text{diam}(\overline{E}) = \text{diam}(E)$
- .

Proof

Since $E \subset \overline{E}$, then $\text{diam}(E) \leq \text{diam}(\overline{E})$.

For $\epsilon > 0$, let $p, q \in E$.

Thus, there are $p', q' \in E$ such that $d(p', p) < \epsilon$ and $d(q', q) < \epsilon$. Thus:
 $d(p, q) \leq d(p, p') + d(p', q') + d(q', q) < 2\epsilon + d(p', q') \leq 2\epsilon + \text{diam}(E)$.

Thus, $\text{diam}(\overline{E}) \leq 2\epsilon + \text{diam}(E)$ so $\text{diam}(\overline{E}) = \text{diam}(E)$.

- (b) If
- K_n
- is a sequence of compact sets of
- X
- such that
- $K_{n+1} \subset K_n$
- and
- $\lim_{n \rightarrow \infty} \text{diam}(K_n) = 0$
- , then
- $\bigcap K_n$
- consist of only one point.

Proof

Let $K = \bigcap K_n$. Since K_n is a sequence of compact sets, then by [corollary 6.3.8](#), K is nonempty.

If K contains more than one point, then $\text{diam}(K) > 0$. But since $K \subset K_n$, then $\text{diam}(K) \leq \text{diam}(K_n)$ which contradicts that $\text{diam}(K_n) \rightarrow 0$.

Theorem 8.3.3: Convergent sequences are cauchy sequences

- (a) Every convergent sequence is a Cauchy sequence.

Proof

If $p_n \rightarrow p$ and $\epsilon > 0$, there is a N such that for all $n \geq N$, $d(p, p_n) < \frac{\epsilon}{2}$. Thus, for $m, n \geq N$:

$$d(p_n, p_m) \leq d(p_n, p) + d(p, p_m) < \epsilon.$$

Thus, $\{p_n\}$ is a Cauchy sequence.

- (b) If
- $\{p_n\}$
- is a Cauchy sequence in compact metric space
- X
- , then
- $\{p_n\}$
- converges to some
- $p \in X$
- .

Proof

Let $\{p_n\}$ be a Cauchy sequence in compact space X .

Let $p_N, p_{N+1}, \dots \in E_N$.

Since $\{p_n\}$ is a Cauchy sequence, then $\lim_{N \rightarrow \infty} \text{diam}(\overline{E_N}) = 0$. Since $\overline{E_N}$ is closed in compact X , then by [theorem 6.3.5](#), $\overline{E_N}$ is compact.

Since $E_{N+1} \subset E_N$, then $\overline{E_{N+1}} \subset \overline{E_N}$ and thus, by [theorem 8.3.2b](#), then there is a unique $p \in \overline{E_N}$ for every N .

Since $p \in \overline{E_N}$, then $d(p, q) < \epsilon$ for every $q \in \overline{E_N}$ so every $q \in E_N$.

Then for $\epsilon > 0$, there is a N_0 such that for $N \geq N_0$, $\text{diam}(\overline{E_N}) < \epsilon$.

Thus, $d(p_n, p) < \epsilon$ for $n \geq N_0$ so $\{p_n\} \rightarrow p$.

- (c) In
- \mathbb{R}^k
- , every Cauchy sequence converges.

Proof

Let $\{x_n\}$ be a Cauchy sequence in \mathbb{R}^k . Let $x_N, x_{N+1}, \dots \in E_N$.

Then for some N , $\text{diam}(E_N) < 1$. Thus, the range of $\{x_n\} = E_N \cup \{x_1, \dots, x_{N-1}\}$. Thus, $\{x_n\}$ is bounded.

Thus, the $\{x_n\}$ is closed and bounded so by [theorem 6.3.13](#), $\overline{\{x_n\}}$ is compact.

Thus, by part b, $\{x_n\}$ converges to some $p \in \mathbb{R}^k$.

Definition 8.3.4: Complete

A metric space where every Cauchy sequence converges is complete.

Thus, by **theorem 8.3.3**, all compact and Euclidean spaces are complete.

Definition 8.3.5: Monotonic Sequences

A sequence $\{s_n\}$ of real numbers is:

- (a) monotonically increasing if $s_n \leq s_{n+1}$
- (b) monotonically decreasing if $s_n \geq s_{n+1}$

Theorem 8.3.6: Monotonic sequences converge if bounded

Suppose $\{s_n\}$ is monotonic. Then $\{s_n\}$ converges if and only if it is bounded

Proof

Suppose $s_n \leq s_{n+1}$. Let E be the range of $\{s_n\}$.

Suppose $\{s_n\}$ is bounded.

Let $s = \sup(E)$ so $s_n \leq s$. For every $\epsilon > 0$, there is a N such that $s - \epsilon < s_N \leq s$ else $s - \epsilon$ would be an upper bound of E which contradicts $s = \sup(E)$.

Since $\{s_n\}$ increases, then for $n \geq N$, $s - \epsilon < s_N \leq s_n \leq s$ so $\{s_n\} \rightarrow s$.

Suppose $\{s_n\}$ converges to s .

Then for $\epsilon > 0$, there is a N such that for $n \geq N$, $s - \epsilon < s_N \leq s_n \leq s$.

Thus, $\{s_n\}$ is bounded from above.

Suppose $s_n \geq s_{n+1}$. Let E be the range of $\{s_n\}$.

Suppose $\{s_n\}$ is bounded.

Let $s = \inf(E)$ so $s_n \geq s$. For every $\epsilon > 0$, there is a N such that $s \leq s_N < s + \epsilon$ else $s + \epsilon$ would be a lower bound of E which contradicts $s = \inf(E)$.

Since $\{s_n\}$ decreases, then for $n \geq N$, $s \leq s_n \leq s_N < s + \epsilon$ so $\{s_n\} \rightarrow s$.

Suppose $\{s_n\}$ converges to s .

Then for $\epsilon > 0$, there is a N such that for $n \geq N$, $s \leq s_n \leq s_N < s + \epsilon$.

Thus, $\{s_n\}$ is bounded from below.

9 Limits and Special Sequences

9.1 Upper and Lower Limits

Definition 9.1.1: Infinite limits

Let $\{s_n\}$ be a sequence of real numbers such that:

For every real M , there is a $N \in \mathbb{Z}$ such that for $n \geq N$, $s_n \geq M$.

Then, $s_n \rightarrow +\infty$.

For every real M , there is a $N \in \mathbb{Z}$ such that for $n \geq N$, $s_n \leq M$.

Then, $s_n \rightarrow -\infty$.

Definition 9.1.2: Upper and Lower Limits


Let $\{s_n\} \subset \mathbb{R}$ and E contain all subsequential limits of $\{s_n\}$ plus possibly $\pm\infty$.

Then, the upper limit of $\{s_n\}$:

$$s^* = \sup(E) \quad \lim_{n \rightarrow \infty} \sup(s_n) = s^*$$

Then, the lower limit of $\{s_n\}$:

$$s_* = \inf(E) \quad \lim_{n \rightarrow \infty} \inf(s_n) = s_*$$



Theorem 9.1.3: Upper and Lower limits are unique

Let $\{s_n\}$ be a sequence of real numbers. Let E be the set of subsequential limits and s^* be the upper limit of $\{s_n\}$. Then:

(a) $s^* \in E$

Proof

If $s^* = +\infty$, then there is a $\{s_{n_k}\} \rightarrow +\infty$ so E is not bounded above.

If $s^* \in \mathbb{R}$, then E is bounded above so $s^* \in E'$.

Then by [theorem 8.2.4](#), $s^* \in E$.

If $s^* = -\infty$, then there are no subsequential limits in E . Thus, for every M , there is a N such that for $n \geq N$, $s_n \leq M$ so $-\infty \in E$.

(b) If $x > s^*$, there is a N such that for $n \geq N$, $s_n < x$

Proof

Suppose there is a $x > s^*$ such that $s_n \geq x$ for infinitely many n .

Then, there is a $y \in E$ where $y \geq x > s^*$ which contradicts $s^* = \sup(E)$.

(c) s^* is the only number that satisfies (a) and (b)

Proof

Suppose p, q satisfy part a and b where $p < q$. Choose x where $p < x < q$. Since p satisfies b, then $s_n < x$ for $n \geq N$. Thus, x is an upper bound for E so $q \notin E$ since $q > x$ contradicting that q satisfies part a.

The same properties are analogous for s_* .

Theorem 9.1.4: Inf & Sup of $s_n \leq t_n$

If $s_n \leq t_n$ for $n \geq$ fixed N , then

$$\lim_{n \rightarrow \infty} \inf(s_n) \leq \lim_{n \rightarrow \infty} \inf(t_n)$$

$$\lim_{n \rightarrow \infty} \sup(s_n) \leq \lim_{n \rightarrow \infty} \sup(t_n)$$

Proof

Let E_1 be the set of extended reals x such that $\{s_{n_k}\} \rightarrow x$ for some $\{s_{n_k}\}$.

Let E_2 be the set of extended reals y such that $\{t_{n_k}\} \rightarrow y$ for some $\{s_{n_k}\}$.

Let $s^* = \sup(E_1)$, $s_* = \inf(E_1)$, $t^* = \sup(E_2)$, and $t_* = \inf(E_2)$.

Since there is a N such that $s_n \leq t_n$ for $n \geq N$, then:

$$x \leftarrow \{s_N, s_{N+1}, \dots\} \leq \{t_N, t_{N+1}, \dots\} \rightarrow y$$

Thus, for $n \geq N$, $\inf(s_n) \leq \inf(t_n)$ and $\sup(s_n) \leq \sup(t_n)$.

9.2 Special Sequences**Theorem 9.2.1: Special sequences**

- (a) If $p > 0$, then $\lim_{n \rightarrow \infty} \frac{1}{n^p} = 0$

Proof

For $\epsilon > 0$, let $N > \sqrt[p]{\frac{1}{\epsilon}}$. Then for $n \geq N$, $\lim_{n \rightarrow \infty} \frac{1}{n^p} \leq \frac{1}{N^p} < \frac{1}{\sqrt[p]{\frac{1}{\epsilon}}} = \epsilon$

- (b) If $p > 0$, then $\lim_{n \rightarrow \infty} \sqrt[p]{p} = 1$

Proof

If $p > 1$, then let $x_n = \sqrt[p]{p} - 1 > 0$.

$$p = (x_n + 1)^n = x_n^n + nx_n^{n-1} + \dots + nx_n + 1 \geq nx_n + 1$$

Thus, $0 < x_n \leq \frac{p-1}{n}$ so $\{x_n\} \rightarrow 0$ and thus, $\{\sqrt[p]{p}\} \rightarrow 1$.

If $p = 1$, then $\lim_{n \rightarrow \infty} \sqrt[p]{p} = \lim_{n \rightarrow \infty} 1 = 1$.

If $0 < p < 1$, then $\frac{1}{p} > 1$. From the proof above for $p > 1$, $\{\sqrt[\frac{1}{p}]{\frac{1}{p}}\} \rightarrow 1$.

Thus, $\{\frac{1}{\sqrt[p]{p}}\} \rightarrow 1$ so $\{\sqrt[p]{p}\} \rightarrow 1$.

- (c) $\lim_{n \rightarrow \infty} \sqrt[n]{n} = 1$

Proof

Let $x_n = \sqrt[n]{n} - 1 \geq 0$. Then, $n = (x_n + 1)^n \geq \frac{n(n-1)}{2} x_n^2$.

Thus, $0 \leq x_n \leq \sqrt{\frac{2}{n-1}}$ so $\{x_n\} \rightarrow 0$ and thus, $\{\sqrt[n]{n}\} \rightarrow 1$.

- (d) If $p > 0$ and $\alpha \in \mathbb{R}$, then $\lim_{n \rightarrow \infty} \frac{n^\alpha}{(1+p)^n} = 0$

Proof

Let $k \in \mathbb{Z}$ such that $k > \alpha$ and $k > 0$. For $n > 2k$:

$$(1+p)^n > \binom{n}{k} p^k = \frac{n(n-1)\dots(n-k+1)}{k!} p^k > \frac{n^k p^k}{2^k k!}$$

Thus, $0 < \frac{n^\alpha}{(1+p)^n} < \frac{2^k k!}{p^k} n^{\alpha-k}$.

Since $\alpha - k < 0$, then $\{n^{\alpha-k}\} \rightarrow 0$ so $\{\frac{n^\alpha}{(1+p)^n}\} \rightarrow 0$.

- (e) If $|x| < 1$, then $\lim_{n \rightarrow \infty} x^n = 0$

Proof

From part d, let $\alpha = 0$.

Thus, $\lim_{n \rightarrow \infty} \frac{1}{(1+p)^n} = 0$ and since $p > 0$, then $\frac{1}{(1+p)^n} = (\frac{1}{1+p})^n < 1$.

Also, $-\lim_{n \rightarrow \infty} \frac{1}{(1+p)^n} = \lim_{n \rightarrow \infty} \frac{-1}{(1+p)^n} = 0$ so $\frac{-1}{(1+p)^n} = (\frac{-1}{1+p})^n > -1$.

10 Series and Convergence Tests

10.1 Series

Definition 10.1.1: Series

For sequence $\{a_n\}$, define $\sum_{n=p}^q a_n = a_p + a_{p+1} + \dots + a_q$.

Then associate $\{a_n\}$ with a sequence $\{s_n\}$ such that $s_n = \sum_{k=1}^n a_k$.

Then $\{s_n\}$ is a series with partial sums s_n .

If $\{s_n\} \rightarrow s$, then $\sum_{n=1}^{\infty} a_n = s$ is the sum of the convergent series.

Note $a_1 = s_1$ and $a_n = s_n - s_{n-1}$.

Theorem 10.1.2: Cauchy Criterion for series

$\sum a_n$ converges if and only if:

For every $\epsilon > 0$, there is a $N \in \mathbb{Z}$ such that for $m \geq n \geq N$, $|\sum_{k=n}^m a_k| \leq \epsilon$

Proof

Suppose $\sum_{k=1}^n a_k$ converges.

Then by [theorem 8.3.3a](#), $\sum_{k=1}^n a_k$ is a Cauchy sequence.

Then for $\epsilon > 0$, there is a N such that for $m \geq n \geq N$:

$$d(\sum_{k=1}^n a_k, \sum_{k=1}^m a_k) = |\sum_{k=1}^m a_k - \sum_{k=1}^n a_k| = |\sum_{k=n}^m a_k| \leq \epsilon$$

Suppose for every $\epsilon > 0$, there is a N such that for $m \geq n \geq N$, $|\sum_{k=n}^m a_k| \leq \epsilon$.

$$|\sum_{k=n}^m a_k| = |\sum_{k=1}^m a_k - \sum_{k=1}^n a_k| = d(\sum_{k=1}^n a_k, \sum_{k=1}^m a_k) \leq \epsilon$$

Thus, $\sum_{k=1}^n a_k$ is a Cauchy sequence and thus, convergent.

Theorem 10.1.3: Convergent $\sum a_n \Rightarrow \{a_n\} \rightarrow 0$

If $\sum a_n$ converges, then $\lim_{n \rightarrow \infty} a_n = 0$.

Proof

Since $\sum a_n$ converges, then by [theorem 10.1.2](#), for $\epsilon > 0$, there is a N such that for $m \geq n \geq N$, $|\sum_{k=n}^m a_k| \leq \epsilon$. Then if $m = n \geq N$, $|\sum_{k=n}^m a_k| = |a_n| \leq \epsilon$ so $\{a_n\} \rightarrow 0$.

Example

$\sum_{n=1}^{\infty} \frac{1}{n}$ diverges

$$\sum_{n=1}^{\infty} \frac{1}{n} = 1 + \frac{1}{2} + (\frac{1}{3} + \frac{1}{4}) + (\frac{1}{5} + \dots + \frac{1}{8}) + (\frac{1}{9} + \dots) \geq 1 + \frac{1}{2} + \frac{1}{2} + \frac{1}{2} + \dots$$

Thus, $s_{2^k} = \sum_{n=1}^{2^k} a_n \geq 1 + k \cdot \frac{1}{2}$ which is unbounded and thus, not convergent.

Theorem 10.1.4: Convergent series \Leftrightarrow Bounded sequence

A series of nonnegative terms converge if and only if its partial sums form a bounded sequence.

Proof

Suppose $\sum a_n$ converges where $a_n \geq 0$.

Since $a_n \geq 0$, then $\{s_n\}$ is monotonic so by [theorem 8.3.6](#), $\{s_n\}$ is bounded above.

Suppose $\{s_n\}$ is bounded where $a_n \geq 0$.

Since $\{s_n\}$ is monotonic and bounded, then by [theorem 8.3.6](#), $\{s_n\}$ converges.

Theorem 10.1.5: Comparison Test

(a) If $|a_n| \leq c_n$ for $n \geq N_0$ and $\sum c_n$ converges, then $\sum a_n$ converges.

Proof

For $\epsilon > 0$, there exists a $N \geq N_0$ such that for $m \geq n \geq N$, $\sum_{k=n}^m c_k \leq \epsilon$.

$$\left| \sum_{k=n}^m a_k \right| \leq \sum_{k=n}^m |a_k| \leq \sum_{k=n}^m c_k \leq \epsilon$$
 Thus, $\sum a_n$ converges.

(b) If $a_n \geq d_n \geq 0$ for $n \geq N_0$ and $\sum d_n$ diverges, then $\sum a_n$ diverges.

Proof

Suppose $\sum a_n$ converges.
 Then from part a, $\sum d_n$ converges which contradicts that $\sum a_n$ diverges.
 Thus, $\sum a_n$ diverges.

10.2 Series of Nonnegative Terms**Theorem 10.2.1: Infinite Geometric Series**

If $x \in [0, 1)$, then:

$$\sum_{n=0}^{\infty} x^n = \frac{1}{1-x}$$

If $x \geq 1$, the series diverges.

Proof

If $x \neq 1$, then using the geometric series $s_n = \sum_{k=0}^n x^k = \frac{1-x^{n+1}}{1-x}$. Let $n \rightarrow \infty$.
 If $x \in [0, 1)$, then by **theorem 9.2.1e**, $s_n = \frac{1}{1-x} (1 - x^{n+1}) = \frac{1}{1-x} (1 - 0) = \frac{1}{1-x}$.
 Also, by **theorem 9.2.1e**, if $x \geq 1$, then the series diverges.

Theorem 10.2.2: Cauchy's Convergence Criterion

Suppose $0 \leq a_{i+1} \leq a_i$.

Then the series $\sum_{n=1}^{\infty} a_n$ converges if and only if the series

$$\sum_{k=0}^{\infty} 2^k a_{2^k} = a_1 + 2a_2 + 4a_4 + 8a_8 + \dots \text{ converges.}$$

Proof

Let $s_n = a_1 + a_2 + \dots + a_n$ and $t_k = a_1 + 2a_2 + \dots + 2^k a_{2^k}$. For $n < 2^k$:

$$\begin{aligned} s_n &\leq a_1 + a_2 + a_3 + a_4 + a_5 + a_6 + a_7 + \dots + a_{2^k} \\ &\leq a_1 + (a_2 + a_3) + (a_4 + a_5 + a_6 + a_7) + \dots + (a_{2^{k-1}} + \dots + a_{2^k-1}) \\ &\leq a_1 + 2a_2 + 4a_4 + \dots + 2^k a_{2^k} = t_k \end{aligned}$$

By **comparison test**, if $\sum_{k=0}^{\infty} 2^k a_{2^k}$ converges, then $\sum_{n=1}^{\infty} a_n$ converges. For $n > 2^k$:

$$\begin{aligned} s_n &\geq a_1 + a_2 + a_3 + a_4 + a_5 + a_6 + a_7 + a_8 + \dots + a_{2^k} \\ &= a_1 + a_2 + (a_3 + a_4) + (a_5 + a_6 + a_7 + a_8) + \dots + (a_{2^{k-1}+1} + \dots + a_{2^k}) \\ &\geq \frac{1}{2}a_1 + a_2 + 2a_4 + \dots + 2^{k-1}a_{2^k} = \frac{1}{2}t_k \end{aligned}$$

By **comparison test**, if $\sum_{n=1}^{\infty} a_n$ converges, then $\sum_{k=0}^{\infty} 2^k a_{2^k}$ converges.

Theorem 10.2.3: P-series

$\sum \frac{1}{n^p}$ converges if $p > 1$ and diverges if $p \leq 1$

Proof

If $p \leq 0$, then by **theorem 10.1.3**, $\sum \frac{1}{n^p}$ diverges.
 If $p > 0$, then by **theorem 10.2.2**, $\sum \frac{1}{n^p}$ converges only if $\sum_{k=0}^{\infty} 2^k \frac{1}{(2^k)^p}$ converges.
 Since $\sum_{k=0}^{\infty} 2^k \frac{1}{(2^k)^p} = \sum_{k=0}^{\infty} 2^{(1-p)k}$, then by **theorem 10.2.1**, $\sum_{k=0}^{\infty} 2^{k(1-p)}$ converges if $2^{1-p} < 1$ so if $1-p < 0$ so $p > 1$.

Theorem 10.2.4: Log P-series

$\sum_{n=2}^{\infty} \frac{1}{n(\log(n))^p}$ converges if $p > 1$ and diverges if $p \leq 1$.

Proof

Since $\frac{1}{n(\log(n))^p}$ decreases, then by **theorem 10.2.2**,
 $\sum_{n=0}^{\infty} \frac{1}{n(\log(n))^p}$ converges if $\sum_{k=1}^{\infty} 2^k \frac{1}{2^k \log(2^k)}$ converges.
 $\sum_{k=1}^{\infty} 2^k \frac{1}{2^k \log(2^k)} = \sum_{k=1}^{\infty} \frac{1}{k \log(2)} = \frac{1}{\log(2)} \sum_{k=1}^{\infty} \frac{1}{k}$
 Then by **theorem 10.2.3**, $\sum_{k=1}^{\infty} 2^k \frac{1}{2^k \log(2^k)}$ converges if $p > 1$ and diverges if $p \leq 1$.
 Thus, $\sum_{n=0}^{\infty} \frac{1}{n(\log(n))^p}$ converges if $p > 1$ and diverges and $p \leq 1$.

Corollary 10.2.5: Log P-series extended

$\sum_{n=3}^{\infty} \frac{1}{n \log(n)(\log(\log(n)))^p}$ converges if $p > 1$ and diverges if $p \leq 1$

Proof

From **theorem 10.2.4**, replace $n = \log(n)$ and multiplying by $\frac{1}{n} \rightarrow \frac{1}{n \log(n)(\log(\log(n)))^p}$.
 Since $\frac{1}{n \log(n)(\log(\log(n)))^p}$ decreases, by **theorem 10.2.2** $\sum_{k=1}^{\infty} 2^k \frac{1}{2^k \log(2^k)(\log(\log(2^k)))^p}$.
 $\sum_{k=1}^{\infty} \frac{1}{\log(2^k)(\log(\log(2^k)))^p} = \frac{1}{\log(2)} \sum_{k=1}^{\infty} \frac{1}{k(\log(k \log(2)))^p} < \frac{1}{\log(2)} \sum_{k=2}^{\infty} \frac{1}{k(\log(k))^p}$
 Since $\sum_{k=2}^{\infty} \frac{1}{k(\log(k))^p}$ converges by **theorem 10.2.4**, $\sum_{n=3}^{\infty} \frac{1}{n \log(n)(\log(\log(n)))^p}$ converges.

10.3 The Number e**Definition 10.3.1: Summation equivalence to e**

$$s_m = \sum_{n=0}^m \frac{1}{n!} = 1 + \sum_{n=1}^m \frac{1}{n!} < 1 + \sum_{n=1}^m \frac{1}{2^{n-1}} < 3$$

$$e = \sum_{n=0}^{\infty} \frac{1}{n!}$$

Theorem 10.3.2: Limit equivalence to e

$$\lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n = e$$

Proof

Let $s_n = \sum_{k=0}^n \frac{1}{k!}$ and $t_n = \left(1 + \frac{1}{n}\right)^n$. Using the binomial theorem:
 $t_n = \sum_{k=0}^n \binom{n}{k} \frac{1}{n^k} = \sum_{k=0}^n \frac{n(n-1)\dots(n-k+1)}{k!} \frac{1}{n^k} = \sum_{k=0}^n \frac{1}{k!} \left(1 - \frac{1}{n}\right)\left(1 - \frac{2}{n}\right)\dots\left(1 - \frac{k-1}{n}\right)$
 Thus, $t_n \leq s_n$ so $\lim_{n \rightarrow \infty} \sup(t_n) \leq e$.
 If $n \geq m$, then $t_n \geq \sum_{k=0}^m \frac{1}{k!} \left(1 - \frac{1}{n}\right)\left(1 - \frac{2}{n}\right)\dots\left(1 - \frac{k-1}{n}\right)$.
 As $n \rightarrow \infty$, then $\lim_{n \rightarrow \infty} \inf(t_n) \geq \sum_{k=0}^m \frac{1}{k!} = s_m$. As $m \rightarrow \infty$, $\lim_{n \rightarrow \infty} \inf(t_n) \geq e$.

Theorem 10.3.3: Rapidity of convergence of e

$$0 < e - s_n < \frac{1}{n!n}$$

Proof

$$e - s_n = \sum_{k=n+1}^{\infty} \frac{1}{k!} < \frac{1}{(n+1)!} \left(1 + \frac{1}{n+1} + \frac{1}{(n+1)^2} + \dots\right) = \frac{1}{(n+1)!} \frac{1}{1 - \frac{1}{n+1}} = \frac{1}{n!n}$$

Theorem 10.3.4: e is irrational

e is irrational

Proof

Suppose r is rational. Then let $e = \frac{p}{q}$ for $p, q \in \mathbb{Z}_+$.
 Thus, by **theorem 10.3.3**, $0 < e - s_q < \frac{1}{q!q}$ so $0 < q!(e - s_q) < \frac{1}{q}$.
 Since $e = \frac{p}{q}$, then $q!e$ is an integer and $q!s_q = q!(1 + \frac{1}{2!} + \dots + \frac{1}{q!})$ is an integer.
 Thus, $q!(e - s_q)$ is an integer which is between 0 and $\frac{1}{q}$ and thus, a contradiction.

10.4 Root and Ratio Tests**Theorem 10.4.1: Root Test**

- For $\sum a_n$, let $\alpha = \lim_{n \rightarrow \infty} \sup(\sqrt[n]{|a_n|})$.
- (a) If $\alpha < 1$, $\sum a_n$ converges
 - (b) If $\alpha > 1$, $\sum a_n$ diverges
 - (c) If $\alpha = 1$, unclear

Proof

If $\alpha < 1$, choose β such that $\beta \in (\alpha, 1)$ and $N \in \mathbb{Z}$ such that $\sqrt[n]{|a_n|} < \beta$ for $n \geq N$.
 Since $\beta \in (0, 1)$, then by **theorem 10.2.1**, $\sum \beta^n$ converges. Then by the **comparison test**, $\sum a_n$ converges.
 If $\alpha > 1$, then there is a a_{n_k} such that $\sqrt[n_k]{|a_{n_k}|} \rightarrow \alpha$.
 Thus, $|a_n| > 1$ for infinitely many n so by **theorem 10.1.3**, $\sum a_n$ doesn't converge.
 $\sum \frac{1}{n}$, $\sum \frac{1}{n^2}$ have $\alpha = 1$, but $\sum \frac{1}{n}$ diverges and $\sum \frac{1}{n^2}$ converges by **theorem 10.2.3**.

Theorem 10.4.2: Ratio Test

- (a) $\sum a_n$ converges if $\lim_{n \rightarrow \infty} \sup(|\frac{a_{n+1}}{a_n}|) < 1$
- (b) $\sum a_n$ diverges if $|\frac{a_{n+1}}{a_n}| \geq 1$ for all $n \geq n_0$ for $n_0 \in \mathbb{Z}$

Proof

If $\lim_{n \rightarrow \infty} \sup(|\frac{a_{n+1}}{a_n}|) < 1$, there is a $\beta < 1$ and N such that for $n \geq N$, $|\frac{a_{n+1}}{a_n}| < \beta$.
 Then $|a_{N+1}| < \beta|a_N|$ so $|a_{N+2}| < \beta|a_{N+1}| < \beta^2|a_N|$.
 Thus, $|a_{N+p}| < \beta^p|a_N|$ so $|a_n| < |a_N|\beta^{-N}\beta^n$.
 Thus, by the **comparison test**, $\sum a_n$ converges.
 If $|a_{n+1}| \geq |a_n| > 0$ for $n \geq n_0$, then by **theorem 10.1.3**, $\sum a_n$ diverges.

Theorem 10.4.3: Ratio convergence \rightarrow Root convergence

$$\lim_{n \rightarrow \infty} \inf(\frac{c_{n+1}}{c_n}) \leq \lim_{n \rightarrow \infty} \inf(\sqrt[n]{c_n})$$

$$\lim_{n \rightarrow \infty} \sup(\sqrt[n]{c_n}) \leq \lim_{n \rightarrow \infty} \sup(\frac{c_{n+1}}{c_n})$$

Proof

Let $\alpha = \lim_{n \rightarrow \infty} \inf(\frac{c_{n+1}}{c_n})$. If $\alpha = -\infty$, then $-\infty \leq \lim_{n \rightarrow \infty} \inf(\sqrt[n]{c_n})$ holds true.
 If α is finite, there is a $\beta \leq \alpha$ and N such that for $n \geq N$, $\frac{c_{n+1}}{c_n} \geq \beta$ so $c_{N+p} \geq \beta^p c_N$.
 Then, $c_n \geq c_N \beta^{-N} \beta^n$ so $\sqrt[n]{c_n} \geq \sqrt[n]{c_N \beta^{-N} \beta^n} = \sqrt[n]{c_N} \beta^{-N/n} \beta$. Thus, $\lim_{n \rightarrow \infty} \inf(\sqrt[n]{c_n}) \geq \beta = \alpha$.
 Let $\alpha = \lim_{n \rightarrow \infty} \sup(\frac{c_{n+1}}{c_n})$. If $\alpha = \infty$, then $\lim_{n \rightarrow \infty} \sup(\sqrt[n]{c_n}) \leq \infty$ holds true.
 If α is finite, there is a $\beta \geq \alpha$ and N such that for $n \geq N$, $\frac{c_{n+1}}{c_n} \leq \beta$ so $c_{N+p} \leq \beta^p c_N$.
 Then, $c_n \leq c_N \beta^{-N} \beta^n$ so $\sqrt[n]{c_n} \leq \sqrt[n]{c_N \beta^{-N} \beta^n} = \sqrt[n]{c_N} \beta^{-N/n} \beta$. Thus, $\lim_{n \rightarrow \infty} \sup(\sqrt[n]{c_n}) \leq \beta = \alpha$.

10.5 Power Series

Definition 10.5.1: Power series

For a sequence $\{c_n\} \in \mathbb{C}$, the series $\sum_{n=0}^{\infty} c_n z^n$ is a power series. c_n are the coefficients and $z \in \mathbb{C}$.

Theorem 10.5.2: Radius of Convergence

For power series $\sum c_n z^n$, let $\alpha = \lim_{n \rightarrow \infty} \sup(\sqrt[n]{|c_n|})$ and $R = \frac{1}{\alpha}$. Then $\sum c_n z^n$ converges if $|z| < R$ and diverges if $|z| > R$.

Proof

Let $a_n = c_n z^n$. Using the **root test**,

$$\lim_{n \rightarrow \infty} \sup(\sqrt[n]{|a_n|}) = \lim_{n \rightarrow \infty} \sup(\sqrt[n]{|c_n z^n|})$$

$$= |z| \lim_{n \rightarrow \infty} \sup(\sqrt[n]{|c_n|}) = \frac{|z|}{R}$$

Thus, $\sum c_n z^n$ converges if $\frac{|z|}{R} < 1$ and diverges if $\frac{|z|}{R} > 1$

10.6 Summation By Parts

Theorem 10.6.1: Summation by parts

For sequences $\{a_n\}$, $\{b_n\}$, let $A_n = \sum_{k=0}^n a_k$. Then for $0 \leq p \leq q$:

$$\sum_{n=p}^q a_n b_n = (\sum_{n=p}^{q-1} A_n(b_n - b_{n+1})) + A_q b_q - A_{p-1} b_p$$

Proof

$$\begin{aligned} \sum_{n=p}^q a_n b_n &= \sum_{n=p}^q (A_n - A_{n-1}) b_n \\ &= \sum_{n=p}^q A_n b_n - \sum_{n=p}^q A_{n-1} b_n = \sum_{n=p}^q A_n b_n - \sum_{n=p-1}^{q-1} A_n b_{n+1} \\ &= \sum_{n=p}^{q-1} A_n b_n - \sum_{n=p}^{q-1} A_n b_{n+1} + A_q b_q - A_{p-1} b_p \\ &= (\sum_{n=p}^{q-1} A_n(b_n - b_{n+1})) + A_q b_q - A_{p-1} b_p \end{aligned}$$

Theorem 10.6.2: Conditions for convergent $\sum a_n b_n$

Suppose for $\{a_n\}$, $\{b_n\}$:

- partial sums A_n of $\sum a_n$ form a bounded sequence
- $b_i \geq b_{i+1}$
- $\lim_{n \rightarrow \infty} b_n = 0$

Then $\sum a_n b_n$ converges.

Proof

Since $\{A_n\}$ is bounded, $|A_n| \leq M$ for all n .
 Since $\{b_n\}$ is monotonically decreasing and $\lim_{n \rightarrow \infty} b_n = 0$, then for $\epsilon > 0$, there is a N such that $b_N \leq \frac{\epsilon}{2M}$. Then for $N \leq p \leq q$:

$$\begin{aligned} |\sum_{n=p}^q a_n b_n| &= |(\sum_{n=p}^{q-1} A_n(b_n - b_{n+1})) + A_q b_q - A_{p-1} b_p| \\ &\leq M |\sum_{n=p}^{q-1} (b_n - b_{n+1}) + b_q + b_p| = 2M b_p \leq 2M b_N \leq \epsilon \end{aligned}$$

Corollary 10.6.3: Convergent series of Alternating Sequences

Suppose for $\{c_n\}$:

- $|c_i| \geq |c_{i+1}|$
- $c_{2i-1} \geq 0$ and $c_{2i} \leq 0$
- $\lim_{n \rightarrow \infty} c_n = 0$

Then $\sum c_n$ converges.

Proof

From **theorem 10.6.2**, let $a_n = (-1)^{n+1}$ and $b_n = |c_n|$.

Corollary 10.6.4: Convergent power series at radius of convergence

Suppose for $\{c_n\}$:

- Radius of convergence of $\sum c_n z^n$ is 1
- $c_i \geq c_{i+1}$
- $\lim_{n \rightarrow \infty} c_n = 0$

Then $\sum c_n z^n$ converges at every point where $|z| = 1$ except possibly $z = 1$.

Proof

From **theorem 10.6.2**, let $a_n = z^n$ and $b_n = c_n$.
 A_n of $\sum a_n$ form a bounded sequence since $|A_n| = |\sum_0^n z^n| = \left| \frac{1-z^{n+1}}{1-z} \right| \leq \frac{2}{|1-z|}$.

10.7 Absolute Convergence**Definition 10.7.1: Absolute convergence**

$\sum a_n$ converges absolutely if $\sum |a_n|$ converges.

If $\sum a_n$ converges, but $\sum |a_n|$ diverges, then $\sum a_n$ converges non-absolutely.

Theorem 10.7.2: Absolute convergence \rightarrow convergence

If $\sum a_n$ converges absolutely, then $\sum a_n$ converges

Proof

Since $\sum a_n$ converges absolutely, then for every $\epsilon > 0$, there is an integer N such that for $m \geq n \geq N$, $|\sum_{k=n}^m |a_k|| = \sum_{k=n}^m |a_k| \leq \epsilon$.
 Thus, $|\sum_{k=n}^m a_k| \leq \sum_{k=n}^m |a_k| \leq \epsilon$ so $\sum a_n$ converges.

10.8 Addition & Multiplication of Series**Theorem 10.8.1: Addition and Scalar Multiplication**

If $\sum a_n = A$ and $\sum b_n = B$, then $\sum (a_n + b_n) = A + B$ and $\sum ca_n = cA$.

Proof

Let $A_n = \sum_{k=0}^n a_k$ and $B_n = \sum_{k=0}^n b_k$.
 Then $A_n + B_n = \sum_{k=0}^n a_k + b_k$ so $\lim_{n \rightarrow \infty} A_n + B_n = A + B$.
 Then $\lim_{n \rightarrow \infty} cA_n = \underbrace{A + \dots + A}_c = cA$

Definition 10.8.2: Cauchy Product

For $\sum a_n$ and $\sum b_n$, let $c_n = \sum_{k=0}^n a_k b_{n-k}$ and the product as $\sum c_n$.

$$\begin{aligned}\sum_{n=0}^{\infty} a_n z^n \sum_{n=0}^{\infty} b_n z^n &= (a_0 + a_1 z + a_2 z^2 + \dots + a_n z^n) (b_0 + b_1 z + b_2 z^2 + \dots + b_n z^n) \\ &= a_0 b_0 + (a_0 b_1 + a_1 b_0) z + (a_0 b_2 + a_1 b_1 + a_2 b_0) z^2 + \dots\end{aligned}$$

Theorem 10.8.3: Conditions $\sum c_n = AB$

Suppose

- $\sum_{n=0}^{\infty} a_n$ converges absolutely
- $\sum_{n=0}^{\infty} a_n = A$
- $\sum_{n=0}^{\infty} b_n = B$
- $c_n = \sum_{k=0}^{\infty} a_k b_{n-k}$

Then $\sum_{n=0}^{\infty} c_n = AB$.

Proof

Let $A_n = \sum_{k=0}^n a_k$, $B_n = \sum_{k=0}^n b_k$, $C_n = \sum_{k=0}^n c_k$, and $\beta_n = B_n - B$.

$$\begin{aligned}C_n &= a_0 b_0 + (a_0 b_1 + a_1 b_0) + \dots + (a_0 b_n + \dots + a_n b_0) \\ &= a_0 B_n + a_1 B_{n-1} + \dots + a_n B_0 \\ &= a_0 (B + \beta_n) + a_1 (B + \beta_{n-1}) + \dots + a_n (B + \beta_0) \\ &= A_n B + a_0 \beta_n + a_1 \beta_{n-1} + \dots + a_n \beta_0\end{aligned}$$

Let $\gamma_n = a_0 \beta_n + a_1 \beta_{n-1} + \dots + a_n \beta_0$ so $C_n = A_n B + \gamma_n$.

Since a_n converges absolutely, then $\sum_{n=0}^{\infty} |a_n| = \alpha$.

Since $\sum_{n=0}^{\infty} b_n = B$, then $\beta_n \rightarrow 0$.

Then for $\epsilon > 0$, there is a N such that $|\beta_n| \leq \frac{\epsilon}{\alpha}$ for $n \geq N$.

$$\begin{aligned}|\gamma_n| &\leq |\beta_0 a_n + \dots + \beta_N a_{n-N}| + |\beta_{N+1} a_{n-N-1} + \dots + \beta_n a_0| \\ &\leq |\beta_0 a_n + \dots + \beta_N a_{n-N}| + |a_{n-N-1} + \dots + a_0| \frac{\epsilon}{\alpha} \\ &\leq |\beta_0 a_n + \dots + \beta_N a_{n-N}| + \alpha \frac{\epsilon}{\alpha}\end{aligned}$$

Thus, with a fixed N , since $a_n \rightarrow 0$, then $\lim_{n \rightarrow \infty} |\gamma_n| \leq \epsilon$ so $\lim_{n \rightarrow \infty} \gamma_n = 0$.

Thus, $\lim_{n \rightarrow \infty} C_n = \lim_{n \rightarrow \infty} A_n B + \gamma_n = AB$.

Theorem 10.8.4: By Cauchy Product, $\sum c_n = C$ implies $C = AB$

If $\sum a_n = A$, $\sum b_n = B$, $\sum c_n = C$ where $c_n = a_0 b_n + \dots + a_n b_0$, then $C = AB$.

11 Continuity


11.1 Limits of Functions

Definition 11.1.1: Limits of functions

For metric spaces X, Y , let $E \subset X$, f maps E into Y , and $p \in E'$.

Then $\lim_{x \rightarrow p} f(x) = q$ if there is a $q \in Y$ such that:

For every $\epsilon > 0$, there is a $\delta > 0$ such that for all $x \in E$ where $d_X(x, p) < \delta$, then $d_Y(f(x), q) < \epsilon$



Theorem 11.1.2: Sequence definition of $\lim_{x \rightarrow p} f(x) = q$

$\lim_{x \rightarrow p} f(x) = q$ if and only if $\lim_{n \rightarrow \infty} f(p_n) = q$ for every sequence $\{p_n\} \in E$ where $p_n \neq p$ and $\lim_{n \rightarrow \infty} p_n = p$.

Proof

Suppose $\lim_{x \rightarrow p} f(x) = q$.

For $\epsilon > 0$, there is a $\delta > 0$ such that $d_Y(f(x), q) < \epsilon$ if $x \in E$ and $d_X(x, p) < \delta$.

Choose $\{p_n\} \in E$ such that $p_n \neq p$ and $\lim_{n \rightarrow \infty} p_n = p$.

Then for $\delta > 0$, there is N such that for $n > N$, then $d_X(p_n, p) < \delta$ so $d_Y(f(p_n), q) < \epsilon$.

Suppose $\lim_{x \rightarrow p} f(x) \neq q$. Then there is a $\epsilon > 0$ such that for every $\delta > 0$, there is a $x \in E$ where $d_Y(f(x), q) \geq \epsilon$, but $d_X(x, p) < \delta$. Let $\delta_n = \frac{1}{n}$ and thus, there is a $\{p_n\}$ where $p_n \neq p$ and $\lim_{n \rightarrow \infty} p_n = p$, but $\lim_{n \rightarrow \infty} f(p_n) \neq q$.

Corollary 11.1.3: A limit of a function is unique

If f has a limit at p , this limit is unique.

Proof

If $\lim_{x \rightarrow p} f(x) = q$, then by **theorem 11.1.2**, $\lim_{n \rightarrow \infty} f(p_n) = q$ for every $\{p_n\} \in E$ where $p_n \neq p$ and $\lim_{n \rightarrow \infty} p_n = p$.

Thus, if there exists $\lim_{x \rightarrow p} f(x) = q'$, then there is a $\{p_n\} \in E$ where $p_n \neq p$ and $\lim_{n \rightarrow \infty} p_n = p$, but $\lim_{n \rightarrow \infty} f(p_n) = q'$ which is a contradiction.

Theorem 11.1.4: Arithmetic operations on functions of limits

Let $E \subset X$, $p \in E'$, and $f(x), g(x) \in \mathbb{C}$ so $\lim_{x \rightarrow p} f(x) = A$, $\lim_{x \rightarrow p} g(x) = B$.

(a) $\lim_{x \rightarrow p} (f + g)(x) = A + B$

(b) $\lim_{x \rightarrow p} (fg)(x) = AB$

(c) $\lim_{x \rightarrow p} \left(\frac{f}{g}\right)(x) = \frac{A}{B}$

11.2 Continuous Functions

Definition 11.2.1: Continuous functions on a set

Suppose X, Y are metric spaces, $E \subset X$, $p \in E$, and f maps E into Y .

f is continuous at p if:


For every $\epsilon > 0$, there is a $\delta > 0$ such that for all $x \in E$

where $d_X(x, p) < \delta$, then $d_Y(f(x), f(p)) < \epsilon$

$f(p)$ have to be defined to be continuous.

If f is continuous at every $p \in E$, then f is continuous on E .

f is continuous at isolated points since regardless of ϵ , there is a $\delta > 0$ such that $d_X(x, p) < \delta$ is $x = p$ so $d_Y(f(x), f(p)) = 0 < \epsilon$.



Theorem 11.2.2: Continuity at $p \Leftrightarrow \lim_{x \rightarrow p} f(x) = f(p)$

Suppose $E \subset X$, $p \in E$, and f maps E into Y . Let $p \in E$.

Then f is continuous at p if and only if $\lim_{x \rightarrow p} f(x) = f(p)$.

Proof

If f is continuous at p , then for every $\epsilon > 0$, there is a $\delta > 0$ such that $d_Y(f(x), f(p)) < \epsilon$ for all $x \in E$ where $d_X(x, p) < \delta$. Thus, $\lim_{x \rightarrow p} f(x) = f(p)$.

If $\lim_{x \rightarrow p} f(x) = f(p)$, then for every $\epsilon > 0$, there is a $\delta > 0$ where $d_Y(f(x), f(p)) < \epsilon$ for all $x \in E$ where $d_X(x, p) < \delta$. Thus, f is continuous at p .

Theorem 11.2.3: Continuity Chain Rule

Suppose $E \subset X$, $f: E \rightarrow Y$, $g: f(E) \rightarrow Z$, and $h: E \rightarrow Z$ where $h(x) = g(f(x))$.

If f is continuous at p and g is continuous at $f(p)$, then h is continuous at p .

Proof

Since g is continuous at $f(p)$, then for $\epsilon > 0$, there is a δ_1 such that:


$d_Z(g(y), g(f(p))) < \epsilon$ for $d_Y(y, f(p)) < \delta_1$ where $y \in f(E)$

Since f is continuous at p , there is a $\delta_2 > 0$ such that:

$d_Y(f(x), f(p)) < \delta_1$ for $d_X(x, p) < \delta_2$ where $x \in E$

Thus, $d_Z(h(x), h(p)) = d_Z(g(f(x)), g(f(p))) < \epsilon$ for $d_X(x, p) < \delta_2$ where $x \in E$.

Thus, h is continuous at p .



Theorem 11.2.4: Continuous functions map open sets to open sets

$f: X \rightarrow Y$ is continuous on X if and only if:

$f^{-1}(V)$ is open in X for every open set V in Y .

Proof

Suppose f is continuous on X and V is an open set in Y .

Suppose $p \in X$ and $f(p) \in V$. Since V is open, there exists $\epsilon > 0$ such that $y \in V$ if $d_Y(y, f(p)) < \epsilon$. Since f is continuous at p , there exists $\delta > 0$ such that $d_Y(f(x), f(p)) < \epsilon$ for $d_X(x, p) < \delta$. Thus, $x \in f^{-1}(V)$ for $d_X(x, p) < \delta$.

Suppose $f^{-1}(V)$ is open in X for every open V in Y .

Fix $p \in X$ and $\epsilon > 0$. Let V be the set of all $y \in Y$ such that $d_Y(y, f(p)) < \epsilon$ so V is open and thus, $f^{-1}(V)$ is open. Thus, there exists $\delta > 0$ such that $x \in f^{-1}(V)$ for $d_X(x, p) < \delta$. Since $x \in f^{-1}(V)$, then $f(x) \in V$ so $d_Y(f(x), f(p)) < \epsilon$.

Corollary 11.2.5: Continuous functions map closed sets to closed sets

$f: X \rightarrow Y$ is continuous on X if and only if:

$f^{-1}(C)$ is closed in X for every closed set C in Y .

Proof

By **theorem 11.2.4**, f is continuous if and only if $f^{-1}(V)$ is open in X for every open set V in Y . Let $C = V^c$. Since V is open, then C is closed.

Since $f^{-1}(C) = f^{-1}(V^c) = (f^{-1}(V))^c$, then $f^{-1}(C)$ is closed since $f^{-1}(V)$ is open.

Theorem 11.2.6: Continuous functions

Let f, g be complex continuous functions on X .

Then $f+g$, fg , and $\frac{f}{g}$ where $g \neq 0$ for all $x \in X$ are continuous on X .

Proof

If x is an isolated point, $f+g$, fg , and $\frac{f}{g}$ are continuous by definition. If x is a limit point, then by **theorems 11.1.4 and 11.2.2**, $f+g$, fg , and $\frac{f}{g}$ are continuous since

- $\lim_{x \rightarrow p} (f + g)(x) = \lim_{x \rightarrow p} f(x) + \lim_{x \rightarrow p} g(x) = f(p) + g(p)$
- $\lim_{x \rightarrow p} (fg)(x) = \lim_{x \rightarrow p} f(x) \lim_{x \rightarrow p} g(x) = f(p)g(p)$
- $\lim_{x \rightarrow p} \left(\frac{f}{g}\right)(x) = \frac{\lim_{x \rightarrow p} f(x)}{\lim_{x \rightarrow p} g(x)} = \frac{f(p)}{g(p)}$

Theorem 11.2.7: Continuous functions on \mathbb{R}^k

(a) Let $f_1, \dots, f_k: X \rightarrow \mathbb{R}$ and $f: X \rightarrow \mathbb{R}^k$ where $f(x) = (f_1(x), \dots, f_k(x))$.

Then f is continuous if and only if f_1, \dots, f_k are continuous.

(b) If f and g are continuous mappings of X into \mathbb{R}^k , then $f + g$ and $f \cdot g$ are continuous on X .

Proof

Since $|f_i(x) - f_i(y)| \leq \sqrt{\sum_{i=1}^k |f_i(x) - f_i(y)|^2} = |f(x) - f(y)|$, then if f is continuous, then each f_i is continuous and vice versa.

Since f, g are continuous, then by part a, each f_i, g_i are continuous. Then by **theorem 11.2.6**, each $f_i + g_i$ and $f_i g_i$ are continuous so by part a, $f + g$ and $f \cdot g$ are continuous.

Thus, every polynomial, rational, and absolute value function is continuous since polynomials are $x_1 \cdot \dots \cdot x_k$ where each x_i is continuous, rationals are polynomials divided by polynomials, and $||x| - |y|| \leq |x - y|$ implies $|x|$ is continuous.

11.3 Continuity and Compactness

Definition 11.3.1: Bounded Functions


$f: E \rightarrow \mathbb{R}^k$ is bounded if there is a $M \in \mathbb{R}$ such that $f(x) \leq M$ for all $x \in E$.

Theorem 11.3.2: Continuous functions from compact spaces are compact

Suppose f is a continuous mapping of a compact metric space X into a metric space Y . Then $f(X)$ is compact.

Proof

Let $\{V_\alpha\}$ be an open cover of $f(X)$. Since f is continuous, then by [theorem 11.2.4](#), each $f^{-1}(V_\alpha)$ is open. Since X is compact, there is n where $X \subset f^{-1}(V_{\alpha_1}) \cup \dots \cup f^{-1}(V_{\alpha_n})$. Thus, $f(X) \subset V_{\alpha_1} \cup \dots \cup V_{\alpha_n}$ so $f(X)$ is compact.



Theorem 11.3.3: Continuous functions from compact to \mathbb{R}^k are bounded

If f is a continuous mapping of a compact metric space X into \mathbb{R}^k , then $f(X)$ is closed and bounded.

Proof

By [theorem 11.2.2](#), $f(X)$ is compact. By [theorem 6.3.13](#), $f(X)$ is closed and bounded.

Theorem 11.3.4: Generalized extreme value theorem

Suppose f is a continuous real function of a compact metric space X such that $M = \sup_{x \in X} f(x)$ and $m = \inf_{x \in X} f(x)$.

Then there exists $p, q \in X$ such that $f(p) = M$ and $f(q) = m$.

Proof

By [theorem 11.3.3](#), $f(X)$ is closed and bounded. Let $M = \sup_{x \in X} f(x)$, $m = \inf_{x \in X} f(x)$. Since $f(X)$ is bounded, then $M, m \in (f(X))'$ and since $f(X)$ is closed, then $M, m \in f(X)$. Thus, there exists $p, q \in X$ such that $f(p) = M$ and $f(q) = m$.

Theorem 11.3.5: If f is continuous 1-1, then f^{-1} is continuous

Suppose f is a continuous 1-1 mapping of a compact metric space X onto a metric space Y . Then f^{-1} is a continuous mapping of Y onto X .

Proof

Let V be an open set in X . Since V^c is closed and $V^c \subset$ compact set X , then by [theorem 6.3.5](#), V^c is compact. Thus by [theorem 11.3.2](#), $f(V^c)$ is a compact subset of Y so $f(V^c)$ is closed. Since f is 1-1 and onto, $f(V^c) = (f(V))^c$ so $f(V)$ is open. Since from any open set V in X , $f(V)$ is open in Y , then by [theorem 11.2.4](#), f^{-1} is continuous.

Definition 11.3.6: Uniformly Continuous

Let $f: X \rightarrow Y$. Then f is uniformly continuous on X if:

For every $\epsilon > 0$, there is a $\delta > 0$ such that for all $p, q \in X$ where $d_X(p, q) < \delta$, then $d_Y(f(p), f(q)) < \epsilon$.

Theorem 11.3.7: Continuous functions on compact are uniformly continuous

Let f be a continuous mapping of a compact metric space X into metric space Y . Then f is uniformly continuous on X .

Proof

For $\epsilon > 0$, since f is continuous, then for each $p \in X$, there is a $\phi(p)$ such that for all $q \in X$ where $d_X(q, p) < \phi(p)$, $d_Y(f(q), f(p)) < \frac{\epsilon}{2}$.

Let $J(p)$ be the set of all $q \in X$ where $d_X(q, p) < \frac{1}{2}\phi(p)$.

Since the set of all $J(p)$ is an open cover of X and since X is compact, then there is a n such that $X \subset J(p_1) \cup \dots \cup J(p_n)$. Let $\delta = \frac{1}{2} \min(\phi(p_1), \dots, \phi(p_n)) > 0$.

Then for $p, q \in X$ where $d_X(p, q) < \delta$, there is a m where $1 \leq m \leq n$ such that $p \in J(p_m)$ so $d_X(p, p_m) < \frac{1}{2}\phi(p_m)$. Thus:

$$\begin{aligned} d_X(q, p_m) &\leq d_X(q, p) + d_X(p, p_m) < \delta + \frac{1}{2}\phi(p_m) \leq \phi(p_m) \\ d_Y(f(p), f(q)) &\leq d_Y(f(p), f(p_m)) + d_Y(f(p_m), f(q)) < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon \end{aligned}$$

Theorem 11.3.8: Continuous functions from noncompact \nrightarrow uniformly continuous

Let E be a noncompact set in \mathbb{R}^1 .

- (a) There exists a continuous function which is not bounded.
- (b) There exists a continuous, bounded function which has no maximum.
- (c) If E is bounded, there exists a continuous function which is not uniformly continuous.

Proof

Suppose E is bounded so there is a $x_0 \in E'$, but $x_0 \notin E$.

Consider $f(x) = \frac{1}{x-x_0}$ which is continuous on E , but unbounded.

For $\epsilon > 0$ and $\delta > 0$, there is a $x \in E$ such that $|x - x_0| < \delta$. Take t close enough to x_0 so $|f(t) - f(x_0)| > \epsilon$, but $|t - x| < \delta$. Thus, f is not uniformly continuous.

Consider $g(x) = \frac{1}{1+(x-x_0)^2}$ which is continuous on E and bounded since $g(x) \in (0,1)$. Since $\sup_{x \in E} g(x) = 1$, but $g(x) < 1$ for all $x \in E$, then g has no maximum on E .

11.4 Continuity and Connectedness**Theorem 11.4.1: Continuous functions map connected to connected**

If f is a continuous mapping of X into Y and E is a connected subset of X , then $f(E)$ is connected.

Proof

Suppose $f(E) = A \cup B$ where A and B are nonempty separated subsets of Y .

Let $G = E \cap f^{-1}(A)$ and $H = E \cap f^{-1}(B)$. Then $E = G \cup H$.

Since $A \subset \bar{A}$, $G \subset f^{-1}(\bar{A})$. Since f is continuous, then $f^{-1}(\bar{A})$ is closed so $\bar{G} \subset f^{-1}(\bar{A})$. Thus, $f(\bar{G}) \subset \bar{A}$.

Since $f(H) = B$ and $\bar{A} \cap B$ is empty, $\bar{G} \cap H$ is empty. Similarly, $G \cap \bar{H}$ is empty so G and H are separated which contradicts that $E = G \cup H$ is connected.

Theorem 11.4.2: Generalized Intermediate Value Theorem

Let f be a continuous real function on $[a, b]$. If $f(a) < c < f(b)$, then there exists $x \in (a, b)$ such that $f(x) = c$.

Proof

Since $[a, b]$ is connected, then by [theorem 11.4.1](#), $f([a, b])$ is a connected subset of \mathbb{R}^1 . Thus, by [theorem 7.2.2](#), any c where $f(a) < c < f(b)$ is $c \in f(x)$ for some $x \in [a, b]$.

11.5 Discontinuities**Definition 11.5.1: Right and Left Limits**

Let f be defined on (a, b) .

Then for any x where $x \in [a, b)$, $f(x+) = q$ if:

$f(t_n) \rightarrow q$ as $n \rightarrow \infty$ for all sequences $\{t_n\}$ in (x, b) such that $t_n \rightarrow x$.

Then for any x where $x \in (a, b]$, $f(x-) = q$ if:

$f(t_n) \rightarrow q$ as $n \rightarrow \infty$ for all sequences $\{t_n\}$ in (a, x) such that $t_n \rightarrow x$.

Then $\lim_{t \rightarrow x} f(t)$ exists if and only if $f(x-) = f(x+) = \lim_{t \rightarrow x} f(t)$.

Definition 11.5.2: Types of discontinuities

If f is discontinuous at x , but $f(x+)$ and $f(x-)$ exists, then f have a simple discontinuity of the first kind else it is a discontinuity of the second kind.

Thus, a simple discontinuity is either:

- $f(x-) \neq f(x+)$
- $f(x-) = f(x+) \neq f(x)$

11.6 Monotonic Functions**Definition 11.6.1: Monotonic**

$f: (a, b) \rightarrow \mathbb{R}$ is monotonically increasing if $f(x) \leq f(y)$ for $a < x < y < b$.

$f: (a, b) \rightarrow \mathbb{R}$ is monotonically decreasing if $f(x) \geq f(y)$ for $a < x < y < b$.

Theorem 11.6.2: Right and Left Limits of monotonic functions on (a, b)

Let f be monotonically increasing on (a, b) .

Then $f(x+)$ and $f(x-)$ exists at every $x \in (a, b)$ where:

$$\sup_{t \in (a, x)} f(t) = f(x-) \leq f(x) \leq f(x+) = \inf_{t \in (x, b)} f(t)$$

Furthermore, for $a < x < y < b$, $f(x+) \leq f(y-)$.

Properties analogous for monotonically decreasing functions.

Proof

Since f is monotonically increasing, then for $t \in (a, x)$, $f(t)$ is bounded above by $f(x)$ and thus, by the least upper bounded property, $\sup_{t \in (a, x)} f(t)$ exists.

For $\epsilon > 0$, there exists a $\delta > 0$ such that $\sup_{t \in (a, x)} f(t) - \epsilon < f(x - \delta) \leq \sup_{t \in (a, x)} f(t)$ for $a < x - \delta < x$. Since $f(x - \delta) \leq f(t) \leq \sup_{t \in (a, x)} f(t)$ for $t \in (x - \delta, x)$, then $|f(t) - \sup_{t \in (a, x)} f(t)| < \epsilon$ for $t \in (x - \delta, x)$ so $f(x-) = \sup_{t \in (a, x)} f(t)$.

For $\epsilon > 0$, there exists a $\delta > 0$ such that $\inf_{t \in (x, b)} f(t) < f(x + \delta) \leq \inf_{t \in (x, b)} f(t) + \epsilon$ for $x < x + \delta < b$. Since $f(x + \delta) \geq f(t) \geq \inf_{t \in (x, b)} f(t)$ for $t \in (x, x + \delta)$, then $|f(t) - \inf_{t \in (x, b)} f(t)| < \epsilon$ for $t \in (x, x + \delta)$ so $f(x+) = \inf_{t \in (x, b)} f(t)$.

Thus, $\sup_{t \in (a, x)} f(t) = f(x-) \leq f(x) \leq f(x+) = \inf_{t \in (x, b)} f(t)$.

If $a < x < y < b$, then:

$$f(x+) = \inf_{t \in (x, b)} f(t) = \inf_{t \in (x, y)} f(t) \leq \sup_{t \in (x, y)} f(t) = \sup_{t \in (a, y)} f(t) = f(y-)$$

Corollary 11.6.3: Monotonics can only have simple discontinuities

Monotonic functions have no discontinuities of the second kind

Proof

By **theorem 11.6.2**, $f(x_-)$ and $f(x_+)$ exists and thus, f can only have simple discontinuities and not discontinuities of the second kind.

Theorem 11.6.4: Discontinuities of monotonics is at most countable

Let f be monotonic on (a,b) .

Then the set of points of (a,b) where f is discontinuous is at most countable.

Proof

Suppose f is increasing. Let E be the set of points where f is discontinuous. Then for $x \in E$, there is a rational $r(x)$ where $f(x_-) < r(x) < f(x_+)$.
 Then for $x_1 < x_2$, by **theorem 11.6.2**, $f(x_1+) \leq f(x_2-)$. Then:

$$f(x_{1-}) < r(x_1) < f(x_{1+}) \leq f(x_{2-}) < r(x_2) < f(x_{2+})$$

 Thus, $r(x_1) \neq r(x_2)$ if $x_1 \neq x_2$.
 Since there is a 1-1 correspondence between E and a subset of rational numbers which is countable, then E is at most countable.
 If f is decreasing, proof is analogous.

11.7 Infinite Limits \ Limits at Infinity**Definition 11.7.1: Neighborhoods in extended reals**

For any real c , a neighborhood of $+\infty = (c, +\infty)$.

For any real c , a neighborhood of $-\infty = (-\infty, c)$.

Definition 11.7.2: Infinite Limits

Let real function f be defined on $E \subset \mathbb{R}$.

Then $f(t) \rightarrow A$ as $t \rightarrow x$ where A and x are extended reals if:

For every neighborhood U of A , there is a neighborhood V of x such that $V \cap E \neq \emptyset$ and $f(t) \in U$ for all $t \in V \cap E$ where $t \neq x$.

Theorem 11.7.3: Arithmetic operations on functions of infinite limits

Let f, g be defined on $E \subset \mathbb{R}$ where $f(t) \rightarrow A$ and $g(t) \rightarrow B$ as $t \rightarrow x$.

(a) If $f(t) \rightarrow A'$, then $A' = A$.

(b) $(f+g)(t) \rightarrow A + B$

(c) $(fg)(t) \rightarrow AB$

(d) $\frac{f}{g}(t) \rightarrow \frac{A}{B}$

12 Differentiation

12.1 Derivative of a function

Definition 12.1.1: Derivative

Let f be defined on any $x \in [a, b]$.

$$\phi(t) = \frac{f(t)-f(x)}{t-x} \text{ for } t \neq x$$

The derivative of f at x :

$$f'(x) = \lim_{t \rightarrow x} \phi(t)$$

if the limit exist as defined by [definition 11.1.1](#).

If f' is defined at x , then f is differentiable at x .

Theorem 12.1.2: Differentiability \rightarrow Continuity

Let f be defined on $[a, b]$.

If f is differentiable at $x \in [a, b]$, then f is continuous at x .

Proof

As $t \rightarrow x$:

$$f(t) - f(x) = \frac{f(t)-f(x)}{t-x} \cdot (t-x) \rightarrow f'(x) \cdot 0 = 0$$

Theorem 12.1.3: Arithmetic operations on differentiation

Suppose f, g are defined on $[a, b]$ and differentiable on $x \in [a, b]$. Then $f+g$, fg , and $\frac{f}{g}$ are differentiable at x :

(a) $(f+g)'(x) = f'(x) + g'(x)$

Proof

$$\begin{aligned} \lim_{t \rightarrow x} \frac{(f+g)(t)-(f+g)(x)}{t-x} &= \lim_{t \rightarrow x} \frac{f(t)-f(x)+g(t)-g(x)}{t-x} \\ &= \lim_{t \rightarrow x} \frac{f(t)-f(x)}{t-x} + \lim_{t \rightarrow x} \frac{g(t)-g(x)}{t-x} = f'(x) + g'(x) \end{aligned}$$

(b) $(fg)'(x) = f'(x)g(x) + f(x)g'(x)$

Proof

$$\begin{aligned} \lim_{t \rightarrow x} \frac{(fg)(t)-(fg)(x)}{t-x} &= \lim_{t \rightarrow x} \frac{f(t)g(t)-f(x)g(x)}{t-x} \\ &= \lim_{t \rightarrow x} \frac{f(t)g(t)-f(x)g(t)+f(x)g(t)-f(x)g(x)}{t-x} \\ &= \lim_{t \rightarrow x} \frac{[f(t)-f(x)]g(t)}{t-x} + \lim_{t \rightarrow x} \frac{f(x)[g(t)-g(x)]}{t-x} \\ &= f'(x)g(x) + f(x)g'(x) \end{aligned}$$

(c) $\left(\frac{f}{g}\right)'(x) = \frac{f'(x)g(x)-f(x)g'(x)}{g^2(x)}$

Proof

$$\begin{aligned} \lim_{t \rightarrow x} \frac{\left(\frac{f}{g}\right)(t)-\left(\frac{f}{g}\right)(x)}{t-x} &= \lim_{t \rightarrow x} \frac{\frac{f(t)}{g(t)}-\frac{f(x)}{g(x)}}{t-x} = \lim_{t \rightarrow x} \frac{f(t)g(x)-f(x)g(t)}{g(t)g(x)(t-x)} \\ &= \lim_{t \rightarrow x} \frac{f(t)g(x)-f(x)g(x)+f(x)g(x)-f(x)g(t)}{g(t)g(x)(t-x)} \\ &= \lim_{t \rightarrow x} \frac{[f(t)-f(x)]g(x)}{g(t)g(x)(t-x)} + \lim_{t \rightarrow x} \frac{f(x)[g(x)-g(t)]}{g(t)g(x)(t-x)} \\ &= \frac{f'(x)g(x)}{g^2(x)} + \frac{f(x)[-g'(x)]}{g^2(x)} = \frac{f'(x)g(x)-f(x)g'(x)}{g^2(x)} \end{aligned}$$

Theorem 12.1.4: Differentiation Chain Rule

Suppose f is continuous on $[a,b]$, $f'(x)$ exists at $x \in [a,b]$, g is defined on interval I containing $f([a,b])$, and g is differentiable at $f(x)$.

If $h(t) = g(f(t))$, then h is differentiable at x and $h'(x) = g'(f(x)) \cdot f'(x)$

Proof

Since f is differentiable at x and g is differentiable at $f(x)$, then:

$$\begin{aligned} f(t) - f(x) &= (t-x) [f'(x) + u(t)] && \text{for } t \in [a,b] \text{ and } \lim_{t \rightarrow x} u(t) = 0 \\ g(s) - g(f(x)) &= (s-f(x)) [g'(f(x)) + v(s)] && \text{for } s \in I \text{ and } \lim_{s \rightarrow f(x)} v(s) = 0 \end{aligned}$$

Thus:

$$\begin{aligned} \lim_{t \rightarrow x} \frac{h(t) - h(x)}{t - x} &= \lim_{t \rightarrow x} \frac{g(f(t)) - g(f(x))}{t - x} \\ &= \lim_{t \rightarrow x} \frac{(f(t) - f(x)) [g'(f(x)) + v(f(t))]}{t - x} \\ &= \lim_{t \rightarrow x} \frac{(t - x) [f'(x) + u(t)] [g'(f(x)) + v(f(t))]}{t - x} \\ &= g'(f(x)) \cdot f'(x) + f'(x) \cdot 0 + g'(f(x)) \cdot 0 + 0 \cdot 0 = g'(f(x)) \cdot f'(x) \end{aligned}$$

12.2 Mean Value Theorems**Definition 12.2.1: Local Extrema**

Let real-valued $f \in X$.

Then f has a local maximum at $p \in X$ if:

There is $\delta > 0$ such that for all $q \in X$ where $d_X(q, p) < \delta$, $f(q) \leq f(p)$.

Then f has a local minimum at $p \in X$ if:

There is $\delta > 0$ such that for all $q \in X$ where $d_X(q, p) < \delta$, $f(q) \geq f(p)$.

Theorem 12.2.2: Derivative at local extrema is 0

Let f be defined on $[a,b]$.

If f has a local maximum at $x \in (a,b)$ and $f'(x)$ exists, then $f'(x) = 0$.

If f has a local minimum at $x \in (a,b)$ and $f'(x)$ exists, then $f'(x) = 0$.

Proof

Suppose x is a local maximum.

Then there is a $\delta > 0$ such that for all $t \in (a,b)$ where $|t - x| < \delta$, then $f(t) \leq f(x)$.

Then for $t < x$, $\frac{f(t) - f(x)}{t - x} \geq 0$. Thus, $\lim_{t \rightarrow x} \frac{f(t) - f(x)}{t - x} = f'(x) \geq 0$.

For $t > x$, $\frac{f(t) - f(x)}{t - x} \leq 0$. Thus, $\lim_{t \rightarrow x} \frac{f(t) - f(x)}{t - x} = f'(x) \leq 0$.

Since $f'(x)$ exists, then $f'(x) = 0$.

Proof is analogous for local minimum.

Theorem 12.2.3: Generalized Mean Value Theorem

If f, g are continuous real functions on $[a,b]$ and differentiable on (a,b) , then there is a $x \in (a,b)$ such that $[f(b) - f(a)] \cdot g'(x) = [g(b) - g(a)] \cdot f'(x)$.

Proof

Let $h(t) = [f(b) - f(a)] \cdot g(t) - [g(b) - g(a)] \cdot f(t)$ for $t \in [a,b]$.

Since f, g are continuous on $[a,b]$ and differentiable on (a,b) , then h is continuous on $[a,b]$ and differentiable on (a,b) . Also, $h(a) = f(b)g(a) - f(a)g(b) = h(b)$.

If h is constant, then $h'(x) = 0$ and thus, theorem holds true for every $x \in (a,b)$.

If $h(t) > h(a)$ for some $t \in (a,b)$, let $x \in [a,b]$ where h attains a local maximum. If $h(t) < h(a)$ for some $t \in (a,b)$, let $x \in [a,b]$ where h attains a local minimum. Then by **theorem 12.2.2**, $h'(x) = 0$ and thus, theorem holds true at local extrema.

Theorem 12.2.4: Mean Value Theorem

If f is a real continuous function on $[a,b]$ and differentiable on (a,b) , then there is a $x \in (a,b)$ such that $f(b) - f(a) = (b-a) f'(x)$.

Proof

From **theorem 12.2.3**, let $g(x) = x$.

Theorem 12.2.5: Sign of derivative determines increasing/decreasing

Suppose f is differentiable on (a,b) .

- (a) If $f'(x) \geq 0$ for all $x \in (a,b)$, then f is monotonically increasing.
- (b) If $f'(x) = 0$ for all $x \in (a,b)$, then f is constant.
- (c) If $f'(x) \leq 0$ for all $x \in (a,b)$, then f is monotonically decreasing

Proof

From **theorem 12.2.4**, $f(x_2) - f(x_1) = (x_2 - x_1) f'(x)$ for $x \in (x_1, x_2) \subset (a,b)$.
 If $f'(x) \geq 0$ for all $x \in (a,b)$, then $f(x_2) - f(x_1) \geq 0$. Since $f(x_2) \geq f(x_1)$ for $x_2 > x_1$, then f is monotonically increasing.
 If $f'(x) = 0$ for all $x \in (a,b)$, then $f(x_2) - f(x_1) = 0$. Since $f(x_2) = f(x_1)$ for $x_2 > x_1$, then f is constant.
 If $f'(x) \leq 0$ for all $x \in (a,b)$, then $f(x_2) - f(x_1) \leq 0$. Since $f(x_2) \leq f(x_1)$ for $x_2 > x_1$, then f is monotonically decreasing.

12.3 Continuity of Derivatives**Theorem 12.3.1: Intermediate values of derivatives exists**

Suppose f is a real differentiable function on $[a,b]$ and $f'(a) < \lambda < f'(b)$.

Then there is a $x \in (a,b)$ such that $f'(x) = \lambda$.

Statement holds true if $f'(a) > f'(b)$.

Proof

Suppose $f'(a) < \lambda < f'(b)$. Let $g(t) = f(t) - \lambda t$.
 Since $f(t), t$ are differentiable on $[a,b]$, then $g(t)$ is differentiable on $[a,b]$.
 Then $g'(a) = f'(a) - \lambda < 0$ so $g(t_1) < g(a)$ for some $t_1 \in (a,b)$.
 Also, $g'(b) = f'(b) - \lambda > 0$ so $g(t_2) < g(b)$ for some $t_2 \in (a,b)$.
 Thus, there is a x where $g(x)$ is a local minimum so $g'(x) = 0$ and thus, $f'(x) = \lambda$.

Corollary 12.3.2: Differentiable functions have no simple discontinuities

If f is differentiable on $[a,b]$, then f' cannot have simple discontinuities on $[a,b]$.

Proof

By **theorem 12.3.1**, $f'(x)$ exists for any $x \in [a,b]$.

12.4 L'Hospital's Rule

Theorem 12.4.1: L'Hospital's Rule

Suppose f, g are real and differentiable on (a, b) and $g'(x) \neq 0$ for all $x \in (a, b)$.

Suppose $\lim_{x \rightarrow a} \frac{f'(x)}{g'(x)} \rightarrow A$. If either:

- $\lim_{x \rightarrow a} f(x) \rightarrow 0$ and $\lim_{x \rightarrow a} g(x) \rightarrow 0$
- $\lim_{x \rightarrow a} g(x) \rightarrow \infty$ or $\lim_{x \rightarrow a} g(x) \rightarrow -\infty$

Then, $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} \rightarrow A$.

Statement holds true if $x \rightarrow b$.

Proof

Consider the case $-\infty \leq A < \infty$.

Choose q such that $A < q$ and r such that $A < r < q$.

Thus, there is a $c \in (a, b)$ such that $a < x < c$ for $\frac{f'(x)}{g'(x)} < r$.

For $a < x < y < c$, then by [theorem 12.2.3](#), there is a $t \in (x, y)$ such that:

$$\frac{f(x)-f(y)}{g(x)-g(y)} = \frac{f'(t)}{g'(t)} < r$$

If $\lim_{x \rightarrow a} f(x) \rightarrow 0$ and $\lim_{x \rightarrow a} g(x) \rightarrow 0$, then as $x \rightarrow a$, $\frac{f(y)}{g(y)} \leq r < q$ for $y \in (a, c)$.

If $\lim_{x \rightarrow a} g(x) \rightarrow \infty$, then keeping y fixed, choose $c_1 \in (a, y)$ such that $g(x) > g(y)$ and $g(x) > 0$ if $a < x < c_1$. Thus:

$$\begin{aligned} \frac{g(x)-g(y)}{g(x)} \cdot \frac{f(x)-f(y)}{g(x)-g(y)} &< \frac{g(x)-g(y)}{g(x)} \cdot r \text{ for } x \in (a, c_1) \\ \frac{f(x)}{g(x)} &< r - r \frac{g(y)}{g(x)} + \frac{f(y)}{g(x)} \end{aligned}$$

Thus as $x \rightarrow a$, there is a $c_2 \in (a, c_1)$ such that $\frac{f(x)}{g(x)} < r < q$ for $x \in (a, c_2)$.

Proof is analogous if $\lim_{x \rightarrow a} g(x) \rightarrow -\infty$.

Thus, $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} \rightarrow A$.

12.5 Derivative of Higher Order

Definition 12.5.1: Derivative of Higher Order

If f has a derivative f' on an interval and f' is differentiable, then the derivative of f' is f'' , the second derivative of f . Then, $f^{(n)}$ is the n th derivative of f .

For $f^{(n)}(x)$ to exist at x , $f^{(n-1)}(t)$ must exist in a neighborhood of x and $f^{(n-1)}$ must be differentiable at x .

If $f^{(n-1)}$ exist in a neighborhood of x , then $f^{(n-2)}$ must be differentiable in that neighborhood and so on until f is differentiable on that neighborhood.

12.6 Taylor's Theorem

Theorem 12.6.1: Taylor's Theorem

Suppose f is a real function on $[a,b]$, $n \in \mathbb{Z}_+$, $f^{(n-1)}$ is continuous on $[a,b]$, $f^n(t)$ exists at every $t \in (a,b)$.

Let $\alpha, \beta \in [a,b]$ be distinct and $P(t) = \sum_{k=0}^{n-1} \frac{f^{(k)}(\alpha)}{k!} (t - \alpha)^k$.

Then there exists a x between α and β such that $f(\beta) = P(\beta) + \frac{f^n(x)}{n!} (\beta - \alpha)^n$

Proof

Let M be the number defined by $f(\beta) = P(\beta) + M(\beta - \alpha)^n$.

Let $g(t) = f(t) - P(t) - M(t - \alpha)^n$ for $t \in [\alpha, \beta]$. Thus, $g^{(n)}(t) = f^{(n)}(t) - n!M$.

Also since $P^{(k)}(\alpha) = f^{(k)}(\alpha)$ for $k = [0, n-1]$, then $g(\alpha) = g'(\alpha) = \dots = g^{(n-1)}(\alpha) = 0$.

Since the choice of M gives $g(\beta) = 0$, then by the Mean Value Theorem, $g'(x_1) = 0$ for some x_1 between α and β .

Since $g'(\alpha) = 0$, then $g''(x_2) = 0$ for some x_2 between α and x_1 .

Thus, $g^{(n)}(x_n) = 0$ for some x_n between α and x_{n-1} so x_n is between α and β .

Thus, there exists an $x_n \in (\alpha, \beta)$ such that:

$$0 = g^{(n)}(x_n) = f^{(n)}(x_n) - n!M$$

$$M = \frac{f^n(x_n)}{n!}$$

12.7 Differentiation of Vector-Valued Functions

Definition 12.7.1: Extending derivative to Vector-Valued Functions

For vector-valued function $f: t \in [a,b] \rightarrow \mathbb{R}^k$, the derivative of f at x :

$$f'(x) = \lim_{t \rightarrow x} \left| \frac{f(t) - f(x)}{t - x} \right|$$

if the limit exist as defined by [definition 14.1.1](#).

If $f = (f_1, \dots, f_k)$, then $f' = (f'_1, \dots, f'_k)$ and f is differentiable at x if and only if f_1, \dots, f_k are differentiable at x .

Thus, by [theorem 11.2.7](#), these theorems hold true for vector-valued functions:

- [12.1.2](#): If f is differentiable at x , then f is continuous at x .
- [12.1.3a](#): If f, g are differentiable at x , then $f+g, f \cdot g$ are differentiable at x .

However, [theorem 12.2.4: Mean Value Theorem](#) and [theorem 12.4.1: L'Hospital's Rule](#) does not always hold true since [theorem 12.1.3b/c](#), multiplying/dividing vectors by vectors, is not defined for vector-valued functions.

Theorem 12.7.2: Mean Value Theorem for \mathbb{R}^k

Suppose f is a continuous mapping of $[a,b]$ into \mathbb{R}^k and f is differentiable on (a,b) . Then there is a $x \in (a,b)$ such that $|f(b) - f(a)| \leq (b-a) |f'(x)|$

Proof

Let $z = f(b) - f(a)$ and define $\phi(t) = z \cdot f(t)$ for $t \in [a,b]$.

Then $\phi(t)$ is real-valued continuous on $[a,b]$ and differentiable on (a,b) .

Then by the Mean Value Theorem, for some $x \in (a,b)$:

$$\phi(b) - \phi(a) = (b-a) \phi'(x) = (b-a) z \cdot f'(x)$$

Since $\phi(b) - \phi(a) = z \cdot f(b) - z \cdot f(a) = z \cdot z = |z|^2$, then by the Schwarz Inequality:

$$|z|^2 = (b-a) |z \cdot f'(x)| \leq (b-a) |z| |f'(x)|$$

$$|z| \leq (b-a) |f'(x)|$$

$$|f(b) - f(a)| \leq (b-a) |f'(x)|$$

References

- [1] Walter Rudin, *Principles of Mathematical Analysis (3rd Edition)*,
ISBN-13: 978-0070542358