

Solutions to Aluffi's Algebra: Chapter 0

Vladyslav Chorny

April 16, 2026

Contents

V: Irreducibility and factorization in integral domains	5
§1 Chain conditions and existence of factorizations	5
VI: Linear Algebra	11
§1 Free modules revisited	11
§2 Homomorphisms of free modules, I	14
§3 Homomorphisms of free modules, II	15
§4 Presentations and resolutions	20
§5 Classification of finitely generated modules over PIDs	25

V: Irreducibility and factorization in integral domains

§1 Chain conditions and existence of factorizations

1.1

Let R be a Noetherian ring, and let I be an ideal of R . Prove that R/I is a Noetherian ring.

Solution.

Suppose J/I is an ideal of R/I , corresponding to an ideal J of R , $J \supset I$. Since R is Noetherian, $J = (a_1, \dots, a_n) \implies J/I = (\bar{a}_1, \dots, \bar{a}_n)$. $\square$

1.2

Prove that if $R[x]$ is Noetherian, so is R . (This is a ‘converse’ to Hilbert’s basis theorem.)

Solution.

$R \simeq \frac{R[x]}{(x)}$, hence R is Noetherian by the previous exercise. $\square$

1.3

Let k be a field, and let $f \in k[x]$, $f \notin k$. For every subring R of $k[x]$ containing k and f , define a homomorphism $\varphi : k[t] \rightarrow R$ by extending the identity on k and mapping t to f . This makes every such R a $k[t]$ -algebra.

- Prove that $k[x]$ is finitely generated as a $k[t]$ -module.
- Prove that every subring R as above is finitely generated as a $k[t]$ -module.

- Prove that every subring of $k[x]$ containing k is a Noetherian ring.

Solution.

Let $\deg f = n$ and let us show that every $g \in k[x]$ can be expressed as

$$g(x) = \sum a_i(x)f(x)^i$$

with $\deg a_i < n$.

The statement trivially holds for all g such that $\deg g < n$. If $\deg g \geq n$, then we can write $g = qf + r$, where $\deg q < \deg g$ and $\deg r < n$. We may then inductively assume that $q = \sum a_i f^i$, with $\deg a_i < n$, which implies

$$g = \left(\sum a_i f^i \right) f + r = \sum a_i f^{i+1} + r = \sum a_i^* f^i$$

It follows then that for all $g \in k[x]$:

$$g(x) = \sum a_i(x)f(x)^i = \sum_{i=0}^m \sum_{j=0}^{n-1} a_{ij} x^j f(x)^i = \sum_j x^j \sum_i a_{ij} f(x)^i = \sum p_j(t) \cdot x^j$$

where $p_j(t) = \sum_i a_{ij} t^i$.

Thus, $k[x] = \langle 1, x, \dots, x^{n-1} \rangle$ as a $k[t]$ -module.

If R is a subring of $k[x]$ containing k and f , then R is a submodule of $k[x]$ over $k[t]$. Since $k[t]$ is a PID and $k[x]$ is finitely generated, R is also finitely generated.

If R is a subring of $k[x]$ containing k , then either $R = k$, in which case R is Noetherian, or there exists $f \in R$ such that $f \notin k$, which defines a $k[t]$ -module structure on $k[x]$, making R into a finitely generated submodule.

Suppose I is an ideal of R . I is a submodule of $k[x]$. Indeed, it is closed under addition and $\forall g \in I \forall p \in k[t]$,

$$p(t) \cdot g = \left(\sum a_i f(x)^i \right) \cdot g(x) \in I,$$

since $\sum a_i f^i \in R$.

Therefore, I is finitely generated as a $k[t]$ -module. Let $g_1, \dots, g_m$ be its generators. It follows that $I = (g_1, \dots, g_m)$ as an ideal of R . $\square$

1.4

Let R be the ring of real-valued continuous functions on the interval $[0, 1]$. Prove that R is not Noetherian.

Solution.

Define $N_A = \{f \in R \mid f(a) = 0 \forall a \in A\}$. For any nonempty $A \subset [0, 1]$, N_A is clearly an ideal.

The chain $N_{[0,1]} \subset N_{[0,\frac{1}{2}]} \subset \cdots \subset N_{[0,\frac{1}{n}]} \subset \cdots$ does not stabilize, hence R is not Noetherian. $\square$

1.8

Prove that every ideal in a Noetherian ring R contains a finite product of prime ideals. (Hint: Let $\mathcal{F}$ be the family of ideals that do not contain finite products of prime ideals. If $\mathcal{F}$ is nonempty, it has a maximal element M since R is Noetherian. Since $M \in \mathcal{F}$, M is not itself prime, so $\exists a, b \in R$ s.t. $a \notin M, b \notin M$, yet $ab \in M$. What's wrong with this?)

Solution.

Since $M + (a)$ and $M + (b)$ strictly contain M , both ideals are not in $\mathcal{F}$. Hence, there are prime ideals $P_1, \dots, P_n, P'_1, \dots, P'_m$ such that $P_1 \dots P_n \subset M + (a)$ and $P'_1 \dots P'_m \subset M + (b) \implies P_1 \dots P_n P'_1 \dots P'_m \subset (M + (a))(M + (b)) \subset M + (ab) = M$. This contradicts the fact that M does not contain a finite product of prime ideals. $\square$

1.9

Let R be a commutative ring, and let $I \subseteq R$ be a proper ideal. The reader will prove in Exercise 3.12 that the set of prime ideals containing I has minimal elements (the *minimal primes* of I). Prove that if R is Noetherian, then the set of minimal primes of I is finite. (Hint: Let $\mathcal{F}$ be the family of ideals that do *not* have finitely many minimal primes. If $\mathcal{F} \neq \emptyset$, note that $\mathcal{F}$ must have a maximal element I , and I is not prime itself. Find ideals J_1, J_2 strictly larger than I , such that $J_1 J_2 \subseteq I$, and deduce a contradiction.)

Solution.

I claim that if $I \subset J \subset P$ and P is a minimal prime of I , then P is a minimal prime of J . Indeed, if there is a prime ideal P' such that $J \subset P' \subsetneq P$, then $I \subset P' \subsetneq P$, which contradicts the minimality of P with respect to I .

Now if I is a maximal element of $\mathcal{F}$ defined as above, then I is not itself prime, for otherwise the set of minimal primes of I would consist of a single element: I . Hence, $\exists a, b \in R$ such that $a \notin I, b \notin I, ab \in I \implies I + (a) \supsetneq I, I + (b) \supsetneq I \implies I + (a)$ and $I + (b)$ have finite sets of minimal primes. Let us denote them by M_a and M_b . We also have $(I + (a))(I + (b)) \subset I + (ab) = I$ and thus, if P is a minimal prime of I , then $(I + (a))(I + (b)) \subset P$, and therefore by exercise III.4.19 $I + (a) \subset P$ or $I + (b) \subset P$, which as discussed before implies that P is a minimal prime of $I + (a)$ or $I + (b)$. This gives us a mapping from the collection M of minimal primes of I into $M_a \cup M_b$, and this mapping is obviously injective. It is impossible, since M is infinite and $M_a \cup M_b$ isn't. $\square$

1.10

By Proposition 1.1, a ring R is Noetherian if and only if it satisfies the a.c.c. for ideals. A ring is *Artinian* if it satisfies the d.c.c. (descending chain condition) for ideals. Prove that if R is Artinian and $I \subseteq R$ is an ideal, then R/I is Artinian. Prove that if R is an Artinian integral domain, then it is a field. (Hint: Let $r \in R$, $r \neq 0$. The ideals (r^n) form a descending sequence; hence $(r^n) = (r^{n+1})$ for some n . Therefore...) Prove that Artinian rings have Krull dimension 0 (that is, prime ideals are maximal in Artinian rings).

Solution.

Suppose R is Artinian, $I \subset R$ is an ideal and $J_1 \supset J_2 \supset \dots$ is a descending chain of ideals in R/I . Recall that there is a natural inclusion-preserving correspondence between ideals of R/I and ideals of R containing I . Therefore, there is a chain $I_1 \supset I_2 \supset \dots$ in R , which must stabilize because R is Artinian. It follows that $J_1 \supset J_2 \supset \dots$ also stabilizes.

Now if R is an Artinian integral domain, then $\forall r \in R, r \neq 0, \exists n \in \mathbb{N}$ such that $(r^n) = (r^{n+1}) \implies \exists a \in R$ such that $r^n = ar^{n+1} \implies r^n(1 - ar) = 0$. Since R is an integral domain and $r \neq 0$, we have $1 - ar = 0$, so r is invertible.

Finally, we need to show that prime ideals are maximal if R is Artinian. Assume $P \subset R$ is prime. Then, as shown before, R/P is also Artinian. Moreover, it is an integral domain because P is prime, and hence it is a field $\implies P$ is maximal. $\square$

1.13

Prove that, for nonzero elements, prime $\iff$ irreducible in $\mathbb{Z}$.

Solution.

Primes are irreducibles in all integral domains, so we only need to show sufficiency. If $p \neq 0$ is irreducible in $\mathbb{Z}$, then (p) is maximal among principal ideals of $\mathbb{Z}$. Since in $\mathbb{Z}$ every ideal is principal, (p) is maximal among all ideals of $\mathbb{Z}$, hence (p) is prime and p is prime by definition. $\square$

1.14

For a, b in a commutative ring R , prove that the class of a in $R/(b)$ is prime if and only if the class of b in $R/(a)$ is prime.

Solution.

Let us denote by $[(a)]_{(b)}$ the image of $(a) \subset R$ under the canonical projection on $R/(b)$. Using the result in III.3.3 we get:

$$\bar{a} \text{ is prime in } R/(b) \iff (\bar{a}) = [(a)]_{(b)} \text{ is prime in } R/(b)$$

$$\begin{aligned} &\Longleftrightarrow \frac{R/(b)}{[(a)]_{(b)}} \simeq \frac{R}{(a) + (b)} \simeq \frac{R/(a)}{[(b)]_{(a)}} \text{ is an integral domain} \\ &\Longleftrightarrow [(b)]_{(a)} = (\bar{b}) \text{ is prime in } R/(a) \Longleftrightarrow \bar{b} \text{ is prime in } R/(a). \end{aligned}$$

□

1.17

Consider the subring of $\mathbb{C}$:

$$\mathbb{Z}[\sqrt{-5}] := \{a + bi\sqrt{5} \mid a, b \in \mathbb{Z}\}.$$

- Prove that this ring is isomorphic to $\mathbb{Z}[t]/(t^2 + 5)$.
- Prove that it is a Noetherian integral domain.
- Define a ‘norm’ N on $\mathbb{Z}[\sqrt{-5}]$ by setting $N(a + bi\sqrt{5}) = a^2 + 5b^2$. Prove that $N(zw) = N(z)N(w)$. (Cf. Exercise III.4.10.)
- Prove that the units in $\mathbb{Z}[\sqrt{-5}]$ are ± 1 . (Use the preceding point.)
- Prove that $2, 3, 1 + i\sqrt{5}, 1 - i\sqrt{5}$ are all irreducible nonassociate elements of $\mathbb{Z}[\sqrt{-5}]$.
- Prove that no element listed in the preceding point is prime. (Prove that the rings obtained by mod-ing out the ideals generated by these elements are not integral domains.)
- Prove that $\mathbb{Z}[\sqrt{-5}]$ is not a UFD.

Solution.

$\varphi : \mathbb{Z}[\sqrt{-5}] \rightarrow \frac{\mathbb{Z}[t]}{(t^2+5)}$ defined by $\varphi(a + bi\sqrt{5}) = \overline{a + bt}$ is an isomorphism, which can be trivially verified. Let’s just check that φ preserves products:

$$\begin{aligned} \varphi((a + bi\sqrt{5})(a' + b'i\sqrt{5})) &= \varphi((aa' - 5bb') + (ab' + a'b)i\sqrt{5}) = \\ &= \overline{(aa' - 5bb') + (ab' + a'b)t} = (\text{since } \bar{t}^2 = -5) = \\ &= \overline{(a + bt)} \cdot \overline{(a' + b't)} = \varphi(a + bi\sqrt{5})\varphi(a' + b'i\sqrt{5}) \end{aligned}$$

$\mathbb{Z}[\sqrt{-5}]$ is an integral domain because $\mathbb{C}$ is and it is Noetherian because $\mathbb{Z}[t]$ is.

We can write N as

$$N(a + bi\sqrt{5}) = a^2 + 5b^2 = (a - bi\sqrt{5})(a + bi\sqrt{5}) = (a + bi\sqrt{5}) \cdot \overline{(a + bi\sqrt{5})},$$

where $\bar{x}$ denotes a complex conjugate, so $N(x) = x\bar{x}$, which implies

$$N(zw) = zw\bar{zw} = z\bar{z}w\bar{w} = N(z)N(w)$$

Note that $N(z) \in \mathbb{Z}_{\geq 0} \forall z \in \mathbb{Z}[\sqrt{-5}]$.

If $u \in \mathbb{Z}[\sqrt{-5}]$ is a unit, then

$$1 = N(1) = N(uu^{-1}) = N(u)N(u^{-1}) \implies N(u) = 1$$

If $a^2 + 5b^2 = 1$ with $a, b \in \mathbb{Z}$, then $(a, b) \in \{(1, 0), (-1, 0)\}$ by size consideration. So the only units are 1 and -1 .

The following trivial claims will help us to deal with irreducibility:

1. $\nexists v \in \mathbb{Z}[\sqrt{-5}] \mid N(v) = 2$ or $N(v) = 3$
2. If $v \mid w$, then $N(v) \mid N(w)$
3. If $N(v) = 1$ then v is a unit

To show that 2 is irreducible, assume $2 = vw \implies N(v) \mid N(2) = 4$. Since $N(v) \neq 2$, we must have $N(v) = 1$ or 4, which forces v or w to be a unit, proving that 2 is irreducible.

The same reasoning yields that 3 and $1 \pm i\sqrt{5}$ are irreducible.

None of the considered elements are prime, because $6 = 2 \cdot 3 = (1 - i\sqrt{5})(1 + i\sqrt{5})$. For example, $1 - i\sqrt{5} \mid 2 \cdot 3$, yet $1 - i\sqrt{5} \nmid 2$ and $1 - i\sqrt{5} \nmid 3$, since $N(1 - i\sqrt{5}) = 6 \nmid 4 = N(2)$ and $6 \nmid 9$.

The same factorization $6 = 2 \cdot 3 = (1 - i\sqrt{5})(1 + i\sqrt{5})$ also shows that $\mathbb{Z}[\sqrt{-5}]$ is not a UFD. $\square$

VI: Linear Algebra

§1 Free modules revisited

1.7

Let R be an integral domain, and let $M = R^{\oplus A}$ be a free R -module. Let K be the field of fractions of R , and view M as a subset of $V = K^{\oplus A}$ in the evident way. Prove that a subset $S \subseteq M$ is linearly independent in M (over R) if and only if it is linearly independent in V (over K). Conclude that the rank of M (as an R -module) equals the dimension of V (as a K -vector space). Prove that if S generates M over R , then it generates V over K . Is the converse true?

Solution.

Let $R \hookrightarrow K$ via $r \mapsto \frac{r}{1}$. This induces an inclusion $R^{\oplus A} \hookrightarrow K^{\oplus A}$ defined by:

$$\sum r_i a_i \mapsto \sum \frac{r_i}{1} a_i$$

$\implies$ Assume $S \subset M$ is linearly independent over R . Suppose we have a linear relation in V :

$$\sum_{i \in I} \frac{r_i}{p_i} s_i = 0$$

where $|I| < +\infty$. Then:

$$\sum_{i \in I} P \frac{r_i}{p_i} s_i = 0, \quad \text{where } P = \prod_{i \in I} p_i$$

For all $k \in I$, the term $\frac{P}{p_k} = \prod_{i \in I \setminus \{k\}} p_i$ is an element of R . Hence, we have obtained a linear relation in M . Since S is linearly independent in M , all coefficients must be zero, which implies $r_i = 0$ for all $i \in I$.

$\longleftarrow$ Obvious.

Suppose that S generates M over R and let $v \in V$. Then we can write v as a finite sum:

$$v = \sum_{i \in I} \frac{r_i}{p_i} a_i$$

Let $P = \prod_{i \in I} p_i$. Multiplying by P clears the denominators, meaning $Pv \in M$. Since S generates M , there exist $r_j \in R$ and $s_j \in S$ such that:

$$Pv = \sum_{j \in J} r_j s_j$$

Dividing by P in V , we get:

$$v = \sum_{j \in J} \frac{r_j}{P} s_j \in \langle S \rangle_K$$

Thus, S generates V over K . The converse is not true. For example, the subset $\{2\}$ generates $\mathbb{Q}$ over $\mathbb{Q}$, but does not generate $\mathbb{Z}$ over $\mathbb{Z}$. $\square$

1.9

Let R be a commutative ring, and let M be an R -module. Let $\mathfrak{m}$ be a maximal ideal in R , such that $\mathfrak{m}M = 0$ (that is, $rm = 0$ for all $r \in \mathfrak{m}$, $m \in M$). Define in a natural way a vector space structure over $R/\mathfrak{m}$ on M .

Solution.

We define scalar multiplication by $\bar{r}m := rm$. We must check that this operation is correctly defined. If $\bar{r} = \bar{r}'$, then:

$$r - r' \in \mathfrak{m} \implies (r - r')m = 0 \implies rm = r'm$$

Thus, the operation is well-defined. The other vector space axioms trivially hold because M is an R -module. $\square$

1.10

Let R be a commutative ring, and let F be a free R -module with basis B . Let $\mathfrak{m}$ be a maximal ideal of R , and let $k = R/\mathfrak{m}$. Prove that $F/\mathfrak{m}F \simeq k^{\oplus B}$ as k -vector spaces.

Solution.

Let $\sum r_i b_i \in F$. If $x \in \mathfrak{m}F$, then $x = \sum m_i f_i = \sum m_i \sum r_{ij} b_j = \sum m'_i b_i$, where $m'_i \in \mathfrak{m}$. On the other hand, all $\sum m_i b_i$ with $m_i \in \mathfrak{m}$ are in $\mathfrak{m}F$. Hence, $\mathfrak{m}F = \mathfrak{m}^{\oplus B}$.

$F/\mathfrak{m}F$ is an R -module, and for all $m \in \mathfrak{m}$ and for all $\bar{f} \in F/\mathfrak{m}F$:

$$m\bar{f} = \overline{mf} = \bar{0}$$

Hence, by Exercise 1.9, $F/\mathfrak{m}F$ is a k -vector space.

Now let's prove that $\pi(B)$ is a basis of $F/\mathfrak{m}F$, where π is the natural projection. Obviously, $\pi(B)$ generates $F/\mathfrak{m}F$ since B generates F . Suppose that $\sum_I (r_i + \mathfrak{m})(b_i + \mathfrak{m}F) = 0 + \mathfrak{m}F$. Then:

$$\left(\sum r_i b_i \right) + \mathfrak{m}F = \bar{0} \implies \sum r_i b_i \in \mathfrak{m}F \implies r_i \in \mathfrak{m} \implies r_i + \mathfrak{m} = \bar{0}$$

This implies that $\pi(B)$ is linearly independent, hence it is a basis. Therefore, $F/\mathfrak{m}F \simeq k^{\oplus B}$. $\square$

1.11

Prove that commutative rings satisfy the IBN (Invariant Basis Number) property.

Solution.

Let R be a commutative ring which has a proper ideal and let F be a free R -module. Suppose $F \simeq R^{\oplus A} \simeq R^{\oplus B}$.

R has a maximal ideal $\mathfrak{m}$ and by Exercise 1.10,

$$k^{\oplus A} \simeq F/\mathfrak{m}F \simeq k^{\oplus B}$$

where $k = R/\mathfrak{m}$. But then it follows that $A \simeq B$, since k is a field. $\square$

1.13

Let A be an abelian group such that $\text{End}_{\text{Ab}}(A)$ is a field of characteristic 0. Prove that $A \simeq \mathbb{Q}$.

Solution.

$\text{End}(A)$ acts on A : $\varphi a := \varphi(a)$ and makes the latter into a vector space.

We have an inclusion $\mathbb{Q} \hookrightarrow \text{End}(A)$, since $\text{char}(\text{End } A) = 0$:

$$\frac{p}{q} \mapsto (p \text{ id}) \circ (q \text{ id})^{-1}$$

Hence, A is a $\mathbb{Q}$ -vector space via

$$\frac{p}{q}a = ((p \text{ id}) \circ (q \text{ id})^{-1})(a) = (q \text{ id})^{-1}(pa)$$

Basically, $p \cdot a = (p \text{ id})(a) = pa$ for integer p , and $\frac{1}{q}a = b$ such that $a = qb$. It exists and is unique because $q \text{ id}$ has an inverse, hence is an isomorphism.

Now, suppose $f \in \text{End}_{\text{Ab}}(A)$. Then

$$\forall n \in \mathbb{Z} \quad f(na) = nf(a)$$

We have also $f(a) = mf\left(\frac{1}{m}a\right) \implies f\left(\frac{1}{m}a\right) = \frac{1}{m}f(a)$ for $m \in \mathbb{N}$.

Hence, $f\left(\frac{n}{m}a\right) = nf\left(\frac{1}{m}a\right) = \frac{n}{m}f(a)$ or simply

$$f(ra) = rf(a) \quad \forall r \in \mathbb{Q}$$

This means that f is actually a $\mathbb{Q}$ -vector space endomorphism, so $\text{End}_{\text{Ab}}(A) = \text{End}_{\mathbb{Q}\text{-Vect}}(A)$.

Suppose S is a basis of A , $|S| > 1$ and $a \in S$. Then $A = V \oplus W$, where $V = \langle a \rangle$ and $W = \langle S \setminus \{a\} \rangle$. Define a map $\sigma : A \rightarrow A$ via

$$A \simeq V \oplus W \xrightarrow{\pi_1} V \hookrightarrow A$$

which sends $\lambda a + \sum_{s \in S \setminus \{a\}} \lambda_s s$ to λa .

It is a $\mathbb{Q}$ -vector space homomorphism as a composition of homomorphisms and has a non-trivial kernel, since $S \setminus \{a\} \subset \ker \sigma$ and $S \setminus \{a\}$ is non-empty and does not contain 0. Hence, σ does not have an inverse, contradicting the fact that $\text{End}(A)$ is a field.

It follows that $|S| = \dim A \leq 1$. The case of 0 is not possible, so $\dim_{\mathbb{Q}} A = 1$ and $A \simeq \mathbb{Q}$.

We can also show that for a $\mathbb{Q}$ -vector space V , $\text{End}_{\mathbb{Q}\text{-Vect}}(V) \simeq M_n(\mathbb{Q})$ as rings, where $n = \dim V$. Thus, in the general case, $\text{End}_{\text{Ab}}(V)$ is not a field, and $\mathbb{Q}$ is the only space with its endomorphism ring being a field. $\square$

1.19

Let k be a field, and let $f(x) \in k[x]$ be any polynomial. Prove that there exists a multiple of $f(x)$ in which all exponents of nonzero monomials are *prime* integers. (Example: for $f(x) = 1 + x^5 + x^6$, $(1 + x^5 + x^6)(2x^2 - x^3 + x^5 - x^8 + x^9 - x^{10} + x^{11}) = 2x^2 - x^3 + x^5 + 2x^7 + 2x^{11} - x^{13} + x^{17}$.)

Solution.

Let $V = k[x]/(f(x))$. This is a k -vector space with basis $\overline{1}, \overline{x}, \dots, \overline{x^{n-1}}$, where $n = \deg f$.

Take $n + 1$ distinct primes $p_1, \dots, p_{n+1}$. The system

$$S = \{\overline{x^{p_1}}, \overline{x^{p_2}}, \dots, \overline{x^{p_{n+1}}}\}$$

is linearly dependent, since $|S| = n + 1 > \dim V$. So, for some $\lambda_i \in k$ (not all zero):

$$\begin{aligned} \sum_{i=1}^{n+1} \lambda_i \overline{x^{p_i}} = \overline{0} &\implies \sum_{i=1}^{n+1} \lambda_i x^{p_i} \in (f(x)) \\ \implies \exists p(x) \in k[x] \quad \text{such that} \quad f(x)p(x) &= \sum_{i=1}^{n+1} \lambda_i x^{p_i} \end{aligned}$$

This multiple has only prime exponents, as desired. $\square$

§2 Homomorphisms of free modules, I

§3 Homomorphisms of free modules, II

3.6

Let R be a commutative ring and $M = \langle m_1, \dots, m_r \rangle$ a finitely generated R -module. Let $A \in M_r(R)$ be a matrix such that

$$A \begin{pmatrix} m_1 \\ \vdots \\ m_r \end{pmatrix} = \begin{pmatrix} 0 \\ \vdots \\ 0 \end{pmatrix}.$$

Prove that $\det(A)m = 0$ for all $m \in M$.

Solution.

Let A^* be the adjoint matrix of A . Recall the fundamental property $A^*A = \det(A)I_r$. Starting from the given equation:

$$A \begin{pmatrix} m_1 \\ \vdots \\ m_r \end{pmatrix} = \begin{pmatrix} 0 \\ \vdots \\ 0 \end{pmatrix}$$

Multiply both sides on the left by A^* :

$$A^*A \begin{pmatrix} m_1 \\ \vdots \\ m_r \end{pmatrix} = A^* \begin{pmatrix} 0 \\ \vdots \\ 0 \end{pmatrix}$$

$$\det(A)I_r \begin{pmatrix} m_1 \\ \vdots \\ m_r \end{pmatrix} = \begin{pmatrix} 0 \\ \vdots \\ 0 \end{pmatrix}$$

This implies $\det(A)m_i = 0$ for every generator m_i . Since any $m \in M$ is an R -linear combination of these generators, $\det(A)m = 0$ for all $m \in M$. $\square$

3.7

Let R be a commutative ring, M a finitely generated R -module, and let J be an ideal of R . Assume $JM = M$. Prove that there exists an element $b \in J$ such that $(1 + b)M = 0$.

Solution.

Since $M = JM$, each generator m_i can be written as $m_i = \sum_{j=1}^r b_{ij}m_j$ with $b_{ij} \in J$. In matrix form, this is $\mathbf{m}^T = B\mathbf{m}^T$, or $(I - B)\mathbf{m}^T = 0$. By Exercise 3.6, $\det(I - B)$ annihilates M .

Expanding the determinant $\det(I - B)$ modulo J , we see that since all entries of B are in J :

$$\det(I - B) \equiv \det(I) \equiv 1 \pmod{J}$$

Thus, $\det(I - B) = 1 + b$ for some $b \in J$. Hence $(1 + b)M = 0$. $\square$

3.8

Let R be a commutative ring, M be a finitely generated R -module, and let J be an ideal of R contained in the Jacobson radical of R (Exercise V.3.14). Prove that $M = 0 \iff JM = M$.

Solution.

$\implies$ If $M = 0$, then $JM = 0 = M$.

$\impliedby$ Assume $JM = M$. By Exercise 3.7, there exists $b \in J$ such that $(1 + b)M = 0$. Since J is contained in the Jacobson radical, $1 + b$ is a unit in R by Exercise V.3.14. Hence, $M = (1 + b)^{-1}(1 + b)M = 0$. $\square$

3.9

Let $(R, \mathfrak{m})$ be a commutative local ring (meaning a ring with a single maximal ideal $\mathfrak{m}$) and M, N be finitely generated R -modules. Prove that if $M = \mathfrak{m}M + N$, then $M = N$.

Solution.

Since M is a finitely generated R -module, its quotient is also finitely generated. For any ideal $I \subseteq R$ and submodule $N \subseteq M$, we have

$$I(M/N) = \frac{IM + N}{N}$$

Applying this to the maximal ideal $\mathfrak{m}$, we have:

$$\mathfrak{m}(M/N) = \frac{\mathfrak{m}M + N}{N} = M/N$$

In a local ring, the Jacobson radical $J(R)$ is equal to the unique maximal ideal $\mathfrak{m}$. Therefore, we have $J(R)(M/N) = M/N$. Since M/N is finitely generated, Nakayama's Lemma implies that $M/N = 0$. Hence, $M = N$. $\square$

3.10

Let R be a commutative local ring, and let M be a finitely generated R -module. Note that $M/\mathfrak{m}M$ is a finite-dimensional vector space over the field $R/\mathfrak{m}$; let $m_1, \dots, m_r \in M$ be elements whose cosets mod $\mathfrak{m}M$ form a basis of $M/\mathfrak{m}M$. Prove that $m_1, \dots, m_r$ generate M .

Solution.

$M/\mathfrak{m}M$ is a vector space over $R/\mathfrak{m}$ by 1.9, and $\dim M/\mathfrak{m}M < +\infty$ since M is finitely generated as an R -module.

Suppose $m \in M$. Since $m_1, \dots, m_r$ is a basis of $M/\mathfrak{m}M$, we have

$$\begin{aligned} m + \mathfrak{m}M &= \sum (s_i + \mathfrak{m})(m_i + \mathfrak{m}M) = \left(\sum s_i m_i \right) + \mathfrak{m}M \\ \implies m &= \sum s_i m_i + m', \quad \text{where } m' \in \mathfrak{m}M \end{aligned}$$

Thus, $M = \langle m_1, \dots, m_r \rangle + \mathfrak{m}M$ and by 3.9

$$M = \langle m_1, \dots, m_r \rangle$$

□

3.16

Every complex $V_\bullet$ determines an element in $K(k\text{-Vect}^f)$ (Grothendieck group), namely

$$\chi_K(V_\bullet) := \sum_i (-1)^i [V_i] \in K(k\text{-Vect}^f).$$

- χ_K 'is an Euler characteristic', in the sense that it satisfies the formula given in Proposition 3.13: $\chi_K(V_\bullet) = \sum_i (-1)^i [H_i(V_\bullet)]$.
- χ_K is a 'universal Euler characteristic', in the following sense. Let G be an abelian group, and let δ be a function associating an element of G to each finite-dimensional vector space, such that $\delta(V) = \delta(V')$ if $V \simeq V'$ and $\delta(V/U) = \delta(V) - \delta(U)$. For $V_\bullet$ a complex, define $\chi_G(V_\bullet) = \sum_i (-1)^i \delta(V_i)$. Then δ induces a (unique) group homomorphism $K(k\text{-Vect}^f) \rightarrow G$ mapping $\chi_K(V_\bullet)$ to $\chi_G(V_\bullet)$.
- In particular, $\delta = \dim$ induces a group homomorphism $K(k\text{-Vect}^f) \rightarrow \mathbb{Z}$ such that $\chi_K(V_\bullet) \mapsto \chi(V_\bullet)$.
- This is in fact an isomorphism.

Solution.

Part 1: χ_K is an Euler characteristic. We prove the formula $\chi_K(V_\bullet) = \sum_i (-1)^i [H_i(V_\bullet)]$ by induction on the length n of the complex $V_\bullet$.

Base case $n = 0$: Consider the complex $0 \rightarrow V_0 \rightarrow 0$. We have $H_0(V_\bullet) = V_0$. Thus:

$$\chi_K(V_\bullet) = [V_0] = [H_0(V_\bullet)]$$

Base case $n = 1$: Consider $0 \rightarrow V_1 \xrightarrow{\alpha} V_0 \rightarrow 0$. The homologies are $H_1(V_\bullet) = \ker \alpha$ and $H_0(V_\bullet) = V_0 / \text{im } \alpha$. We have the short exact sequences:

$$0 \rightarrow \ker \alpha \rightarrow V_1 \rightarrow \text{im } \alpha \rightarrow 0 \implies [V_1] = [\ker \alpha] + [\text{im } \alpha]$$

$$0 \rightarrow \operatorname{im} \alpha \rightarrow V_0 \rightarrow V_0 / \operatorname{im} \alpha \rightarrow 0 \implies [V_0] = [\operatorname{im} \alpha] + [V_0 / \operatorname{im} \alpha]$$

Computing the Euler characteristic:

$$\begin{aligned} \chi_K(V_\bullet) &= [V_0] - [V_1] = ([\operatorname{im} \alpha] + [V_0 / \operatorname{im} \alpha]) - ([\ker \alpha] + [\operatorname{im} \alpha]) = [V_0 / \operatorname{im} \alpha] - [\ker \alpha] \\ &= [H_0(V_\bullet)] - [H_1(V_\bullet)] = \sum_{i=0}^1 (-1)^i [H_i(V_\bullet)] \end{aligned}$$

Inductive step: Assume the statement holds for complexes of length n . Consider a complex $V_\bullet$ of length $n+1$:

$$0 \rightarrow V_{n+1} \xrightarrow{\alpha_{n+1}} V_n \xrightarrow{\alpha_n} \dots \xrightarrow{\alpha_1} V_0 \rightarrow 0$$

Let $V'_\bullet$ be its truncated complex $0 \rightarrow V_n \xrightarrow{\alpha_n} \dots \rightarrow V_0 \rightarrow 0$. By the induction hypothesis, $\chi_K(V'_\bullet) = \sum_{i=0}^n (-1)^i [H_i(V'_\bullet)]$. By definition, $\chi_K(V_\bullet) = \chi_K(V'_\bullet) + (-1)^{n+1} [V_{n+1}]$. Let us compare the homologies of $V_\bullet$ and $V'_\bullet$:

- $H_i(V_\bullet) = H_i(V'_\bullet)$ for all $i < n$.
- $H_n(V'_\bullet) = \ker \alpha_n$.
- $H_n(V_\bullet) = \ker \alpha_n / \operatorname{im} \alpha_{n+1}$.
- $H_{n+1}(V_\bullet) = \ker \alpha_{n+1}$.

The exact sequence $0 \rightarrow \ker \alpha_{n+1} \rightarrow V_{n+1} \rightarrow \operatorname{im} \alpha_{n+1} \rightarrow 0$ gives $[V_{n+1}] = [\ker \alpha_{n+1}] + [\operatorname{im} \alpha_{n+1}]$. The exact sequence $0 \rightarrow \operatorname{im} \alpha_{n+1} \rightarrow \ker \alpha_n \rightarrow \ker \alpha_n / \operatorname{im} \alpha_{n+1} \rightarrow 0$ gives $[\ker \alpha_n] = [\operatorname{im} \alpha_{n+1}] + [\ker \alpha_n / \operatorname{im} \alpha_{n+1}]$, which rewrites as $[H_n(V'_\bullet)] = [\operatorname{im} \alpha_{n+1}] + [H_n(V_\bullet)]$.

Now, sum the alternating homologies for $V_\bullet$:

$$\sum_{i=0}^{n+1} (-1)^i [H_i(V_\bullet)] = \sum_{i=0}^{n-1} (-1)^i [H_i(V'_\bullet)] + (-1)^n [H_n(V_\bullet)] + (-1)^{n+1} [H_{n+1}(V_\bullet)]$$

Substitute $[H_n(V_\bullet)] = [H_n(V'_\bullet)] - [\operatorname{im} \alpha_{n+1}]$ and $[H_{n+1}(V_\bullet)] = [\ker \alpha_{n+1}]$:

$$\begin{aligned} &= \sum_{i=0}^{n-1} (-1)^i [H_i(V'_\bullet)] + (-1)^n ([H_n(V'_\bullet)] - [\operatorname{im} \alpha_{n+1}]) + (-1)^{n+1} [\ker \alpha_{n+1}] \\ &= \sum_{i=0}^n (-1)^i [H_i(V'_\bullet)] + (-1)^{n+1} ([\operatorname{im} \alpha_{n+1}] + [\ker \alpha_{n+1}]) \\ &= \chi_K(V'_\bullet) + (-1)^{n+1} [V_{n+1}] = \chi_K(V_\bullet) \end{aligned}$$

This completes the induction.

Part 2: χ_K is a universal Euler characteristic. Recall that $K(k\text{-Vect})^f = F/E$, where F is the free abelian group on isomorphism classes of finite-dimensional vector spaces, and E is the subgroup generated by relations from short exact sequences. Define a set-map from the generators of F to G via $[V] \mapsto \delta(V)$. Because $\delta(V) = \delta(V')$ whenever $V \simeq V'$, this map is correctly defined and extends linearly to a unique group homomorphism $\varphi : F \rightarrow G$.

If $0 \rightarrow U \rightarrow V \rightarrow W \rightarrow 0$ is exact, then $W \simeq V/U$. Thus:

$$\varphi([V] - [U] - [W]) = \delta(V) - \delta(U) - \delta(V/U) = 0$$

Hence, the subgroup $E \subset \ker \varphi$. By the universal property of quotients, φ factors through to a unique induced morphism $\varphi^* : K(k\text{-Vect}^f) \rightarrow G$ such that $\varphi^*([V]) = \delta(V)$. Applying this to a complex $V_\bullet$:

$$\varphi^*(\chi_K(V_\bullet)) = \varphi^*\left(\sum_i (-1)^i [V_i]\right) = \sum_i (-1)^i \varphi^*([V_i]) = \sum_i (-1)^i \delta(V_i) = \chi_G(V_\bullet)$$

To see φ^* is unique, suppose $\sigma : K(k\text{-Vect}^f) \rightarrow G$ also maps $\chi_K(V_\bullet)$ to $\chi_G(V_\bullet)$ for all complexes. Consider the trivial complex $0 \rightarrow V \rightarrow 0$. Here $\chi_K(V_\bullet) = [V]$ and $\chi_G(V_\bullet) = \delta(V)$. This forces $\sigma([V]) = \delta(V)$, so $\sigma = \varphi^*$.

Part 3 & 4: $\delta = \dim$ induces an isomorphism to $\mathbb{Z}$. Since $\dim(V/U) = \dim V - \dim U$ and dimension is an isomorphism invariant, we apply Part 2. The function $\delta = \dim$ induces a group homomorphism $\varphi : K(k\text{-Vect}^f) \rightarrow \mathbb{Z}$ mapping $\chi_K(V_\bullet) \mapsto \chi(V_\bullet)$. We show φ is an isomorphism.

First, observe a key property in $K(k\text{-Vect}^f)$. For any spaces V, U , there is an exact sequence

$$0 \rightarrow U \rightarrow V \oplus U \rightarrow V \rightarrow 0$$

It follows that $[V \oplus U] = [V] + [U]$. For any n -dimensional vector space V , $V \simeq k^{\oplus n}$. Therefore, $[V] = n[k] = (\dim V)[k]$.

Surjectivity: The morphism φ maps $[V] \mapsto \dim V$. It is clearly surjective onto $\mathbb{Z}$, as choosing $V = k^{\oplus n}$ yields n .

Injectivity: Let $x \in K(k\text{-Vect}^f)$ be any element. We can write x as a finite linear combination $\sum \lambda_i [V_i]$ for some $\lambda_i \in \mathbb{Z}$. Suppose $x \in \ker \varphi$. Then:

$$0 = \varphi\left(\sum \lambda_i [V_i]\right) = \sum \lambda_i \varphi([V_i]) = \sum \lambda_i \dim V_i$$

Using the identity $[V_i] = (\dim V_i)[k]$ derived above, we rewrite our element x in K :

$$\sum \lambda_i [V_i] = \sum \lambda_i (\dim V_i \cdot [k]) = \left(\sum \lambda_i \dim V_i\right) [k] = 0 \cdot [k] = 0$$

Thus, $\ker \varphi = \{0\}$, meaning φ is injective. Therefore, φ is an isomorphism. $\square$

§4 Presentations and resolutions

4.3

Prove that an integral domain R is a PID if and only if every submodule of R itself is free.

Solution.

$\implies$ Let $I \subset R$ be a submodule. By definition, submodules of R are exactly the ideals of R . Since R is a PID, $I = (a)$ for some $a \in R$. If $I = 0$, it is free (with an empty basis). If $I \neq 0$, the map $R \rightarrow I$ given by $r \mapsto ra$ is an isomorphism of R -modules. Alternatively, the set $\{a\}$ is linearly independent since R is an integral domain ($ra = 0 \implies r = 0$ for $a \neq 0$), and it generates I . Thus, it is a basis of I , which implies $I \simeq R^1$.

$\Leftarrow$ Suppose $I \subset R$ is an ideal. By assumption, it is free as an R -module. Every basis of I would be a linearly independent system in R . However, any two non-zero elements $x, y \in R$ are linearly dependent over R , since $y \cdot x - x \cdot y = 0$. Therefore, the maximum size of any linearly independent set in R is 1.

This means $\text{rk } I \leq 1$. Hence, the basis is either empty ($I = 0$) or consists of a single element ($I = (a)$ for some $a \in R$). Since every ideal is generated by at most one element, R is a PID. $\square$

4.4

Let R be a commutative ring and M an R -module.

- Prove that $\text{Ann}(M)$ is an ideal of R .
- If R is an integral domain and M is finitely generated, prove that M is torsion if and only if $\text{Ann}(M) \neq 0$.
- Give an example of a torsion module M over an integral domain, such that $\text{Ann}(M) = 0$. (Of course this example cannot be finitely generated!)

Solution.

$$\text{Ann}(M) = \{r \in R \mid rm = 0 \quad \forall m \in M\}$$

$$a \in R, r \in \text{Ann}(M) \implies \forall m \in M, \quad (ar)m = a(rm) = a0 = 0.$$

$$\forall r, r' \in \text{Ann}(M), \forall m \in M, \quad (r + r')m = rm + r'm = 0 + 0 = 0.$$

More is true: $\forall m \in M$ $\text{Ann}(m)$ is an ideal of R and $\text{Ann}(M) = \bigcap_{m \in M} \text{Ann}(m)$

Note that if R is an integral domain and M is finitely generated, there is a surjective morphism $R^{\oplus A} \rightarrow M$, $A = \{a_1, \dots, a_n\}$, hence $M \simeq R^{\oplus A}/N$ for some submodule N .

$\Leftarrow$ If $\text{Ann}(M) \neq 0$, then $\exists r \in R \mid r \neq 0, \quad rm = 0 \quad \forall m \in M \implies \forall m \in M, \quad m \text{ is torsion.}$

$\implies$ If M is torsion, then $\exists r_i \in R, \quad i = 1, \dots, n \mid r_i \bar{a}_i = \bar{0}. \implies \forall m \in M, \quad m = \sum p_i \bar{a}_i \implies rm = 0$, where $r = \prod r_i \neq 0 \implies \text{Ann}(M) \neq 0$.

$\mathbb{Q}$ is not finitely generated as a $\mathbb{Z}$ -module. $\mathbb{Z} \subset \mathbb{Q}$ is a submodule. $\mathbb{Q}/\mathbb{Z}$ as a $\mathbb{Z}$ -module is torsion:

$$q \left(\frac{p}{q} + \mathbb{Z} \right) = p + \mathbb{Z} = \mathbb{Z}$$

If $n \in \mathbb{Z}$ such that $\forall \bar{s} \in \mathbb{Q}/\mathbb{Z}, \quad n\bar{s} = \bar{0}$, then $\forall m \in \mathbb{N}, \quad n \left(\frac{1}{m} + \mathbb{Z} \right) = \frac{n}{m} + \mathbb{Z} = \mathbb{Z} \implies m \mid n \quad \forall m \in \mathbb{N} \implies n = 0$.

Hence, $\text{Ann}_{\mathbb{Z}}(\mathbb{Q}/\mathbb{Z}) = 0$. □

4.5

Let M be a module over a commutative ring R . Prove that an ideal I of R is the annihilator of an element of M if and only if M contains an isomorphic copy of R/I (viewed as an R -module).

The *associated primes* of M are the prime ideals among the ideals $\text{Ann}(m)$, for $m \in M$. The set of the associated primes of a module M is denoted $\text{Ass}_R(M)$. Note that every prime in $\text{Ass}_R(M)$ contains $\text{Ann}_R(M)$.

Solution.

$\implies$ If $I = \text{Ann}(m)$ is an ideal, then $\ker \varphi = I$, where $\varphi : R \rightarrow M$ defined by $\varphi(r) = rm$ is an R -module homomorphism. Hence, we have an inclusion $R/I \simeq \text{im } \varphi \hookrightarrow M$.

$\Leftarrow$ If there is an inclusion $\varphi : R/I \hookrightarrow M$, then consider $m = \varphi(\bar{1})$. If $rm = 0$, then

$$0 = r\varphi(\bar{1}) = \varphi(r\bar{1}) = \varphi(\bar{r}) \implies \bar{r} = \bar{0} \quad (\text{since } \ker \varphi = 0) \implies r \in I$$

This argument works in the reverse direction as well, so $I = \text{Ann}(m)$.

Now, suppose that $\text{Ann}(m)$ is an associated prime. We need to show that $\text{Ann}(m) \supset \text{Ann}(M)$. It is obvious, since

$$\text{Ann}(M) = \bigcap_{m \in M} \text{Ann}(m)$$

□

4.6

Let M be a module over a commutative ring R , and consider the family of ideals $\text{Ann}(m)$, as m ranges over the nonzero elements of M . Prove that the maximal elements in this family are prime ideals of R . Conclude that if R is Noetherian, then $\text{Ass}_R(M) \neq \emptyset$.

Solution.

Note that $\forall r \in R, \quad \text{Ann}(m) \subset \text{Ann}(rm)$, for if $sm = 0$, then $s(rm) = r(sm) = 0$.

Suppose $\text{Ann}(m)$ is maximal among annihilators and that $rs \in \text{Ann}(m)$. If $r \notin \text{Ann}(m)$, then $rm \neq 0$.

$s(rm) = 0 \implies s \in \text{Ann}(rm)$. Since $\text{Ann}(m)$ is maximal and $\text{Ann}(m) \subset \text{Ann}(rm)$ (with $rm \neq 0$), we must have $\text{Ann}(m) = \text{Ann}(rm)$. Hence, $s \in \text{Ann}(m)$ and $\text{Ann}(m)$ is prime.

The family of ideals $\mathcal{F} = \{\text{Ann}(m) \mid m \in M \setminus \{0\}\}$ is non-empty.

In a Noetherian ring, every non-empty family of ideals has a maximal element with respect to inclusion. Let P be a maximal element of $\mathcal{F}$. By the argument above, P is a prime ideal. Because $P \in \mathcal{F}$, $P = \text{Ann}(m)$ for some non-zero $m \in M$, meaning P is an associated prime of M .

Therefore, $\text{Ass}_R(M) \neq \emptyset$. □

4.13

Let R be a commutative ring. A tuple $(a_1, a_2, \dots, a_n)$ of elements of R is a *regular sequence* if a_1 is a non-zero-divisor in R , a_2 is a non-zero-divisor modulo (a_1) , a_3 is a non-zero-divisor modulo (a_1, a_2) , and so on.

For a, b in R , consider the following complex of R -modules:

$$(*) \quad 0 \rightarrow R \xrightarrow{d_2} R \oplus R \xrightarrow{d_1} R \xrightarrow{\pi} \frac{R}{(a, b)} \rightarrow 0$$

where π is the canonical projection, $d_1(r, s) = ra + sb$, and $d_2(t) = (bt, -at)$. Put otherwise, d_1 and d_2 correspond, respectively, to the matrices

$$\begin{pmatrix} a & b \end{pmatrix}, \quad \begin{pmatrix} b \\ -a \end{pmatrix}.$$

- Prove that this is indeed a complex, for every a and b .
- Prove that if (a, b) is a regular sequence, this complex is *exact*.

The complex $(*)$ is called the *Koszul complex* of (a, b) . Thus, when (a, b) is a regular sequence, the Koszul complex provides us with a free resolution of the module $R/(a, b)$.

Solution.

$$[d_1 \circ d_2] = \begin{pmatrix} a & b \end{pmatrix} \begin{pmatrix} b \\ -a \end{pmatrix} = 0, \quad \text{so } d_1 \circ d_2 = 0$$

$\pi \circ d_1 = 0$ is obvious, and canonical projection is an epimorphism.

If (a, b) is regular and $d_2(t) = 0$, then

$$(bt, -at) = (0, 0) \implies at = 0 \implies t = 0$$

since a is a non-zero-divisor. Hence, $\ker d_2 = 0$.

If $d_1(r, s) = ra + sb = 0$, then consider this equation modulo (a) :

$$\bar{s}\bar{b} = \bar{0} \implies \bar{s} = \bar{0}$$

for $\bar{b}$ is a non-zero-divisor.

$$\implies s = at \quad \text{for some } t \in R.$$

Then

$$0 = ra + atb = a(r + tb) \implies r = -tb$$

Thus, $(r, s) = (-bt, at) = d_2(-t) \in \text{im } d_2 \implies \ker d_1 = \text{im } d_2$.

Finally, $\ker \pi = (a, b) = \text{im } d_1$, even without regularness, and π is an epimorphism. $\square$

4.16

Let $\varphi : R^n \rightarrow R^m$ and $\psi : R^p \rightarrow R^q$ be two R -module homomorphisms, and let

$$\varphi \oplus \psi : R^n \oplus R^p \rightarrow R^m \oplus R^q$$

be the morphism induced on direct sums. Prove that

$$\text{coker}(\varphi \oplus \psi) = \text{coker } \varphi \oplus \text{coker } \psi.$$

Solution.

$$\begin{array}{ccccccc}
 R^n & \xrightarrow{\varphi} & R^m & \xrightarrow{\varphi'} & \text{coker } \varphi & & \\
 \downarrow i_1 & & \downarrow i'_1 & & \downarrow i''_1 & \searrow \sigma_1 & \\
 R^n \oplus R^p & \xrightarrow{\varphi \oplus \psi} & R^m \oplus R^q & \xrightarrow{\tau} & \text{coker } \varphi \oplus \text{coker } \psi & \xrightarrow{\lambda} & M \\
 \uparrow i_2 & & \uparrow i'_2 & & \uparrow \tau^* & \nearrow \sigma_2 & \\
 R^p & \xrightarrow{\psi} & R^q & \xrightarrow{\psi'} & \text{coker } \psi & &
 \end{array}$$

Let $v \in R^n$, $w \in R^p$.

$$\tau \circ (\varphi \oplus \psi) \circ i_1(v) = i_1'' \circ \varphi' \circ \varphi(v) = i_1''(0) = (0, 0)$$

Similarly, $\tau \circ (\varphi \oplus \psi) \circ i_2 = 0$.

Hence, $\forall (v, w) \in R^n \oplus R^p$,

$$\begin{aligned} \tau \circ (\varphi \oplus \psi)(v, w) &= \tau \circ (\varphi \oplus \psi)(v, 0) + \tau \circ (\varphi \oplus \psi)(0, w) \\ &= \tau \circ (\varphi \oplus \psi) \circ i_1(v) + \tau \circ (\varphi \oplus \psi) \circ i_2(w) = (0, 0), \end{aligned}$$

so τ indeed annihilates $\varphi \oplus \psi$.

Moreover, if $\tau^* : R^m \oplus R^q \rightarrow M$ such that $\tau^* \circ (\varphi \oplus \psi) = 0$, then

$$\tau^* \circ i_2' \circ \psi = \tau^* \circ (\varphi \oplus \psi) \circ i_2 = 0$$

Since $\tau^* \circ i_2'$ annihilates ψ , it induces a unique $\sigma_2 : \text{coker } \psi \rightarrow M$ such that

$$\tau^* \circ i_2' = \sigma_2 \circ \psi'$$

Similarly, we obtain $\sigma_1 : \text{coker } \varphi \rightarrow M$ such that $\tau^* \circ i_1' = \sigma_1 \circ \varphi'$.

Together σ_1 and σ_2 induce $\lambda : \text{coker } \varphi \oplus \text{coker } \psi \rightarrow M$ such that

$$\sigma_1 = \lambda \circ i_1'', \quad \sigma_2 = \lambda \circ i_2''$$

We have then

$$\lambda \circ \tau \circ i_2' = \lambda \circ i_2'' \circ \psi' = \sigma_2 \circ \psi' = \tau^* \circ i_2'$$

and similarly $\lambda \circ \tau \circ i_1' = \tau^* \circ i_1'$.

It follows that $\forall (v, w) \in R^m \oplus R^q$,

$$\begin{aligned} \lambda \circ \tau(v, w) &= \lambda \circ \tau \circ i_1'(v) + \lambda \circ \tau \circ i_2'(w) \\ &= \tau^* \circ i_1'(v) + \tau^* \circ i_2'(w) \\ &= \tau^*(v, w) \implies \tau^* = \lambda \circ \tau \end{aligned}$$

If λ^* is another morphism such that $\tau^* = \lambda^* \circ \tau$, then $\sigma_2 \circ \psi' = \tau^* \circ i_2' = \lambda^* \circ \tau \circ i_2' = \lambda^* \circ i_2'' \circ \psi'$.

Since ψ' is an epimorphism, this would imply $\sigma_2 = \lambda^* \circ i_2''$. Similarly, we get $\sigma_1 = \lambda^* \circ i_1''$, and we must have $\lambda^* = \lambda$.

So, $\text{coker } \varphi \oplus \text{coker } \psi$ satisfies the universal property for $\text{coker}(\varphi \oplus \psi)$. □

§5 Classification of finitely generated modules over PIDs

5.5

Recall (Exercise V.4.11) that a commutative ring is local if it has a single maximal ideal $\mathfrak{m}$. Let R be a local ring, and let M be a direct summand of a finitely generated free R -module: that is, there exists an R -module N such that $M \oplus N$ is a free R -module.

- Choose elements $m_1, \dots, m_r \in M$ whose cosets mod $\mathfrak{m}M$ are a basis of $M/\mathfrak{m}M$ as a vector space over the field $R/\mathfrak{m}$. By Nakayama's lemma, $M = \langle m_1, \dots, m_r \rangle$ (Exercise 3.10).
- Obtain a surjective homomorphism $\pi : F = R^{\oplus r} \rightarrow M$.
- Show that π splits, giving an isomorphism $F \simeq M \oplus \ker \pi$. (Apply Exercise III.6.9 to the surjective homomorphism π and the free module $M \oplus N$ to obtain a splitting $M \rightarrow F$; then use Proposition III.7.5.)
- Show $\ker \pi / \mathfrak{m} \ker \pi = 0$. Use Nakayama's lemma (Exercise 3.8) to deduce that $\ker \pi = 0$.
- Conclude that $M \simeq F$ is in fact free.

Solution.

First point is already proved in 3.10.

Second point is obvious.

By III.6.9, there is a morphism $\varphi : M \oplus N \rightarrow F$ such that the diagram

$$\begin{array}{ccc} & M \oplus N & \\ \varphi \swarrow & & \searrow \pi_1 \\ F & \xrightarrow{\pi} & M \end{array}$$

commutes. Define $\tau : M \rightarrow F$ by $\tau = \varphi \circ i_1$. We have $\pi \circ \tau = \pi \circ \varphi \circ i_1 = \pi_1 \circ i_1 = \text{id}$, so τ is right-inverse for π and $F = M \oplus \ker \pi$.

Now I claim that

$$F/\mathfrak{m}F \simeq M/\mathfrak{m}M \oplus \ker \pi / \mathfrak{m} \ker \pi$$

as $R/\mathfrak{m}$ -vector spaces and

$$\dim F/\mathfrak{m}F = \dim M/\mathfrak{m}M = r$$

It follows then that $\ker \pi / \mathfrak{m} \ker \pi = 0$, as desired.

To verify my claim, consider an R -module homomorphism

$$\begin{aligned} \psi : F &\rightarrow M/\mathfrak{m}M \oplus \ker \pi / \mathfrak{m} \ker \pi \\ (m, k) &\mapsto (m + \mathfrak{m}M, k + \mathfrak{m} \ker \pi) \end{aligned}$$

If $\psi(m, k) = 0$, then $m \in \mathfrak{m}M$, $k \in \mathfrak{m} \ker \pi$, hence for some $r_i, r'_i \in \mathfrak{m}$, $m'_i \in M$, $k_i \in \ker \pi$,

$$(m, k) = \left(\sum r'_i m'_i, \sum r_i k_i \right) = \sum r'_i (m'_i, 0) + \sum r_i (0, k_i) \in \mathfrak{m}F$$

On the other hand, if $f \in \mathfrak{m}F$, then

$$f = \sum r_i f_i = \sum r_i (m'_i, k_i) = \left(\sum r_i m'_i, \sum r_i k_i \right)$$

and $\psi(f) = 0$. Since ψ is clearly surjective, we have an induced isomorphism of R -modules

$$\psi^* : F/\mathfrak{m}F \rightarrow M/\mathfrak{m}M \oplus \ker \pi / \mathfrak{m} \ker \pi$$

Moreover, ψ^* is a homomorphism of $R/\mathfrak{m}$ -vector spaces. Indeed

$$\psi^*((r + \mathfrak{m})\bar{f}) = \psi^*(r\bar{f}) = r\psi^*(\bar{f}) = (r + \mathfrak{m})\psi^*(\bar{f})$$

and ψ^* also preserves sum as an R -module homomorphism. Consequently,

$$F/\mathfrak{m}F \simeq M/\mathfrak{m}M \oplus \ker \pi / \mathfrak{m} \ker \pi$$

as $R/\mathfrak{m}$ vector spaces.

Let $e_1, \dots, e_r$ be a basis of F . The corresponding classes generate $F/\mathfrak{m}F$ over $R/\mathfrak{m}$ and if $\sum (r_i + \mathfrak{m})\bar{e}_i = \bar{0}$, then $\sum r_i e_i \in \mathfrak{m}F$, which implies $r_i \in \mathfrak{m}$ and $r_i + \mathfrak{m} = 0 + \mathfrak{m}$. Hence, $\bar{e}_1, \dots, \bar{e}_r$ are linearly independent in $F/\mathfrak{m}F$ over $R/\mathfrak{m}$ and $\dim F/\mathfrak{m}F = r$.

As discussed above, we have now $\ker \pi / \mathfrak{m} \ker \pi = 0$ as $R/\mathfrak{m}$ -vector space, and hence as an R -module. It implies that $\ker \pi = \mathfrak{m} \ker \pi$ and $\ker \pi = 0$ by Nakayama's lemma ($\ker \pi \simeq F/M$ is finitely generated).

Thus, π is an isomorphism and $M \simeq R^r$. □

5.6

Let R be an integral domain, and let $M = \langle m_1, \dots, m_r \rangle$ be a finitely generated module. Prove that $\text{rk } M \leq r$.

Solution.

Since M is generated by r elements, there is a surjective homomorphism $R^r \rightarrow M$, meaning $M \simeq R^r/N$ for some submodule N . Every linear relation in R^r reduces modulo N to a relation in M , so there can't be more than r linearly independent elements in M . □

5.7

Let R be an integral domain, and let M be a finitely generated module over R . Prove that $\text{rk } M = \text{rk}(M/\text{Tor}(M))$.

Solution.

If a system of elements in M is linearly dependent, then its image in $M/\text{Tor}(M)$ under canonical projection is also linearly dependent. Hence, $\text{rk } M \geq \text{rk}(M/\text{Tor}(M))$.

For the reverse direction, consider $\overline{m_1}, \dots, \overline{m_r} \in M/\text{Tor}(M)$ such that

$$\sum r_i \overline{m_i} = \overline{0}$$

It follows that $\sum r_i m_i \in \text{Tor}(M)$, so there is some $s \in R \setminus \{0\}$ such that $\sum sr_i m_i = 0$, meaning that $\{m_1, \dots, m_r\}$ is linearly dependent in M . Hence, $\text{rk } M \leq \text{rk}(M/\text{Tor}(M))$. $\square$

5.8

Let R be an integral domain, and let M be a finitely generated module over R . Prove that $\text{rk } M = r$ if and only if M has a free submodule $N \simeq R^r$, such that M/N is torsion.

If R is a PID, then N may be chosen so that $0 \rightarrow N \rightarrow M \rightarrow M/N \rightarrow 0$ splits.

Solution.

$\Rightarrow$ Suppose $m_1, \dots, m_r$ are linearly independent. They generate

$$N = \langle m_1, \dots, m_r \rangle \simeq R^r$$

If $\overline{m} \in M/N$, $\overline{m} \neq \overline{0}$, then $m_1, \dots, m_r, m$ are linearly dependent $\Rightarrow \exists r \neq 0, r_i$ not all zeros, such that $\sum r_i m_i + rm = 0$. $\Rightarrow r\overline{m} = -\sum r_i \overline{m_i} = \overline{0}$, so M/N is torsion.

$\Leftarrow$ Let N be a submodule of M such that $N \simeq R^r$, M/N is torsion, and let $n_1, \dots, n_r$ be its basis.

$n_1, \dots, n_r$ is a maximal linearly independent system of M . Indeed, if $m \in M$, then $\exists r \in R$ such that $rm \in N$, since M/N is torsion. $rm = \sum r_i n_i$, so $n_1, \dots, n_r, m$ is linearly dependent.

Now if $m_1, \dots, m_s$ is linearly independent, then $s \leq r$, so $\text{rk } M = r$.

Suppose now R is a PID and let K be a finitely generated torsion-free R -module. Let us show that K is free. K has a free submodule $F \simeq R^s$, such that K/F is torsion. By 4.4, $\exists a \in \text{Ann}(K/F)$ such that $a \neq 0$. The morphism $\varphi : K \rightarrow F$, $k \mapsto ak$ is injective, since K is torsion-free, and identifies K with a submodule of F

$$K \simeq \text{im } \varphi \subset F$$

Since over a PID every submodule of a free module is free, we conclude that K is free.

If M is a finitely generated module over a PID R with $\text{rk } M = r$, then $M/\text{Tor}(M)$ is torsion-free, hence free, and by 5.7

$$\text{rk}(M/\text{Tor}(M)) = \text{rk } M = r \Rightarrow M/\text{Tor}(M) \simeq R^r$$

Let $m_1, \dots, m_r \in M$ be elements whose classes form a basis of $M/\text{Tor}(M)$ and consider

$$N = \langle m_1, \dots, m_r \rangle \simeq R^r$$

It follows that $N + \text{Tor}(M) = M \implies M \simeq N \oplus \text{Tor}(M)$, since $N \cap \text{Tor}(M) = 0$.

Thus, $M/N \simeq \text{Tor}(M)$ is torsion and

$$0 \rightarrow N \rightarrow M \rightarrow M/N \rightarrow 0$$

splits. □

5.9

Let R be an integral domain, and let

$$0 \longrightarrow M_1 \longrightarrow M_2 \longrightarrow M_3 \longrightarrow 0$$

be an exact sequence of finitely generated R -modules. Prove that $\text{rk } M_2 = \text{rk } M_1 + \text{rk } M_3$.

Deduce that ‘rank’ defines a homomorphism from the Grothendieck group of the category of finitely generated R -modules to $\mathbb{Z}$

Solution.

Denote $\text{rk } M_i = r_i$ and suppose that $\{m_1, \dots, m_{r_1}\}$ and $\{m''_1, \dots, m''_{r_3}\}$ are linearly independent systems in M_1 and M_3 respectively. Let us denote the morphisms in our sequence by λ and λ' .

Consider elements $m'_1, \dots, m'_{r_3} \in M_2$ such that $\lambda'(m'_k) = m''_k$. Then the system

$$\lambda(m_1), \dots, \lambda(m_{r_1}), m'_1, \dots, m'_{r_3}$$

is linearly independent. Indeed, if $\sum s_k \lambda(m_k) + \sum s'_k m'_k = 0$, then

$$0 = \lambda' \left(\sum s_k \lambda(m_k) + \sum s'_k m'_k \right) = \sum s'_k \lambda'(m'_k) = \sum s'_k m''_k \implies s'_k = 0,$$

since $m''_1, \dots, m''_{r_3}$ is linearly independent. We have now

$$\lambda \left(\sum s_k m_k \right) = \sum s_k \lambda(m_k) = 0 \implies \sum s_k m_k = 0,$$

since λ is an inclusion $\implies s_k = 0$.

Now if $m' \in M_2$, then there exist $s'' \neq 0$ and s''_k for $1 \leq k \leq r_3$, such that

$$s'' \lambda'(m') + \sum s''_k \lambda'(m'_k) = 0$$

Since $\ker \lambda' = \text{im } \lambda$, it follows that there exists $m \in M_1$ such that

$$\lambda(m) = s'' m' + \sum s''_k m'_k$$

On the other hand, there exist $s \neq 0$ and s_k for $1 \leq k \leq r_1$, such that $sm + \sum s_k m_k = 0$. We have then

$$ss'' m' + \sum ss''_k m'_k + \sum s_k \lambda(m_k) = 0$$

Because R is an integral domain and $s \neq 0, s'' \neq 0$, their product $ss'' \neq 0$. Thus,

$$\lambda(m_1), \dots, \lambda(m_{r_1}), m'_1, \dots, m'_{r_3}$$

is a maximal linearly independent system in M_2 , and hence every other linearly independent system in M_2 has $\leq r_1 + r_3$ elements, which concludes the proof. $\square$

5.13

Let M be a finitely generated module over an integral domain R . Prove that if R is a PID, then M is torsion-free if and only if it is free. Prove that this property characterizes PIDs. (Cf. Exercise 4.3.)

Solution.

If M is torsion-free, then it is free as shown in 5.8.

If M is free with basis $b_1, \dots, b_r$ and has a torsion m , so that $rm = 0, r \neq 0$, then

$$\begin{aligned} 0 = rm &= r \sum r_i b_i = \sum rr_i b_i \implies rr_i = 0 \quad \forall i \\ &\implies r_i = 0 \quad \forall i \quad \text{and} \quad m = 0, \end{aligned}$$

so M is torsion-free.

Now we want to show that if a ring R has a property that for any finitely generated R -module M ,

$$M \text{ is torsion-free} \iff M \text{ is free}$$

Then R is a PID.

Indeed, R is free as an R -module: $\{1\}$ is a linearly independent system, for if $r \cdot 1 = 0$, then $r = 0$; and $R = (1)$. Consequently, R is torsion-free, so it is an integral domain.

Consider an ideal $I \subset R$. Since R is an integral domain, I is torsion-free, hence free with $\text{rk } I \leq 1$. It follows that $I = (a)$ for some a . $\square$