




PERSONAL HEALTH SMART REPORT

A comprehensive analysis of your health using
Blood, Physicals, and Health Questionnaire data



Prepared for

ARVIND

Basic Info

Male /41 Yrs

Patient ID

HYD572178

Report released on

26/06/2025

Date of Test

25/06/2025




Table of contents

Your smart report includes the following sections.

S. No.	Section	Page No
01	Summary for Doctors	03
02	Your Wellbeing Index	07
03	Glance of Important Parameters	08
04	Wellness Recommendations	15
05	References	16
06	Lab Report	

Get personalized insights & trends of your lab reports

Now on Tata 1mg app!

View trends & insights



Scan QR to view

Disclaimer


- This is an electronically generated report and is not a substitute for medical advice.
- While following the recommendations, please be careful of any allergies or intolerances.
- If you are pregnant or lactating, some of the recommendations and analyzed information in the Smart Report may not directly apply to you. Please consult a doctor regarding your test results and recommendations.
- Analysis uses the attached blood test report and Well Being Index Questionnaire data, if present, and urine analysis report, if present.
- Tata 1mg is not liable for any direct, indirect, special, consequential, or other damages. This report cannot be used for any medico-legal purposes. Partial reproduction of the test results is not permitted. Also, TATA 1mg Labs is not responsible for any misinterpretation or misuse of the information.

Doctor Summary For

Comprehensive Platinum Full Body Checkup with Smart Report

For
Arvind
Male /41 Yrs

Note This is an electronically generated summary of the attached report. It is advised to read this summary in conjunction with the attached report and to correlate it clinically. For the trends section, the out of range values are highlighted with respect to the bio reference range of respective reports.

Test Name	Result, 25/06/25	Bio. Ref. Interval	Trends (For last three tests)			
Complete Blood Count			Date 1	Date 2	Date 3	
Hemoglobin	16.2 g/dL	13.0 - 17.0	<div></div> <p>We don't have any of your previous lab results for these tests in our records</p>			
RBC	5.26 10^6/cu.mm	4.5 - 5.5				
HCT	48.9 %	40 - 50				
Total Leucocyte Count	8.38 10^3/ÂµL	4 - 10				
Neutrophils	55.1 %	40 - 80				
Lymphocytes	31.1 %	20 - 40				
Monocytes	▲ 10.1 %	2 - 10				
Eosinophils	3.1 %	1 - 6				
Basophils	0.6 %	0 - 2				
Platelet Count	319 10^3/ÂµL	150 - 410				
PDW	▲ 17.5 fL	9 - 17				
Inflammatory markers						
Erythrocyte Sedimentation Rate	4 mm/hr	0 - 10				
C-Reactive Protein (Quantitative)	5.10 mg/L	0.1 - 10				
Iron Studies						
Iron Serum	84 Âµg/dL	65 - 175				
Total Iron Binding Capacity (TIBC)	442.88 Âµg/dL	250 - 460				
Unsaturated Iron Binding Capacity	▲ 359 Âµg/dL	111 - 343				
Ferritin	220.00 ng/mL	22 - 322				
Diabetes Profile						
Glycosylated Hemoglobin (HbA1c)	5.3 %	4 - 5.6				
Glucose - Fasting	80 mg/dL	70 - 99				



We don't have any of your previous lab results for these tests in our records

Doctor Summary For
Comprehensive Platinum Full Body Checkup with Smart Report

For
Arvind
Male /41 Yrs

Note This is an electronically generated summary of the attached report. It is advised to read this summary in conjunction with the attached report and to correlate it clinically. For the trends section, the out of range values are highlighted with respect to the bio reference range of respective reports.

Test Name	Result, 25/06/25	Bio. Ref. Interval	Trends (For last three tests)		
Diabetes Profile			Date 1	Date 2	Date 3
Microalbumin-Albumin	6.07 mg/L	0 - 29.99			
Microalbumin-Albumin/Creatinine Ratio	9.58 mg/g Creatinine	0 - 29.99			
Kidney Function Test					
Blood Urea Nitrogen	▼ 8 mg/dL	9 - 23			
Urea	▼ 16.24 mg/dL	19.26 - 49.22			
Creatinine	0.88 mg/dL	0.7 - 1.3			
Uric Acid	6.4 mg/dL	3.5 - 7.2			
Sodium	142 mEq/L	132 - 146			
Potassium	3.92 mEq/L	3.5 - 5.5			
Lipid Profile					
Cholesterol - Total	92 mg/dL	<= 199.9			
Triglycerides	▲ 159 mg/dL	<= 149.9			
Cholesterol - HDL	▼ 36 mg/dL	>= 39.9			
Cholesterol - LDL	24 mg/dl	<= 99.9			
Cholesterol- VLDL	▲ 32 mg/dl	<= 29.9			
Non HDL Cholesterol	56 mg/dL	<= 129.9			
Cardiac Profile					
Homocysteine	▲ 41.24 Åµmol/L	0 - 14.9			
High sensitivity CRP	▲ 3.44 mg/L	0 - 3			
Lipoprotein (a)	16.05 mg/dL	0 - 29.99			
Apolipoprotein - A1	99.00 mg/dL	79 - 169			
Apolipoprotein - B	57.00 mg/dL	46 - 174			
Apolipoprotein B/A1 Ratio	0.58 Ratio				
Liver Function Test					
Bilirubin - Total	0.74 mg/dL	0.3 - 1.2			



We don't have any of your previous lab results for these tests in our records

Doctor Summary For

Comprehensive Platinum Full Body Checkup with Smart Report

For

Arvind

Male /41 Yrs

Note This is an electronically generated summary of the attached report. It is advised to read this summary in conjunction with the attached report and to correlate it clinically. For the trends section, the out of range values are highlighted with respect to the bio reference range of respective reports.

Test Name	Result, 25/06/25	Bio. Ref. Interval	Trends (For last three tests)		
Liver Function Test			Date 1	Date 2	Date 3
Protein, Total	7.35 g/dL	5.7 - 8.2			
Albumin	4.71 g/dL	3.4 - 4.8			
Aspartate Transaminase (SGOT)	▲ 54 U/L	0 - 33.9			
Alanine Transaminase (SGPT)	▲ 64 U/L	10 - 49			
Alkaline Phosphatase	71 U/L	46 - 116			
Gamma Glutamyltransferase (GGT)	57 U/L	<= 72			
Pancreas Profile					
Amylase	86 U/L	30 - 118			
Lipase	37.0 U/L	12 - 53			
Urine Routine & Microscopy					
Appearance	Slightly Hazy	CLEAR			
Specific gravity	1.020	1.003 - 1.035			
pH	6.0	4.6 - 8			
Glucose	Negative	NEGATIVE			
Protein	Negative	NEGATIVE			
Ketones	Negative	NEGATIVE			
Leucocyte Esterase	Positive	NEGATIVE			
Nitrite	Positive	NEGATIVE			
Pus cells	7-8 /hpf	0 - 5			
Red blood cell	Nil /hpf	0 - 2			
Epithelial cells	2-3 /hpf	FEW			
Casts	Nil /lpf	NIL			
Crystals	Nil	NIL			
Bacteria	Present	NIL			



We don't have any of your previous lab results for these tests in our records

Doctor Summary For

Comprehensive Platinum Full Body Checkup with Smart Report

For

Arvind

Male /41 Yrs

Note This is an electronically generated summary of the attached report. It is advised to read this summary in conjunction with the attached report and to correlate it clinically. For the trends section, the out of range values are highlighted with respect to the bio reference range of respective reports.

Test Name	Result, 25/06/25	Bio. Ref. Interval	Trends (For last three tests)		
Calcium and Bone Health			Date 1	Date 2	Date 3
Calcium	▲ 11.1 mg/dL	8.7 - 10.4			
Vitamin D (25-OH)	30.5 ng/mL	30 - 100			
Phosphorus, Serum	2.50 mg/dl	2.4 - 5.1			
Vitamin Profile					
Vitamin B12	248.0 pg/ml	211 - 911			
Vitamin B9	16.42 ng/ml	>= 5.38			
Thyroid Function Test					
T3, Total	1.22 ng/mL	0.60 - 1.81			
T4, Total	12.1 Åµg/dl	4.5 - 12.6			
Thyroid Stimulating Hormone - Ultra Sensitive	▲ 4.818 uIU/ml	0.55 - 4.78			
Free T4	1.23 ng/dL	0.89 - 1.76			
Free T3	3.35 pg/mL	2.3 - 4.2			
Arthritis Screening					
Rheumatoid Factor - Quantitative	4.5 IU/mL	0 - 14			
Allergy Panel					
Immunoglobulin E (IgE) Total	▲ 202 IU/mL	0 - 158			



We don't have any of your previous lab results for these tests in our records

Wellbeing Index

Important Findings from your Wellbeing Index

For
Arvind
Male / 41 Yrs



Physicals

Height

Data not available

Weight

Data not available

Waist

Data not available

BMI

Data not available

Heart Age

Data not available

BP

Data not available



Disease Risks

Diabetes

Survey not taken yet

Hypertension

Survey not taken yet

Stroke

Survey not taken yet

CVD

Survey not taken yet

Depression

Survey not taken yet

Anxiety

Survey not taken yet

Stress

Survey not taken yet

* Embark on a better you by completing the wellbeing index. [Here](#)



Lifestyle Data

Habits

Data not available

Family History

Data not available

Important Parameters

From your Comprehensive Platinum Full Body Checkup with Smart Report

For
Arvind
Male / 41 Yrs



Complete Blood Count

Gives an insight into the health of blood and blood cells which are essential to carry out various bodily functions like transporting oxygen, fighting infections, and clotting blood after an injury.

Hemoglobin 16.2 g/dL Range: 13.0 – 17.0	RBC 5.26 $10^6/\text{cu.mm}$ Range: 4.5 – 5.5	HCT 48.9 % Range: 40 – 50	Total Leucocyte Count 8.38 $10^3/\mu\text{L}$ Range: 4 – 10
--	--	--	--

Neutrophils 55.1 % Range: 40 – 80	Lymphocytes 31.1 % Range: 20 – 40	Monocytes ▲ 10.1 % Range: 2 – 10	Eosinophils 3.1 % Range: 1 – 6	Basophils 0.6 % Range: 0 – 2
--	--	---	---	---

Platelet Count 319 $10^3/\mu\text{L}$ Range: 150 – 410	PDW ▲ 17.5 fL Range: 9 – 17
---	--



Inflammatory markers

Helps to understand presence of an inflammation in the body. Inflammation is bodies defence against infection or injury.

Erythrocyte Sedimentation Rate 4 mm/hr Range: 0 – 10	C-Reactive Protein (Quantitative) 5.10 mg/L Range: 0.1 – 10
---	--

Important Parameters

From your Comprehensive Platinum Full Body Checkup with Smart Report

For
Arvind
Male / 41 Yrs



Iron Studies

Iron is a vital mineral. It helps our blood cells to transport oxygen. Iron studies are used to assess level of iron in blood and blood's ability to attach itself to iron.

Iron Serum

84 µg/dL

Range: 65 - 175

Total Iron Binding Capacity (TIBC)

442.88 µg/dL

Range: 250 - 460

Unsaturated Iron Binding Capacity

▲ **359** µg/dL

Range: 111 - 343

Ferritin

220.00 ng/mL

Range: 22 - 322



Diabetes Profile

Measures the level of glucose in the body and helps identify the body's ability to process glucose. It can be used for screening as well as monitoring the treatment of diabetes.

Glycosylated Hemoglobin (HbA1c)

5.3 %

Range: 4 - 5.6

Glucose - Fasting

80 mg/dL

Range: 70 - 99

Microalbumin-Albumin

6.07 mg/L

Range: 0 - 29.99

Microalbumin-Albumin/Creatinine Ratio

9.58 mg/g Creatinine

Range: 0 - 29.99

Important Parameters

From your Comprehensive Platinum Full Body Checkup with Smart Report

For
Arvind
Male / 41 Yrs



Kidney Function Test

Performed to determine how well the kidneys are working. Kidneys regulate elimination of waste from our body and maintain electrolyte balance.

Blood Urea Nitrogen

▼ **8** mg/dL

Range: 9 - 23

Urea

▼ **16.24** mg/dL

Range: 19.26 - 49.22

Creatinine

0.88 mg/dL

Range: 0.7 - 1.3

Uric Acid

6.4 mg/dL

Range: 3.5 - 7.2

Sodium

142 mEq/L

Range: 132 - 146

Potassium

3.92 mEq/L

Range: 3.5 - 5.5



Lipid Profile

Measures the amount of Cholesterol and Triglycerides in your blood. This gives an insight into the health of heart and blood vessels.

Cholesterol - Total

92 mg/dL

Range: <= 199.9

Triglycerides

▲ **159** mg/dL

Range: <= 149.9

Cholesterol - HDL

▼ **36** mg/dL

Range: >= 39.9

Cholesterol - LDL

24 mg/dL

Range: <= 99.9

Cholesterol- VLDL

▲ **32** mg/dL

Range: <= 29.9

Non HDL Cholesterol

56 mg/dL

Range: <= 129.9

Important Parameters

From your Comprehensive Platinum Full Body Checkup with Smart Report

For
Arvind
Male / 41 Yrs



Cardiac Profile

A comprehensive blood test that offers detailed information about the risk of cardiovascular disease (CVD) and the overall health of the heart.

Homocysteine

▲ **41.24** $\mu\text{mol/L}$

Range: 0 - 14.9

High sensitivity CRP

▲ **3.44** mg/L

Range: 0 - 3

Lipoprotein (a)

16.05 mg/dL

Range: 0 - 29.99

Apolipoprotein - A1

99.00 mg/dL

Range: 79 - 169

Apolipoprotein - B

57.00 mg/dL

Range: 46 - 174

Apolipoprotein B/A1 Ratio

0.58 Ratio



Liver Function Test

Group of blood tests commonly performed to evaluate the function of the liver which is essential to digest food and removing toxins from the body.

Bilirubin - Total

0.74 mg/dL

Range: 0.3 - 1.2

Protein, Total

7.35 g/dL

Range: 5.7 - 8.2

Albumin

4.71 g/dL

Range: 3.4 - 4.8

Aspartate Transaminase (SGOT)

▲ **54** U/L

Range: 0 - 33.9

Alanine Transaminase (SGPT)

▲ **64** U/L

Range: 10 - 49

Alkaline Phosphatase

71 U/L

Range: 46 - 116

Gamma Glutamyltransferase (GGT)

57 U/L

Range: ≤ 72

Important Parameters

From your Comprehensive Platinum Full Body Checkup with Smart Report

For
Arvind
Male / 41 Yrs



Pancreas Profile

Measures the levels of digestive enzymes, lipase and amylase, produced by the pancreas which can be used to monitor the pancreatic health.

Amylase

86 U/L

Range: 30 - 118

Lipase

37.0 U/L

Range: 12 - 53



Urine Routine & Microscopy

Microscopic examination of urine sample to check for the presence of blood cells, crystals, bacteria, parasites, and cells from tumors in it.

Appearance

Slightly Hazy

Range: CLEAR

Specific gravity

1.020

Range: 1.003 - 1.035

pH

6.0

Range: 4.6 - 8

Glucose

Negative

Range: NEGATIVE

Protein

Negative

Range: NEGATIVE

Ketones

Negative

Range: NEGATIVE

Leucocyte Esterase

Positive

Range: NEGATIVE

Nitrite

Positive

Range: NEGATIVE

Pus cells

7-8 /hpf

Range: 0 - 5

Red blood cell

Nil /hpf

Range: 0 - 2

Epithelial cells

2-3 /hpf

Range: FEW

Casts

Nil /lpf

Range: NIL

Crystals

Nil

Range: NIL

Bacteria

Present

Range: NIL

Important Parameters

From your Comprehensive Platinum Full Body Checkup with Smart Report

For
Arvind
Male / 41 Yrs



Calcium and Bone Health

Measures the levels of calcium and vitamin D in the blood which are responsible for keeping bones, teeth, and muscles healthy.

Calcium

▲ **11.1** mg/dL

Range: 8.7 - 10.4

Vitamin D (25-OH)

30.5 ng/mL

Range: 30 - 100

Phosphorus, Serum

2.50 mg/dl

Range: 2.4 - 5.1



Vitamin Profile

Vitamins are the essential nutrients for human life. This profile offers tests to check level of different types of vitamin B, vitamin D, vitamin E and vitamin K.

Vitamin B12

248.0 pg/ml

Range: 211 - 911

Vitamin B9

16.42 ng/ml

Range: >= 5.38



Thyroid Function Test

Window to the health of the butterfly shaped gland - Thyroid, which determines how the body uses energy.

T3, Total

1.22 ng/mL

Range: 0.60 - 1.81

T4, Total

12.1 Åµg/dl

Range: 4.5 - 12.6

Thyroid Stimulating Hormone - Ultra Sensitive

▲ **4.818** uIU/ml

Range: 0.55 - 4.78

Free T4

1.23 ng/dL

Range: 0.89 - 1.76

Important Parameters

From your Comprehensive Platinum Full Body Checkup with Smart Report

For
Arvind
Male / 41 Yrs

Free T3

3.35 pg/mL

Range: 2.3 - 4.2



Arthritis Screening

Measures the amount of rheumatoid factor (RF) and Anti-CCP Antibody in the blood, which helps diagnose or monitor rheumatoid arthritis (RA) and differentiates it from other types of arthritis.

Rheumatoid Factor - Quantitative

4.5 IU/mL

Range: 0 - 14



Allergy Panel

This test aids in detecting various allergies, including seasonal, food, insect sting, and certain disorders of immune system.

Immunoglobulin E (IgE) Total

▲ **202** IU/mL

Range: 0 - 158

Recommendations

Care for better health and wellbeing

For
Arvind
Male / 41 Yrs



Lifestyle

Healthy eating



Do's

Cook At Home More Often

Cook more often to control ingredients and use healthier methods like steaming, grilling, or baking.

Avoid High-Calorie Meals And Stay Hydrated

Prioritize low-calorie meals and maintain adequate hydration for a healthier lifestyle.

Do's

Regular Bedtime And Rise Time

Maintain consistent bedtime and wake time to regulate sleep patterns and prevent sleep irregularities.

Don'ts

Limit Alcohol

Avoid alcohol before bed as its initial sedative effect wears off, disrupting sleep patterns.

Sleep hygiene



Exercise



Do's

Get Off One Stop Early

Get off the bus or train one stop early and walk the rest of the way.

Take Stairs

Take the stairs instead of the elevator.

References

From trusted sources

For

Arvind

Male / 41 Yrs

01

Estimation of 10-year Cardiovascular Disease (CVD) Risk
 D'Agostino RB Sr, et al. General cardiovascular risk profile for use in primary care: the Framingham Heart Study.Circulation. 2008 Feb 12;117(6):743-53

02

Framingham Heart Study: Hypertension Risk
 Parikh NI, et al. A risk score for predicting near-term incidence of hypertension: the Framingham Heart Study.Ann Intern Med. 2008;148(2):102-110.

03

Framningham Heart Study. Stroke Risk
 D'Agostino RB, et al. Stroke risk profile: adjustment for antihypertensive medication. The Framingham Study. Stroke. 1994;25(1):40-3.

04

Depression: Patient Health Questionnaire-2 (PHQ-2)
 Kroenke K, et al. The Patient Health Questionnaire-2: validity of a two-item depression screener.Med Care. 2003;41(11):1284-1292.

05

Anxiety: Generalized Anxiety Disorder 2-item (GAD-2)
 Kroenke K, et al. Anxiety disorders in primary care: prevalence, impairment, comorbidity, and detection.Ann Intern Med. 2007;146(5):317-325.

06

Anxiety: Generalized Anxiety Disorder 7-item (GAD-7)
 Spitzer RL, et al. A brief measure for assessing generalized anxiety disorder: the GAD-7.Arch Intern Med. 2006;166:1092-7.

07

Indian Diabetes Risk Score [IDRS]
 Mohan V, et al. A simplified Indian Diabetes Risk Score for screening for undiagnosed diabetic subjects. J Assoc Physicians India. 2005;53:759-763.

08

Dietary Guidelines for Indians
 Dietary Guidelines for Indians - A Manual, Second Edition, 2011.ICMR-National Institute of Nutrition, Hyderabad.

09

My plate for the day
 R. Hemalatha. Promotionof 'My Plate for the Day' and physical activity among the population to prevent all forms of malnutrition and NCDs in the country, 2023.ICMR-National Institute of Nutrition, Hyderabad.

10

Healthy Eating Plate
 Building a Healthy and Balanced DietThe Nutrition Source, Department of Nutrition, Harvard T.H. Chan School of Public Health.

11

Top 10 Take-Home Messages for the Primary Prevention of Cardiovascular Disease
 2019 ACC/AHA Guideline on the Primary Prevention of Cardiovascular Disease. Circulation. 2019 Sep 10;140(11).

12

Smoking cessation
 Age-friendly Primary Health Care Centres Toolkit. World Health Organization

13

Sleep Hygiene
 Irish LA, et al. The role of sleep hygiene in promoting public health: A review of empirical evidence. Sleep Med Rev. 2015;22:23-36.

14

Body mass index (BMI)
 Nutritional Status of Women and Men, 2019-21 India.National Family Health Survey (NFHS - 5), 2019-21.



PO No :PO3372032243-231



Customer Name	: Mr.ARVIND	Collected Via	: TATA IMG HYDERABAD
Age/Gender	: 41/Male	Referred By	: Dr.
Lab Visit ID	: HYD572178	Collection Date	: 25/Jun/2025 06:38AM
Barcode ID/Order ID	: D18948389 / 13257890	Report Date	: 25/Jun/2025 03:34PM
Sample Type	: EDTA	Report Status	: Final Report

HAEMATOLOGY

COMPREHENSIVE PLATINUM FULL BODY CHECKUP WITH SMART REPORT

Test Name	Result	Unit	Bio. Ref. Interval	Method
Complete Blood Count				
Hemoglobin	16.2	g/dL	13.0-17.0	Spectrophotometry (Cyanide-free)
RBC	5.26	10 ⁶ /cu.mm	4.5 - 5.5	Impedance
HCT	48.9	%	40 - 50	Calculated
MCV	93.0	fL	83 - 101	Calculated
MCH	30.8	pg	27 - 32	Calculated
MCHC	33.1	g/dL	31.5 - 34.5	Calculated
RDW-CV	14.0	%	11.5-14	Calculated
Total Leucocyte Count	8.38	10 ³ /μL	4 - 10	Impedance
Differential Leucocyte Count				
Neutrophils	55.1	%	40-80	DHSS/Microscopy
Lymphocytes	31.1	%	20-40	DHSS/Microscopy
Monocytes	10.1	%	2-10	DHSS/Microscopy
Eosinophils	3.1	%	1-6	DHSS/Microscopy
Basophils	0.6	%	0-2	Impedance/Microscopy
Absolute Leucocyte Count				
Absolute Neutrophil Count	4.62	10 ³ /μL	2 - 7	Calculated
Absolute Lymphocyte Count	2.61	10 ³ /μL	1-3	Calculated
Absolute Monocyte Count	0.85	10 ³ /μL	0.2 - 1	Calculated
Absolute Eosinophil Count	0.26	10 ³ /μL	0.02 - 0.5	Calculated
Absolute Basophil Count	0.05	10 ³ /μL	0.02-0.1	Calculated
Platelet Count	319	10 ³ /μL	150-410	Impedance /Microscopy
MPV	9.4	fL	6.5 - 12	Calculated
PDW	17.5	fL	9 - 17	Calculated

Comment:

As per the recommendation of International council for Standardization in Hematology, the differential leucocyte counts are additionally being reported as absolute numbers of each cell in per unit volume of blood.

DHSS : Double Hydrodynamic Sequential System Flowcytometry

Calculated parameters are either derived from Impedance measure, RBC pulse measurement, RBC/platelet histograms or formula derived.

NABL certificate
and scope

This test has been performed at

TATA 1MG HYDERABAD

Address: SCB Door No. 3-14-011, 1st Floor,
Patny Square, SP Road, Rasoolpura,
Secunderabad, Telangana - 500003

Dr. Vittal Sri Navya
MBBS, MD (Pathology)
Consultant Pathologist
Reg. No: 85499

Scan for
digital copy

Page 1 of 28



PO No :PO3372032243-231



Customer Name	: Mr.ARVIND	Collected Via	: TATA IMG HYDERABAD
Age/Gender	: 41/Male	Referred By	: Dr.
Lab Visit ID	: HYD572178	Collection Date	: 25/Jun/2025 06:38AM
Barcode ID/Order ID	: D18948389 / 13257890	Report Date	: 25/Jun/2025 03:34PM
Sample Type	: EDTA	Report Status	: Final Report

HAEMATOLOGY

COMPREHENSIVE PLATINUM FULL BODY CHECKUP WITH SMART REPORT

Test Name	Result	Unit	Bio. Ref. Interval	Method
Erythrocyte Sedimentation Rate				
Erythrocyte Sedimentation Rate	4	mm/hr	0-10	Modified Westergren

Comment:

- ESR provides an index of progress of the disease and is widely used as an indicator of inflammation, infection, trauma, or malignant diseases. Changes are more significant than a single abnormal test
- It is specifically indicated to monitor the course or response to the treatment of diseases like rheumatoid arthritis, tuberculosis bacterial endocarditis, acute rheumatic fever, Hodgkins disease, temporal arthritis, and systemic lupus erythematosus; and to diagnose and monitor giant cell arteritis and polymyalgia rheumatica.
- An elevated ESR may also be associated with many other conditions, including autoimmune disease, anemia, infection, malignancy, pregnancy, multiple myeloma, menstruation, and hypothyroidism.
- Although a normal ESR cannot be taken to exclude the presence of organic disease, its rate is dependent on various physiologic and pathologic factors.
- The most important component influencing ESR is the composition of plasma. High level of C-Reactive Protein, fibrinogen, haptoglobin, alpha-1 antitrypsin, ceruloplasmin and immunoglobulins causes the elevation of Erythrocyte Sedimentation Rate.
- Drugs that may cause increase ESR levels include: dextran, methyl dopa, oral contraceptives, penicillamine, procainamide, theophylline, and Vitamin A. Drugs that may cause decrease levels include: aspirin, cortisone, and quinine

NABL certificate
and scope

This test has been performed at
TATA 1MG HYDERABAD
Address: SCB Door No. 3-14-011, 1st Floor,
Patny Square, SP Road, Rasoolpura,
Secunderabad, Telangana - 500003

Dr. Vittal Sri Navya
MBBS, MD (Pathology)
Consultant Pathologist
Reg. No: 85499

Scan for
digital copy

Page 2 of 28



PO No :PO3372032243-231



Customer Name	: Mr.ARVIND	Collected Via	: TATA IMG HYDERABAD
Age/Gender	: 41/Male	Referred By	: Dr.
Lab Visit ID	: HYD572178	Collection Date	: 25/Jun/2025 06:38AM
Barcode ID/Order ID	: D18948389 / 13257890	Report Date	: 25/Jun/2025 03:19PM
Sample Type	: WHOLE BLOOD-EDTA	Report Status	: Final Report

HAEMATOLOGY

COMPREHENSIVE PLATINUM FULL BODY CHECKUP WITH SMART REPORT

Test Name	Result	Unit	Bio. Ref. Interval	Method
HbA1c (Glycosylated Hemoglobin)				
Glycosylated Hemoglobin (HbA1c)	5.3	%	4-5.6	HPLC (NGSP certified)
Estimated average glucose (eAG)	105.41	mg/dL		Calculated

Comment:

Interpretation: HbA1c%

≤5.6	Normal
5.7-6.4	At Risk For Diabetes
≥6.5	Diabetes

Adapted from American Diabetes Association.

Comments:

A 3 to 6 monthly monitoring is recommended in diabetics. People with diabetes should get the test done more often if their blood sugar stays too high or if their healthcare provider makes any change in the treatment plan. HbA1c concentration represent the integrated values for blood glucose over the preceding 8-12 weeks and is not affected by daily glucose fluctuation, exercise & recent food intake.

Please note, Glycemic goal should be individualized based on duration of diabetes, age/life expectancy, comorbid conditions, known CVD or advanced microvascular complications, hypoglycemia unawareness, and individual patient considerations.

Factors that interfere with HbA1c Measurement: Hemoglobin variants, elevated fetal hemoglobin (HbF) and chemically modified derivatives of hemoglobin (e.g. carbamylated Hb in patients with renal failure) can affect the accuracy of HbA1c measurements.

Factors that affect interpretation of HbA1c Measurement: Any condition that shortens erythrocyte survival or decrease mean erythrocyte age (e. g., recovery from acute blood loss, hemolytic anemia, HbSS, HbCC, and HbSC) will falsely lower HbA1c test results regardless of the assay method used. Iron deficiency anemia is associated with higher HbA1c.

Note: Presence of Hemoglobin variants and/or conditions that affect red cell turnover must be considered, particularly when the HbA1c result does not correlate with the patient's blood glucose levels.

- HPLC - High performance liquid chromatography

NABL certificate
and scope

This test has been performed at
TATA 1MG HYDERABAD
Address: SCB Door No. 3-14-011, 1st Floor,
Patny Square, SP Road, Rasoolpura,
Secunderabad, Telangana - 500003

Dr. K Madhuri
MBBS, MD (Pathology)
Consultant Pathologist
Reg. No: 80149

Scan for
digital copy

Page 3 of 28



PO No :PO3372032243-231



Customer Name	: Mr.ARVIND	Collected Via	: TATA IMG HYDERABAD
Age/Gender	: 41/Male	Referred By	: Dr.
Lab Visit ID	: HYD572178	Collection Date	: 25/Jun/2025 06:38AM
Barcode ID/Order ID	: D18948391 / 13257890	Report Date	: 25/Jun/2025 04:22PM
Sample Type	: Fluoride Plasma F	Report Status	: Final Report

BIOCHEMISTRY

COMPREHENSIVE PLATINUM FULL BODY CHECKUP WITH SMART REPORT

Test Name	Result	Unit	Bio. Ref. Interval	Method
FBS (Fasting Blood Sugar)				
Glucose - Fasting	80	mg/dL	70.0 - 99.0	Hexokinase

Fasting Plasma Glucose (mg/dL)	2 hr plasma Glucose (mg/dL)	Diagnosis
99 or below	139 or below	Normal
100 to 125	140 to 199	Pre-Diabetes (IGT)
126 or above	200 or above	Diabetes

Reference : American Diabetes Association

Comment:

Impaired glucose tolerance (IGT) fasting, means a person has an increased risk of developing type 2 diabetes but does not have it yet. A level of 126 mg/dL or above, confirmed by repeating the test on another day, means a person has diabetes. IGT (2 hrs Post meal), means a person has an increased risk of developing type 2 diabetes but does not have it yet. A 2-hour glucose level of 200 mg/dL or above, confirmed by repeating the test on another day, means a person has diabetes

Plasma Glucose Goals	For people with Diabetes
Before meal	70-130 mg/dL
2 Hours after meal	Less than 180 mg/dL
HbA1c	Less than 7%

NABL certificate
and scope

This test has been performed at
TATA 1MG HYDERABAD
Address: SCB Door No. 3-14-011, 1st Floor,
Patny Square, SP Road, Rasoolpura,
Secunderabad, Telangana - 500003

Dr. K Madhuri
MBBS, MD (Pathology)
Consultant Pathologist
Reg. No: 80149

Scan for
digital copy

Page 4 of 28



PO No :PO3372032243-231



Customer Name	: Mr.ARVIND	Collected Via	: TATA IMG HYDERABAD
Age/Gender	: 41/Male	Referred By	: Dr.
Lab Visit ID	: HYD572178	Collection Date	: 25/Jun/2025 06:38AM
Barcode ID/Order ID	: D18948383 / 13257890	Report Date	: 25/Jun/2025 09:28PM
Sample Type	: Serum	Report Status	: Final Report

BIOCHEMISTRY

COMPREHENSIVE PLATINUM FULL BODY CHECKUP WITH SMART REPORT

Test Name	Result	Unit	Bio. Ref. Interval	Method
Lipid Profile				
Cholesterol - Total	92	mg/dL	Low (desirable): < 200 mg/dL Moderate (borderline) 200–239 mg/dL High: >= 240 mg/dL	Enzymatic
Triglycerides	159	mg/dL	Normal: <150, Borderline: 150 - 199, High:200-499, Very High>=500	GPO
Cholesterol - HDL	36	mg/dL	Undesirable/high risk <40mg/dL Desirable/low risk>=60mg/dl	Accelerator Selective Detergent
Cholesterol - LDL	24	mg/dl	Desirable: <100 Above desirable: 100 - 129 Borderline high : 130 - 159 High : 160 - 189 Very high : >=190	Calculated
Cholesterol- VLDL	32	mg/dl	<30	Calculated
Cholesterol : HDL Cholesterol	2.6	Ratio	Desirable : 3.5-4.5 High Risk : >5	Calculated
LDL : HDL Cholesterol	0.67	Ratio	Desirable : 2.5-3.0 High risk : >3.5	Calculated
Non HDL Cholesterol	56	mg/dL	Desirable:< 130, Above Desirable:130 - 159, Borderline High:160 - 189, High:190 - 219,	Calculated

NABL certificate and scope



This test has been performed at
TATA 1MG HYDERABAD
 Address: SCB Door No. 3-14-011, 1st Floor,
 Patny Square, SP Road, Rasoolpura,
 Secunderabad, Telangana - 500003

Dr. Vittal Sri Navya

Dr. Vittal Sri Navya
 MBBS, MD (Pathology)
 Consultant Pathologist
 Reg. No: 85499

Scan for digital copy





PO No :PO3372032243-231



Customer Name	: Mr.AR.VIND	Collected Via	: TATA IMG HYDERABAD
Age/Gender	: 41/Male	Referred By	: Dr.
Lab Visit ID	: HYD572178	Collection Date	: 25/Jun/2025 06:38AM
Barcode ID/Order ID	: D18948383 / 13257890	Report Date	: 25/Jun/2025 09:28PM
Sample Type	: Serum	Report Status	: Final Report

BIOCHEMISTRY

COMPREHENSIVE PLATINUM FULL BODY CHECKUP WITH SMART REPORT

Test Name	Result	Unit	Bio. Ref. Interval	Method
Very High: >= 220				

Comment:

- Lipid profile measurements in the same patient can show physiological & analytical variations. It is recommended that 3 serial samples 1 week apart may be tested.
- Indians are at a high risk of developing atherosclerotic cardiovascular disease (ASCVD); at a much earlier age and more severe with high mortality. Dyslipidemia (abnormal lipid profile) is the major risk factor and found in almost 80% Indians.
- Total cholesterol** is the total amount of cholesterol in blood comprising of HDL, LDL-C, and VLDL.
- LDL Cholesterol (LDL-C)** or "bad" cholesterol contributes most significantly to atherosclerosis leading to heart disease or stroke and is the primary target for reducing risk for cardiovascular disease.
- High-density lipoprotein (HDL)** or "good" cholesterol can lower risk of heart disease and stroke.
- Triglyceride (TG)** level also plays a major role in CVD. Indians are more prone to Atherogenic dyslipidemia, a condition associated with high TG, low HDL-C and high LDL-C; this is associated with diabetes, metabolic syndrome and insulin resistance. Hence high triglyceride levels also need to be treated.
- Non-HDL-Cholesterol (Non-HDL-C)** measures all plaque forming lipoproteins (e.g. remnants, LDL-C, VLDL, Lp(a), Apo-B). Monitoring of Non-HDL-C is important in patients with high TG (e.g. diabetics, obese persons) and those already on statin therapy.
- Lipid Association of India (LAI-2020) recommends:-**

- Screening of all Indians above the age of 20 years for CVD risk factors, esp. lipid profile.
- Identification of Risk factors: Age (male ≥45 years, female ≥55 years); Family h/o heart disease at younger age (<55 yrs in males, <65 yrs in female), Smoking/tobacco use, High blood pressure, Low HDL (males <40 mg/dl and females <50mg/dl).
- Fasting lipid profile is not mandatory for screening. Both fasting and non-fasting lipid profiles are equally important for managing Indian patients.
- Non-HDL-C should be calculated in every subject. LAI recommends LDL-C as the primary target and Non-HDL-C as the co-primary target for initiating drug therapy.
- Lifestyle modifications are of first and foremost importance for management and prevention of dyslipidemia. Among low risk groups, treatment is started only after 3 months of lifestyle changes.
- Testing for Apolipoprotein B, hsCRP, Lp(a) should be considered for patients in moderate risk group.
- Newer treatment goals based on Risk Groups and values of LDL-C and Non-HDL-C

New treatment goals by Lipid Association of India (2020)

	CONSIDER THERAPY (cut-off level)		TREATMENT GOALS	
Risk groups	LDL-C (mg/dL)	Non-HDL-C (mg/dL)	LDL-C (mg/dL)	Non-HDL-C (mg/dL)
Extreme Risk Gp Cat. A	≥50	≥80	<50 (Optional ≤30)	<80 (Optional ≤60)
Extreme Risk Gp Cat. B	>30	>60	≤30	≤60
Very High Risk	≥50	≥80	<50	<80
High Risk	≥70	≥100	<70	<100
Moderate Risk	≥100	≥130	<100	<130
Low risk	≥130*	≥160*	<100	<130

NABL certificate and scope



This test has been performed at
TATA 1MG HYDERABAD
 Address: SCB Door No. 3-14-011, 1st Floor,
 Patny Square, SP Road, Rasoolpura,
 Secunderabad, Telangana - 500003

Dr. Vittal Sri Navya

Dr. Vittal Sri Navya
 MBBS, MD (Pathology)
 Consultant Pathologist
 Reg. No: 85499

Scan for digital copy





PO No :PO3372032243-231



Customer Name	: Mr.ARVIND	Collected Via	: TATA IMG HYDERABAD
Age/Gender	: 41/Male	Referred By	: Dr.
Lab Visit ID	: HYD572178	Collection Date	: 25/Jun/2025 06:38AM
Barcode ID/Order ID	: D18948383 / 13257890	Report Date	: 25/Jun/2025 09:28PM
Sample Type	: Serum	Report Status	: Final Report

BIOCHEMISTRY

COMPREHENSIVE PLATINUM FULL BODY CHECKUP WITH SMART REPORT

Test Name	Result	Unit	Bio. Ref. Interval	Method
*After an adequate non-pharmacological intervention for at least 3 months				

● **As per NCEP Expert Panel (2011) guidelines**, universal screening for dyslipidemia is recommended for children between 9 - 11 yrs (repeat at 17-21 yrs). Screening is not recommended before the age of 2yrs. Above the age of 2 yrs, selective screening is done in children with family history of premature CVD or risk factors like obesity, diabetes, and hypertension.

Note: Reference Interval as per National Cholesterol Education Program (NCEP) Report.

LIVER FUNCTION TEST

Liver Function Test

Bilirubin-Total	0.74	mg/dL	0.3 – 1.2	Vanadate oxidation
Bilirubin-Direct	0.16	mg/dL	0.0-0.3	Vanadate oxidation
Bilirubin-Indirect	0.58	mg/dL	0.2-0.8	Calculated
Protein, Total	7.35	g/dL	5.7 - 8.2	Biuret
Albumin	4.71	g/dL	3.4 - 4.8	BCG Dye Binding
Globulin	2.6	g/dL	2.1 - 3.9	Calculated
A/G Ratio	1.78	Ratio	0.8 - 2.1	Calculated
Aspartate Transaminase (SGOT)	54	U/L	<34	Modified IFCC
SGPT (Alanine Transaminase)	64	U/L	10-49	Modified IFCC
SGOT/SGPT	0.84	Ratio		Calculated
Alkaline Phosphatase	71	U/L	46-116	IFCC Standardization
Gamma Glutamyltransferase (GGT)	57	U/L	<73	Modified IFCC

Comment:

- Raised ALT and AST indicate hepatocellular damage (e.g. viral or drugs etc). ALT is more liver-specific while AST is also found in heart, skeletal muscle, and kidney. Mild elevation (less than twice normal) often resolves on its own. Fatty liver disease (especially with metabolic syndrome) is a common cause in asymptomatic cases. Certain drugs (paracetamol, statins), herbal supplements, energy drinks, and antibiotics may also affect liver function.
- SGOT/SGPT Ratio: Typically <1 in healthy individuals (vary between 0.7-1.4; higher in women than men). High SGPT (ratio <1) seen in acute or chronic hepatitis, autoimmune disorders, medications, toxins while ratio >1 indicates alcoholic hepatitis, cirrhosis, metastasis or non-hepatic issues (hemolytic diseases, CVS disorders).
- Elevated Alkaline Phosphatase and GGT: Suggest cholestatic diseases (e.g. bile duct obstruction, primary biliary

NABL certificate and scope



This test has been performed at
TATA 1MG HYDERABAD
Address: SCB Door No. 3-14-011, 1st Floor,
Patny Square, SP Road, Rasoolpura,
Secunderabad, Telangana - 500003

Dr. Vittal Sri Navya
MBBS, MD (Pathology)
Consultant Pathologist
Reg. No: 85499

Scan for digital copy



Page 7 of 28



PO No :PO3372032243-231



Customer Name	: Mr.ARVIND	Collected Via	: TATA IMG HYDERABAD
Age/Gender	: 41/Male	Referred By	: Dr.
Lab Visit ID	: HYD572178	Collection Date	: 25/Jun/2025 06:38AM
Barcode ID/Order ID	: D18948383 / 13257890	Report Date	: 25/Jun/2025 09:28PM
Sample Type	: Serum	Report Status	: Final Report

BIOCHEMISTRY

COMPREHENSIVE PLATINUM FULL BODY CHECKUP WITH SMART REPORT

Test Name	Result	Unit	Bio. Ref. Interval	Method
cirrhosis etc.) and can also be due to bone disease, pregnancy, chronic renal failure, malignancy, and congestive heart failure.				
<ul style="list-style-type: none">High Bilirubin: Indicates jaundice due to increased RBC breakdown, liver damage (e.g., infections, toxins), or cholestasis (e.g., gallstones, tumors).High Protein Levels: Seen in dehydration (e.g., severe vomiting, diarrhea) or increased production (e.g., inflammation, hematopoietic neoplasms). Low protein and albumin: Result from impaired synthesis (liver disease), decreased intake, tissue damage, malabsorption, or increased renal excretion.				

TATA 1mg Labs

NABL certificate
and scope

This test has been performed at
TATA 1MG HYDERABAD
Address: SCB Door No. 3-14-011, 1st Floor,
Patny Square, SP Road, Rasoolpura,
Secunderabad, Telangana - 500003

Dr. Vittal Sri Navya
MBBS, MD (Pathology)
Consultant Pathologist
Reg. No: 85499

Scan for
digital copy

Page 8 of 28

PO No :PO3372032243-231



Customer Name	: Mr.ARVIND	Collected Via	: TATA IMG HYDERABAD
Age/Gender	: 41/Male	Referred By	: Dr.
Lab Visit ID	: HYD572178	Collection Date	: 25/Jun/2025 06:38AM
Barcode ID/Order ID	: D18948383 / 13257890	Report Date	: 25/Jun/2025 07:55PM
Sample Type	: Serum	Report Status	: Final Report

BIOCHEMISTRY

COMPREHENSIVE PLATINUM FULL BODY CHECKUP WITH SMART REPORT

Test Name	Result	Unit	Bio. Ref. Interval	Method
Kidney Function Test with eGFR (18 Years & Above)				
Blood Urea Nitrogen	8	mg/dL	9.0 - 23.0	Urease with GLDH
Urea	16.24	mg/dL	19.26 - 49.22	Calculated
Creatinine	0.88	mg/dL	0.7-1.3	Alkaline picrate-kinetic
Uric Acid	6.4	mg/dL	3.5-7.2	Uricase/Peroxidase
Sodium	142	mEq/L	132.0-146.0	Indirect ISE
Potassium	3.92	mEq/L	3.5 - 5.5	Indirect ISE
Chloride	105.3	mmol/L	98-107	Indirect ISE
BUN/Creatinine Ratio	8.6	Ratio	12:1 - 20:1	Calculated
Glomerular Filtration Rate (estimated)	111	mL/min/1.73m ²		Calculated

- Calculation of eGFR is based on 2021 CKD-EPI creatinine equation, recommended method for estimating GFR in adults (≥ 18 yrs). Estimates GFR using serum creatinine, age and sex.
- GFR categories defined acc. to KDIGO 2024 Clinical Practice Guideline for the Evaluation and Management of CKD.
- eGFR_{creat} levels less than 60 mL/min/1.73 m² should be considered as "decreased GFR".
- It is advised to estimate GFR using Cystatin C for confirmation of CKD in patients with eGFR_{creat} value 45-59 mL/min/1.73 sq.m without evidence of kidney damage and in specific circumstances when eGFR_{creat} is less accurate e.g. race, muscle mass, exercise, pregnancy, dehydration and nutritional status.

Interpretation:

GFR categories in CKD

Category	GFR (mL/min/1.73 sq.m)	Terms
G1	≥ 90	Normal or high
G2	60-89	Mildly decreased
G3a	45-59	Mildly to moderately decreased
G3b	30-44	Moderately to severely decreased
G4	15-29	Severely decreased
G5	< 15	Kidney failure

Criteria for CKD (Either of the following present for > 3 months)

Albuminuria (ACR > 30 mg/g)
Urine sediment abnormalities

This test has been performed at

TATA 1MG HYDERABADAddress: SCB Door No. 3-14-011, 1st Floor,
Patny Square, SP Road, Rasoolpura,
Secunderabad, Telangana - 500003Dr. Vittal Sri Navya
MBBS, MD (Pathology)
Consultant Pathologist
Reg. No: 85499NABL certificate
and scopeScan for
digital copy

Page 9 of 28

State of the Art Lab Facilities

Delhi • Gurgaon • Jaipur • Lucknow • Patna • Dehradun • Ahmedabad • Mumbai • Pune • Bangalore
Hyderabad • Chennai • Kolkata • Bhubaneswar • Angul

PO No :PO3372032243-231



Customer Name	: Mr.ARVIND	Collected Via	: TATA IMG HYDERABAD
Age/Gender	: 41/Male	Referred By	: Dr.
Lab Visit ID	: HYD572178	Collection Date	: 25/Jun/2025 06:38AM
Barcode ID/Order ID	: D18948383 / 13257890	Report Date	: 25/Jun/2025 07:55PM
Sample Type	: Serum	Report Status	: Final Report

BIOCHEMISTRY

COMPREHENSIVE PLATINUM FULL BODY CHECKUP WITH SMART REPORT

Test Name	Result	Unit	Bio. Ref. Interval	Method
Markers of kidney damage (one or more)	Electrolyte and other abnormalities due to tubular disorders Abnormalities detected by histology Structural abnormalities detected by imaging History of kidney transplantation			
Decreased GFR	GFR <60 ml/min/1.73 sq.m (GFR categories G3a-G5)			

Source: KDIGO 2024 Clinical Practice Guideline.



NABL certificate and scope



This test has been performed at
TATA 1MG HYDERABAD
Address: SCB Door No. 3-14-011, 1st Floor,
Patny Square, SP Road, Rasoolpura,
Secunderabad, Telangana - 500003

Dr. Vittal Sri Navya
MBBS, MD (Pathology)
Consultant Pathologist
Reg. No: 85499

Scan for digital copy





PO No :PO3372032243-231



Customer Name	: Mr.ARVIND	Collected Via	: TATA IMG HYDERABAD
Age/Gender	: 41/Male	Referred By	: Dr.
Lab Visit ID	: HYD572178	Collection Date	: 25/Jun/2025 06:38AM
Barcode ID/Order ID	: D18948383 / 13257890	Report Date	: 25/Jun/2025 10:03PM
Sample Type	: Serum	Report Status	: Final Report

BIOCHEMISTRY

COMPREHENSIVE PLATINUM FULL BODY CHECKUP WITH SMART REPORT

Test Name	Result	Unit	Bio. Ref. Interval	Method
Calcium	11.1	mg/dL	8.7-10.4	Arsenazo III

Comment:

Increased in: Hyperparathyroidism primary and secondary, Acute and chronic renal failure, Following renal transplantation, Osteomalacia with malabsorption, Acute osteoporosis, Malignant tumours (specially of breast, lung and kidney), Drugs: Vit. D and A intoxication, Diuretics, estrogen, androgen, tamoxifen, lithium

Decreased in: Hypoparathyroidism, Surgical and Idiopathic, Pseudohypoparathyroidism, Chronic renal disease with uremia and phosphate retention, Malabsorption of Calcium and Vit.D, obstructive jaundice, Bone Disease (Osteomalacia and rickets), Drugs: Cancer chemotherapy drugs, calcitonin, loop-actives diuretics, Hypomagnesemia, Hypoalbuminemia

TATA 1mg Labs

NABL certificate
and scope

This test has been performed at
TATA 1MG HYDERABAD
Address: SCB Door No. 3-14-011, 1st Floor,
Patny Square, SP Road, Rasoolpura,
Secunderabad, Telangana - 500003

Dr. Vittal Sri Navya
MBBS, MD (Pathology)
Consultant Pathologist
Reg. No: 85499

Scan for
digital copy

Page 11 of 28

PO No :PO3372032243-231



Customer Name	: Mr.ARVIND	Collected Via	: TATA IMG HYDERABAD
Age/Gender	: 41/Male	Referred By	: Dr.
Lab Visit ID	: HYD572178	Collection Date	: 25/Jun/2025 06:38AM
Barcode ID/Order ID	: D18948383 / 13257890	Report Date	: 25/Jun/2025 09:28PM
Sample Type	: Serum	Report Status	: Final Report

BIOCHEMISTRY

COMPREHENSIVE PLATINUM FULL BODY CHECKUP WITH SMART REPORT

Test Name	Result	Unit	Bio. Ref. Interval	Method
Phosphorus, Serum	2.50	mg/dl	2.4 - 5.1	Phosphomolybdate

Comment:

Phosphate metabolism is under the regulation of PTH, Vitamin D metabolites, and Fibroblast growth factor-23. Serum phosphate concentrations are about 50% higher in infants than in adults and decline throughout childhood as a consequence of the ability of growth hormone to increase the renal phosphate threshold.

Increased in:

Decreased Renal filtration (Acute or chronic renal failure) or increased reabsorption (e.g., hypoparathyroidism)
Increased Phosphate load (e.g., Oral or iv administration, Phosphate-containing laxatives or enemas, Vitamin D intoxication)
Cell Lysis (e.g., hemolysis, leukemias, chemotherapy, rhabdomyolysis)
Bone disease (e.g., healing fractures, multiple myeloma, Paget disease, osteolytic tumors)
Genetic (e.g., Hypoparathyroidism, Tumoral calcinosis)

Decreased in:

Intracellular Shift (e.g., Oral or intravenous Glucose, Insulin, Diabetic ketoacidosis, Respiratory alkalosis, Alcoholism, Severe burns)
Lowered Renal Phosphate Threshold (e.g., Primary or secondary hyperparathyroidism, Renal tubular defects)
Decreased Intestinal Absorption (Malabsorption syndrome, Vitamin D deficiency) or increased loss (Vomiting, Diarrhea)
Drugs (e.g., Salicylate, Paracetamol, Estrogens, Diuretics, Bisphosphonates, Anticonvulsants, Phosphate binding antacids, Antiviral drugs etc)

Note:

Because a significant diurnal variation in plasma phosphate has been reported, fasting morning specimens are recommended. Levels are influenced by dietary intake, meals, and exercise.

NABL certificate
and scope



This test has been performed at
TATA 1MG HYDERABAD
Address: SCB Door No. 3-14-011, 1st Floor,
Patny Square, SP Road, Rasoolpura,
Secunderabad, Telangana - 500003

Dr. Vittal Sri Navya
MBBS, MD (Pathology)
Consultant Pathologist
Reg. No: 85499

Scan for
digital copy





PO No :PO3372032243-231



Customer Name	: Mr.ARVIND	Collected Via	: TATA IMG HYDERABAD
Age/Gender	: 41/Male	Referred By	: Dr.
Lab Visit ID	: HYD572178	Collection Date	: 25/Jun/2025 06:38AM
Barcode ID/Order ID	: D18948383 / 13257890	Report Date	: 26/Jun/2025 12:52AM
Sample Type	: Serum	Report Status	: Final Report

BIOCHEMISTRY

COMPREHENSIVE PLATINUM FULL BODY CHECKUP WITH SMART REPORT

Test Name	Result	Unit	Bio. Ref. Interval	Method
Iron Studies, Comprehensive				
Iron Serum	84	µg/dL	65-175	Ferrozine
Total Iron Binding Capacity (TIBC)	442.88	µg/dL	250-460	Calculated
Unsaturated Iron Binding Capacity	359	µg/dL	111-343	Ferene
Transferrin saturation	18.96	%	16-50	Calculated
Ferritin	220.00	ng/mL	22 - 322	CLIA

Comment:

Iron is an essential trace mineral element which forms an important component of hemoglobin, metallocompounds and Vitamin A. Deficiency of iron is seen in iron deficiency and anaemia of chronic disorders. Increased iron concentration are seen in hemolytic anaemias, hemochromatosis and acute liver disease. Serum Iron alone is unreliable due to considerable physiologic diurnal variation in the results with highest values in the morning and lowest values in the evening as well as variation in response to iron therapy.

Total Iron Binding capacity (TIBC) is a direct measure of the protein Transferrin which transports iron from the gut to storage sites in the bone marrow. Increased levels of TIBC suggest that total iron body stores are low, increased concentration may be the sign of Iron deficiency anaemia, polycythemia vera, and may occur during the third trimester of pregnancy. Decreased levels may be seen in hemolytic anaemia, hemochromatosis, chronic liver disease, hypoproteinemia, malnutrition.

Unsaturated Iron Binding Capacity (UIBC) is increased in low iron state and decreased in high iron concentration such as hemochromatosis. In case of anaemia of chronic disease the patient may be anaemic but has adequate iron reserve and a low UIBC.

Transferrin Saturation occurs in Idiopathic hemochromatosis and Transfusional hemosiderosis where no unsaturated iron binding capacity is available for iron mobilization. Similar condition is seen in congenital deficiency of Transferrin.

*Please note change in BRI of Ferritin.

NABL certificate
and scope

This test has been performed at
TATA 1MG HYDERABAD
Address: SCB Door No. 3-14-011, 1st Floor,
Patny Square, SP Road, Rasoolpura,
Secunderabad, Telangana - 500003

Dr. Vittal Sri Navya
MBBS, MD (Pathology)
Consultant Pathologist
Reg. No: 85499

Scan for
digital copy



PO No :PO3372032243-231



Customer Name	: Mr.ARVIND	Collected Via	: TATA IMG HYDERABAD
Age/Gender	: 41/Male	Referred By	: Dr.
Lab Visit ID	: HYD572178	Collection Date	: 25/Jun/2025 06:38AM
Barcode ID/Order ID	: D18948383 / 13257890	Report Date	: 25/Jun/2025 07:55PM
Sample Type	: Serum	Report Status	: Final Report

BIOCHEMISTRY

COMPREHENSIVE PLATINUM FULL BODY CHECKUP WITH SMART REPORT

Test Name	Result	Unit	Bio. Ref. Interval	Method
Lipase				
Lipase	37.0	U/L	12-53	Colorimetric rate

Comment:

Pancreas is the major and primary source of serum lipase, though lipase is also secreted by the gastric and intestinal mucosa. Lipase measurement in serum is used to diagnose acute pancreatitis. After an attack of acute pancreatitis, serum Lipase activity increases within 4 to 8 hours, peaks at about 24 hours, and decreases over 8 to 14 days. Concentrations often remain elevated longer than those of Amylase. The increase in serum Lipase activity is not necessarily proportional to the severity of the attack.

Increased levels are seen in:

- Acute & Chronic Pancreatitis.
- Obstruction of Pancreatic duct.
- Non pancreatic conditions like renal disease, intestinal obstruction, acute cholecystitis, duodenal ulcer, alcoholism, diabetic ketoacidosis and following endoscopic retrograde cholangiopancreatography (ERCP).

Amylase

Amylase	86	U/L	30.0 - 118.0	Ethylidene Blocked-pNPG7
---------	----	-----	--------------	--------------------------

Comment:

- Amylase is a digestive enzyme mainly secreted by pancreas and salivary glands.
- An elevation of serum amylase beyond three times the upper limit of normal, combined with either clinical symptoms and/or imaging findings, may indicate acute pancreatitis. Serum amylase levels typically rise within 6 to 48 hours and usually return to baseline within 3 to 7 days. However, because of its short half-life, amylase levels may normalize as quickly as 24 hours after onset. Additionally, around 20% of patients with acute pancreatitis may have normal or near-normal amylase levels. Therefore, lipase, which is more specific to pancreatitis, should be measured alongside amylase to improve diagnostic accuracy.
- Elevated amylase levels can also be associated with conditions such as pancreatic duct obstruction, pancreatic carcinoma, or pancreatic pseudocysts. Additionally, increased amylase levels may occur in cholecystitis, renal disease, acute alcohol poisoning, following procedures like endoscopic retrograde cholangiopancreatography (ERCP) and even in non-pancreatic conditions like penetrating peptic ulcers, duodenal obstruction, mumps, ectopic pregnancy, and severe diabetic

NABL certificate
and scope

This test has been performed at
TATA 1MG HYDERABAD
Address: SCB Door No. 3-14-011, 1st Floor,
Patny Square, SP Road, Rasoolpura,
Secunderabad, Telangana - 500003

Dr. Vittal Sri Navya
MBBS, MD (Pathology)
Consultant Pathologist
Reg. No: 85499

Scan for
digital copy



PO No :PO3372032243-231



Customer Name	: Mr.ARVIND	Collected Via	: TATA IMG HYDERABAD
Age/Gender	: 41/Male	Referred By	: Dr.
Lab Visit ID	: HYD572178	Collection Date	: 25/Jun/2025 06:38AM
Barcode ID/Order ID	: D18948383 / 13257890	Report Date	: 25/Jun/2025 07:55PM
Sample Type	: Serum	Report Status	: Final Report

BIOCHEMISTRY

COMPREHENSIVE PLATINUM FULL BODY CHECKUP WITH SMART REPORT

Test Name	Result	Unit	Bio. Ref. Interval	Method
ketoacidosis.				
<ul style="list-style-type: none"> In asymptomatic individuals, elevated amylase levels may be attributed to macroamylasemia or idiopathic hyperamylasemia; amylase levels may fluctuate. Transient increases in amylase may also result from inflammation, alcohol consumption, or medications such as aspirin, diuretics, oral contraceptives, corticosteroids, indomethacin, and opiates. Low amylase levels are seen in chronic pancreatitis, congestive heart failure, 2nd & 3rd trimester of pregnancy, gastrointestinal cancer & bone fractures. Highly lipemic samples may show falsely low amylase levels. 				

Lipoprotein (a)

Lipoprotein(a)	16.05	mg/dL	<30	Latex microparticle-enhanced immunoturbidimetry
----------------	-------	-------	-----	---

Comment:

Note: Lipoprotein(a) [Lp(a)] is considered an important risk factor for Coronary Heart Disease (CHD).

- * Lipoprotein (a) consists of an LDL particle that is covalently bound to an additional protein, apolipoprotein (a). Apo(a) has high-sequence homology with the coagulation factor plasminogen and, like LDL, Lp(a) contains apolipoprotein B100. Thus, Lp(a) is both proatherogenic and prothrombotic. Lp(a) is an independent risk factor for CHD, Ischemic Stroke, and Aortic Valve Stenosis.
- * Lp(a) is highly heterogeneous molecule; the degree of atherogenicity of the Lp(a) particle may depend on the molecular size of the Lp(a)-specific protein.
- * Serum concentrations of Lp(a) are related to genetic factors, and are largely unaffected by diet, exercise and lipid -lowering pharmaceuticals. However, in a patient with additional modifiable CHD risk factors, more aggressive therapy to normalize these factors may be indicated if the Lp(a) value is also increased.

Usage:

Evaluation of increased risk for cardiovascular disease and events:

- * In individuals at intermediate risk for cardiovascular disease
- * In patients with early atherosclerosis
- * In patients with strong family history of early CHD

High Sensitive CRP

High sensitivity CRP	3.44	mg/L	Healthy Individuals: <= 3 Low Risk: < 1 Average Risk: > 1 and < 3 High Risk: > 3	Latex enhanced immunoturbidimetric
----------------------	------	------	---	------------------------------------

NABL certificate and scope



This test has been performed at
TATA 1MG HYDERABAD
Address: SCB Door No. 3-14-011, 1st Floor,
Patny Square, SP Road, Rasoolpura,
Secunderabad, Telangana - 500003

Dr. Vittal Sri Navya
MBBS, MD (Pathology)
Consultant Pathologist
Reg. No: 85499

Scan for digital copy





PO No :PO3372032243-231



Customer Name	: Mr.ARVIND	Collected Via	: TATA IMG HYDERABAD
Age/Gender	: 41/Male	Referred By	: Dr.
Lab Visit ID	: HYD572178	Collection Date	: 25/Jun/2025 06:38AM
Barcode ID/Order ID	: D18948383 / 13257890	Report Date	: 25/Jun/2025 07:55PM
Sample Type	: Serum	Report Status	: Final Report

BIOCHEMISTRY

COMPREHENSIVE PLATINUM FULL BODY CHECKUP WITH SMART REPORT

Test Name	Result	Unit	Bio. Ref. Interval	Method
-----------	--------	------	--------------------	--------

Comment:

High Sensitivity C- Reactive protein (hs-CRP) is used as a marker for determining and performing risk assessment of cardiovascular disease (good marker for inflammation), often along with tests for Lipid profile.

The American Heart Association and US Centers for Disease Control and Prevention have defined risk groups as follows:

- <1.0 Low Risk
- 1.0 - 3.0 - Average Risk
- >3.0 High Risk

These values are only a part of the total evaluation process for cardiovascular diseases.

To assess vascular risk, it is recommended to test hsCRP levels 2 or more weeks apart and calculate the average

Additional risk factors to be considered are elevated levels of lipids & glucose, smoking, high blood pressure (hypertension). Anti inflammatory drugs (like aspirin, ibuprofen, and naproxen) or statins may reduce CRP levels in blood. It is important that any person undergoing this test must be in a healthy state in order for the results to be of diagnostic value in predicting the risk of coronary artery disease or heart attack. Any recent illness, tissue injury, infection, or other general inflammation will raise the amount of hsCRP and give a falsely elevated estimate of risk.

Women on hormone replacement therapy have been shown to have elevated hs-CRP levels.

Note:

Since the hs-CRP and CRP tests measure the same molecule, people with chronic inflammation, such as those with arthritis, should not have hs-CRP levels measured. Their CRP levels will be very high due to the arthritis/often too high to be measured or meaningful using the hs-CRP test.

Apolipoproteins A1 & B

Apolipoprotein - A1	99.00	mg/dL	79-169	PEG immunturbidimetric
Apolipoprotein - B	57.00	mg/dL	46- 174	PEG immunturbidimetric
Apolipoprotein B/A1 Ratio	0.58	Ratio		Calculated

Comment:

Apolipoprotein A1

- Apolipoproteins A1 (Apo A1) is the major apolipoprotein attached to HDL and is found in greater proportion than Apo A2 (3:1).
- It is inversely related to the risk of coronary artery disease (CAD).
- It may be a better predictor of atherogenic risk than HDL.

Apo A1 may be increased with

Apo A1 may be decreased with

This test has been performed at

TATA 1MG HYDERABAD

Address: SCB Door No. 3-14-011, 1st Floor,
Patny Square, SP Road, Rasoolpura,
Secunderabad, Telangana - 500003

Dr. Vittal Sri Navya
MBBS, MD (Pathology)
Consultant Pathologist
Reg. No: 85499

NABL certificate and scope



Scan for digital copy





PO No :PO3372032243-231



Customer Name	: Mr.ARVIND	Collected Via	: TATA IMG HYDERABAD
Age/Gender	: 41/Male	Referred By	: Dr.
Lab Visit ID	: HYD572178	Collection Date	: 25/Jun/2025 06:38AM
Barcode ID/Order ID	: D18948383 / 13257890	Report Date	: 25/Jun/2025 07:55PM
Sample Type	: Serum	Report Status	: Final Report

BIOCHEMISTRY

COMPREHENSIVE PLATINUM FULL BODY CHECKUP WITH SMART REPORT

Test Name	Result	Unit	Bio. Ref. Interval	Method
Drugs (carbamazepine, estrogens, ethanol, statins, niacin, oral contraceptives, phenobarbital)	Chronic renal failure			
Familial hyper alpha-lipoproteinemia	Coronary artery disease and peripheral vascular disease			
Physical exercise	Drugs (androgens, beta blockers, diuretics and progestins)			
Pregnancy	Familial hypo alpha-lipoproteinemia			
Weight reduction	Smoking & Uncontrolled diabetes 2			

Apolipoprotein B

- Apolipoprotein B (Apo B) is a major protein component of low density lipoprotein (LDL), comprising >90% of the LDL. It is a more powerful independent predictor of coronary artery disease (CAD) than LDL cholesterol. It is useful in assessing the risk of CAD and to classify Hyperlipidemias.
- Apolipoprotein studies help in monitoring coronary bypass surgery patients with regard to risk and severity of restenosis. They are also useful in assessing risk of re-infarction in patients with Myocardial infarction.
- In patients with hyperapobetalipoproteinemia (HALB), a disorder associated with increased risk of developing CHD and with an estimated prevalence of 30% in patients with premature CAD, Apo B is increased disproportionately in LDL cholesterol. Apo B quantitation is used in distinguishing HALB from another common lipoprotein abnormality, Familial combined hyperlipidemia.

Apolipoprotein B:A1 Ratio

Elevated ApoB/ApoA1 ratio confers increased risk of atherosclerotic cardiovascular disease independently of LDL and HDL cholesterol concentrations.

Apo B to A1 ratio	
Ratio	Remarks
0.35- 0.98	Desirable
>0.98	Increased CAD risk

C-Reactive Protein Quantitative

C-Reactive Protein (Quantitative)	5.10	mg/L	<10	Latex Immuno Turbidimetric
-----------------------------------	------	------	-----	----------------------------

This test has been performed at

TATA 1MG HYDERABAD

Address: SCB Door No. 3-14-011, 1st Floor,
Patny Square, SP Road, Rasoolpura,
Secunderabad, Telangana - 500003Dr. Vittal Sri Navya
MBBS, MD (Pathology)
Consultant Pathologist
Reg. No: 85499

NABL certificate and scope



Scan for digital copy





PO No :PO3372032243-231



Customer Name	: Mr.ARVIND	Collected Via	: TATA IMG HYDERABAD
Age/Gender	: 41/Male	Referred By	: Dr.
Lab Visit ID	: HYD572178	Collection Date	: 25/Jun/2025 06:38AM
Barcode ID/Order ID	: D18948383 / 13257890	Report Date	: 25/Jun/2025 07:55PM
Sample Type	: Serum	Report Status	: Final Report

BIOCHEMISTRY

COMPREHENSIVE PLATINUM FULL BODY CHECKUP WITH SMART REPORT

Test Name	Result	Unit	Bio. Ref. Interval	Method
-----------	--------	------	--------------------	--------

Comment:

- C-Reactive Protein [CRP] is an acute phase reactant ,hepatic secretion of which is stimulated in response to inflammatory cytokines.
- CRP is a very sensitive but nonspecific marker of inflammation and infection.
- The CRP test is useful in patient with Inflammatory bowel disease, arthritis, Autoimmune diseases, Pelvic inflammatory disease (PID), tissue injury or necrosis and infections.
- CRP levels can be elevated in the later stages of pregnancy as well as with use of birth control pills or hormone replacement therapy i.e. estrogen. Higher levels of CRP have also been observed in the obese.
- As compared to ESR, CRP shows an earlier rise in inflammatory disorders which begins in 4-6 hrs, the intensity of the rise being higher than ESR and the recovery being earlier than ESR. Unlike ESR, CRP levels are not influenced by hematologic conditions like Anemia, Polycythemia.

Rheumatoid Factor - Quantitative

Rheumatoid Factor - Quantitative	4.5	IU/mL	0-14	Turbidimetry
----------------------------------	-----	-------	------	--------------

Comment:

- The detection of Rheumatoid factor (RF) is one of the criteria of the American Rheumatism Association (ARA) for the diagnosis of Rheumatoid Arthritis (RA).
- RF are heterogeneous group of auto antibodies directed against Fc- region of IgG molecules.
- They are useful in diagnosis of Rheumatoid Arthritis, but can also be found in other inflammatory diseases and in various non-rheumatic diseases.
- These occur in all the immunoglobulin classes, although the usual analytical methods are limited to the detection of Rheumatoid Factors of the IgM type. Healthy individuals >65 years of age may also show positive RF results.

NABL certificate
and scope

This test has been performed at
TATA 1MG HYDERABAD
Address: SCB Door No. 3-14-011, 1st Floor,
Patny Square, SP Road, Rasoolpura,
Secunderabad, Telangana - 500003

Dr. Vittal Sri Navya
MBBS, MD (Pathology)
Consultant Pathologist
Reg. No: 85499

Scan for
digital copy

Page 18 of 28

PO No :PO3372032243-231



Customer Name	: Mr.ARVIND	Collected Via	: TATA IMG HYDERABAD
Age/Gender	: 41/Male	Referred By	: Dr.
Lab Visit ID	: HYD572178	Collection Date	: 25/Jun/2025 06:38AM
Barcode ID/Order ID	: D18948385 / 13257890	Report Date	: 25/Jun/2025 05:03PM
Sample Type	: Urine	Report Status	: Final Report

BIOCHEMISTRY

COMPREHENSIVE PLATINUM FULL BODY CHECKUP WITH SMART REPORT

Test Name	Result	Unit	Bio. Ref. Interval	Method
Microalbumin Creatinine Ratio, Urine				
Microalbumin-Albumin	6.07	mg/L	0-30	Turbidimetric
Urinary Creatinine	63.35	mg/dL	22-328	Kinetic Alkaline Picrate
Microalbumin-Albumin/Creatinine Ratio	9.58	mg/g Creatinine	<30	Calculated

Comment:

Microalbumin/Albumin-to-Creatinine Ratio (UACR) Categories

ACR Category	UACR (mg/g creatinine)	Terms
A1	<30	Normal
A2	30 - 299	Microalbuminuria
A3	>= 300	Clinical Albuminuria

Note: ACR categories: A1 - normal to mildly increased; A2 - moderately increased; A3 - severely increased.

(Source- American Diabetes Association (ADA):Standards of Care in Diabetes-2024)

- As per ADA, due to high biological variability (>20%) between measurements of urinary albumin excretion; two out of three specimens collected within a 3-to 6-month period should be abnormal before considering albuminuria (after excluding non-renal causes).
- Certain factors may raise UACR even without kidney damage - **physiological** like exercise within 24 hours, menstruation, pregnancy, benign postural proteinuria or **pathological** like infection (UTI), hematuria, fever, marked hyperglycemia, congestive heart failure, marked hypertension & poor metabolic control. A high albumin-to-creatinine ratio can be due to low urinary creatinine seen in females, low muscle mass, low protein intake or acute kidney injury.
- A random spot urine sample can be used, but due to high variability, it is recommended that abnormal UACR (>= 30 mg/g) should be confirmed with subsequent first morning midstream sample or 24 hr urine collection.
- Due to inherent day to day variability in albumin excretion, UACR is a better indicator than urine albumin alone. Microalbuminuria is defined as the small but abnormal increase in the excretion of urinary albumin (30-300 mg/g creatinine), but it is recommended to use the term albuminuria for ACR >= 30 mg/g creatinine.
- Persistent albuminuria present for a minimum of 3 months is one of the diagnostic markers of kidney damage and used for classification of chronic kidney disease (CKD).

Clinical Utility: Useful in early screening of diabetic nephropathy, as a risk marker for stroke & heart disease and also for classification and progression of CKD.

NABL certificate
and scope

This test has been performed at
TATA 1MG HYDERABAD
Address: SCB Door No. 3-14-011, 1st Floor,
Patny Square, SP Road, Rasoolpura,
Secunderabad, Telangana - 500003

Madhuri

Dr. K Madhuri
MBBS, MD (Pathology)
Consultant Pathologist
Reg. No: 80149

Scan for
digital copy

Page 19 of 28



PO No :PO3372032243-231



Customer Name	: Mr.ARVIND	Collected Via	: TATA IMG HYDERABAD
Age/Gender	: 41/Male	Referred By	: Dr.
Lab Visit ID	: HYD572178	Collection Date	: 25/Jun/2025 06:38AM
Barcode ID/Order ID	: D18948383 / 13257890	Report Date	: 26/Jun/2025 12:44AM
Sample Type	: Serum	Report Status	: Final Report

IMMUNOLOGY

COMPREHENSIVE PLATINUM FULL BODY CHECKUP WITH SMART REPORT

Test Name	Result	Unit	Bio. Ref. Interval	Method
Immunoglobulin E (IgE) Total	202	IU/mL	0 - 158	CLIA

Comment:

- Immunoglobulin E (IgE) is the most important trigger molecule for allergic information.
- As IgE is a mediator of allergic response, quantitative measurement can provide useful information for differential diagnosis of atopic and non-atopic disease.
- The level of IgE is low during the first year of life, gradually increases with age and reaches adult level after 10 years.

Uses

- For Allergy testing.
- Evaluation of children and adults suspected of having allergic respiratory disease
- To confirm clinical expression of sensitivity to foods in patients with Anaphylactic sensitivity or with Asthma, Angioedema or Cutaneous disease.
- To confirm the presence of IgE antibodies to certain occupational allergens

Increased Levels:

Atopic/Non-atopic allergy, Hyper IgE syndrome, Parasitic infections, IgE Myeloma, Bronchopulmonary Aspergillosis, Immunodeficiency states & Autoimmune diseases, Hodgkin's disease, etc.

Decreased Levels:

Hereditary deficiencies, Acquired immunodeficiency, Ataxia Telangiectasia, Non IgE Myeloma

Note:

Normal levels of IgE does not eliminate the possibility of allergic diseases
No close correlation has been demonstrated between severity of allergic reaction and IgE levels.

Thyroid profile Total

T3, Total	1.22	ng/mL	0.60-1.81	CLIA
T4, Total	12.1	µg/dl	4.5-12.6	CLIA
Thyroid Stimulating Hormone - Ultra Sensitive	4.818	uIU/ml	0.55-4.78	CLIA

NABL certificate
and scope

This test has been performed at
TATA 1MG HYDERABAD
Address: SCB Door No. 3-14-011, 1st Floor,
Patny Square, SP Road, Rasoolpura,
Secunderabad, Telangana - 500003

Dr. Vittal Sri Navya
MBBS, MD (Pathology)
Consultant Pathologist
Reg. No: 85499

Scan for
digital copy



PO No :PO3372032243-231



Customer Name	: Mr.AR.VIND	Collected Via	: TATA IMG HYDERABAD
Age/Gender	: 41/Male	Referred By	: Dr.
Lab Visit ID	: HYD572178	Collection Date	: 25/Jun/2025 06:38AM
Barcode ID/Order ID	: D18948383 / 13257890	Report Date	: 26/Jun/2025 12:44AM
Sample Type	: Serum	Report Status	: Final Report

IMMUNOLOGY

COMPREHENSIVE PLATINUM FULL BODY CHECKUP WITH SMART REPORT

Test Name	Result	Unit	Bio. Ref. Interval	Method
-----------	--------	------	--------------------	--------

Comment:

- Below mentioned are the guidelines for pregnancy related reference ranges for TSH, total T3 & Total T4.

Pregnancy			
	TSH (μ IU/mL) (as per American Thyroid Association)	Total T3 (ng/mL)	Total T4(μ g/dL)
1st trimester	0.1-2.5	0.81-1.90	7.33-14.8
2nd trimester	0.2-3.0	1.00-2.60	7.93-16.1
3rd trimester	0.3-3.0	1.00-2.60	6.95-15.7

- TSH levels are subject to circadian variation, reaching peak levels between 2 - 4.a.m. and at a minimum between 6-10 pm .
- The variation is of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.
- TSH is secreted in a dual fashion: Intermittent pulses constitute 60-70% of total amount, background continuous secretion is 30-40%.These pulses occur regularly every 1-3 hrs.
- Total T3 & T4 concentrations are altered by physiological or pathological changes in thyroxine binding globulin (TBG) capacity .
- The determination of free T3 & free T4 has the advantage of being independent of changes in the concentrations and binding properties of the binding proteins.
- Changes in thyroid status are typically associated with concordant changes in T3, T4 and TSH levels.
- Unexpectedly abnormal or discordant thyroid test values may be seen with some rare, but clinically significant conditions such as central hypothyroidism, TSH-secreting pituitary tumors, thyroid hormone resistance, or the presence of heterophilic antibodies (HAMA) or thyroid hormone autoantibodies.
- For diagnostic purposes, results should be used in conjunction with other data.

TSH	T3	T4	Interpretation
High	Normal	Normal	Subclinical Hypothyroidism
Low	Normal	Normal	Subclinical Hyperthyroidism

NABL certificate and scope



This test has been performed at
TATA 1MG HYDERABAD
 Address: SCB Door No. 3-14-011, 1st Floor,
 Patny Square, SP Road, Rasoolpura,
 Secunderabad, Telangana - 500003

Dr. Vittal Sri Navya

Dr. Vittal Sri Navya
 MBBS, MD (Pathology)
 Consultant Pathologist
 Reg. No: 85499

Scan for digital copy





PO No :PO3372032243-231



Customer Name	: Mr.ARVIND	Collected Via	: TATA IMG HYDERABAD
Age/Gender	: 41/Male	Referred By	: Dr.
Lab Visit ID	: HYD572178	Collection Date	: 25/Jun/2025 06:38AM
Barcode ID/Order ID	: D18948383 / 13257890	Report Date	: 26/Jun/2025 12:44AM
Sample Type	: Serum	Report Status	: Final Report

IMMUNOLOGY

COMPREHENSIVE PLATINUM FULL BODY CHECKUP WITH SMART REPORT

Test Name	Result	Unit	Bio. Ref. Interval	Method
High	High	High	Secondary Hyperthyroidism	
Low	High/Normal	High/Normal	Hyperthyroidism	
Low	Low	Low	Non thyroidal illness / Secondary Hypothyroidism	

Free T4 1.23 ng/dL 0.89 - 1.76 CLIA

Comment:

- Below mentioned are the guidelines for pregnancy related reference ranges for free T4.

Pregnancy	Reference Ranges(ng/dL)
1st trimester	0.7-2.0
2nd trimester	0.5-1.6
3rd trimester	0.5-1.6

- FT4 is the biologically active fraction of thyroxine in circulating blood.
- In patients with hyperthyroidism, the FT4 concentration increases, whereas in patients with hypothyroidism it generally decreases.
- Patients on hormone replacement therapy may have an elevation of FT4, although clinically they are euthyroid.
- The determination of free T4 has the advantage of being independent of changes in the concentrations and binding properties of the binding proteins.
- For diagnostic purposes, the results should always be assessed in conjunction with the patient's medical history, clinical examination and other findings.

TSH	T3 /FT3	T4/FT4	Interpretation
High	Normal	Normal	Subclinical Hypothyroidism
Low	Normal	Normal	Subclinical Hyperthyroidism
High	High	High	Secondary Hyperthyroidism
Low	High/Normal	High/Normal	Hyperthyroidism

NABL certificate and scope



This test has been performed at
TATA 1MG HYDERABAD
 Address: SCB Door No. 3-14-011, 1st Floor,
 Patny Square, SP Road, Rasoolpura,
 Secunderabad, Telangana - 500003

Dr. Vittal Sri Navya
 MBBS, MD (Pathology)
 Consultant Pathologist
 Reg. No: 85499

Scan for digital copy





PO No :PO3372032243-231



Customer Name	: Mr.ARVIND	Collected Via	: TATA IMG HYDERABAD
Age/Gender	: 41/Male	Referred By	: Dr.
Lab Visit ID	: HYD572178	Collection Date	: 25/Jun/2025 06:38AM
Barcode ID/Order ID	: D18948383 / 13257890	Report Date	: 26/Jun/2025 12:44AM
Sample Type	: Serum	Report Status	: Final Report

IMMUNOLOGY

COMPREHENSIVE PLATINUM FULL BODY CHECKUP WITH SMART REPORT

Test Name	Result	Unit	Bio. Ref. Interval	Method
Low	Low	Low	Non thyroidal illness / Secondary Hypothyroidism	

Free T3 3.35 pg/mL 2.3 - 4.2 CLIA

Comment:

- Below mentioned are the guidelines for pregnancy related reference ranges for free T3.

Pregnancy	Reference Ranges(pg/mL)
1st trimester	2.0-3.8
2nd trimester	2.0-3.8
3rd trimester	2.0-3.8

- Free T3 measurements support the differential diagnosis of thyroid disorders, are needed to distinguish different forms of hyperthyroidism, to identify patients with T3 thyrotoxicosis, monitoring of patients with hypothyroidism treated with Thyroxine and antithyroid agents.
- The determination of free T3 has the advantage of being independent of changes in the concentrations and binding properties of the binding proteins.
- For diagnostic purposes, the results should always be assessed in conjunction with the patient's medical history, clinical examination and other findings.

TSH	T3 /FT3	T4/FT4	Interpretation
High	Normal	Normal	Subclinical Hypothyroidism
Low	Normal	Normal	Subclinical Hyperthyroidism
High	High	High	Secondary Hyperthyroidism
Low	High/Normal	High/Normal	Hyperthyroidism
Low	Low	Low	Non thyroidal illness / Secondary Hypothyroidism

NABL certificate and scope



This test has been performed at
TATA 1MG HYDERABAD
 Address: SCB Door No. 3-14-011, 1st Floor,
 Patny Square, SP Road, Rasoolpura,
 Secunderabad, Telangana - 500003

Dr. Vittal Sri Navya

MBBS, MD (Pathology)
 Consultant Pathologist
 Reg. No: 85499

Scan for digital copy





PO No :PO3372032243-231



Customer Name	: Mr.ARVIND	Collected Via	: TATA IMG HYDERABAD
Age/Gender	: 41/Male	Referred By	: Dr.
Lab Visit ID	: HYD572178	Collection Date	: 25/Jun/2025 06:38AM
Barcode ID/Order ID	: D18948383 / 13257890	Report Date	: 26/Jun/2025 12:44AM
Sample Type	: Serum	Report Status	: Final Report

IMMUNOLOGY

COMPREHENSIVE PLATINUM FULL BODY CHECKUP WITH SMART REPORT

Test Name	Result	Unit	Bio. Ref. Interval	Method
Vitamin D (25-OH)				
Vitamin D (25-OH)	30.5	ng/mL	Deficiency:< 20, Insufficiency:20-29, Sufficiency:30 - 100, Toxicity possible:> 100	CLIA

Comment:

- Vitamin D is a fat-soluble steroid prohormone involved in the intestinal absorption of calcium and the regulation of calcium homeostasis.
- Two forms of vitamin D are biologically relevant - vitamin D3 (Cholecalciferol) and vitamin D2 (Ergocalciferol).
- Both vitamins D3 and D2 can be absorbed from food but only an estimated 10-20perc. of vitamin D is supplied through nutritional intake.
- Vitamin D is converted to the active hormone 1,25-(OH)2-vitamin D (Calcitriol) through two hydroxylation reactions. The first hydroxylation converts vitamin D into 25-OH vitamin D and occurs in the liver. The second hydroxylation converts 25-OH vitamin D into the biologically active 1,25-(OH)2-vitamin D and occurs in the kidneys as well as in many other cells of the body.
- Most cells express the vitamin D receptor and about 3perc. of the human genome is directly or indirectly regulated by the vitamin D endocrine system.
- The major storage form of vitamin D is 25-OH vitamin D and is present in the blood at up to 1,000 fold higher concentration compared to the active 1,25-(OH)2-vitamin D. 25-OH vitamin D has a half-life of 2-3 weeks vs. 4 hours for 1,25-(OH)2-vitamin D. Therefore, 25-OH vitamin D is the analyte of choice for determination of the vitamin D status.
- Risk factors for vitamin D deficiency include low sun exposure, inadequate intake, decreased absorption, abnormal metabolism, vitamin D resistance and liver or kidney diseases.
- Vitamin D deficiency is a cause of secondary hyperparathyroidism and diseases resulting in impaired bone metabolism (like rickets, osteomalacia).
- Recently, many chronic diseases such as cancer, high blood pressure, osteoporosis and several autoimmune diseases have been linked to vitamin D deficiency.
- The assay measures both D2 (Ergocalciferol) and D3 (Cholecalciferol) metabolites of vitamin D

Utility Quantitative determination of 25-hydroxyvitamin D (25-OH vitamin D).

Vitamin B12

This test has been performed at
TATA 1MG HYDERABAD
Address: SCB Door No. 3-14-011, 1st Floor,
Patny Square, SP Road, Rasoolpura,
Secunderabad, Telangana - 500003

Dr. Vittal Sri Navya
MBBS, MD (Pathology)
Consultant Pathologist
Reg. No: 85499

NABL certificate
and scopeScan for
digital copy



PO No :PO3372032243-231



Customer Name	: Mr.ARVIND	Collected Via	: TATA IMG HYDERABAD
Age/Gender	: 41/Male	Referred By	: Dr.
Lab Visit ID	: HYD572178	Collection Date	: 25/Jun/2025 06:38AM
Barcode ID/Order ID	: D18948383 / 13257890	Report Date	: 26/Jun/2025 12:44AM
Sample Type	: Serum	Report Status	: Final Report

IMMUNOLOGY

COMPREHENSIVE PLATINUM FULL BODY CHECKUP WITH SMART REPORT

Test Name	Result	Unit	Bio. Ref. Interval	Method
Vitamin B12	248.0	pg/ml	211-911	CLIA

Comment:

- Vitamin B12** along with **folate** is essential for DNA synthesis and myelin formation.
- Decreased levels** are seen in anaemia, term pregnancy, vegetarian diet, intrinsic factor deficiency, partial gastrectomy/ileal damage, celiac disease, oral contraceptive use, parasitic infestation, pancreatic deficiency, treated epilepsy, smoking, hemodialysis and advanced age.
- Increased levels** are seen in renal failure, hepatocellular disorders, myeloproliferative disorders and at times with excess supplementation of vitamins pills.

Vitamin B9 (Folic Acid)

Vitamin B9 (Folic Acid)	16.42	ng/ml	0.35-3.37 Deficient 3.38-5.38 Indeterminate >5.38 Normal	CLIA
-------------------------	-------	-------	--	------

Comment:

Folate plays an important role in the synthesis of purine & pyrimidines in the body and is important for the maturation of erythrocytes. It is widely available from plants and to a lesser extent organ meats, but more than half the folate content of food is lost during cooking. Folate deficiency is commonly prevalent in alcoholic liver disease, pregnancy, and the elderly. It may result from poor intestinal absorption, nutrition deficiency, excessive demand as in pregnancy or in malignancy, and in response to certain drugs like Methotrexate & anticonvulsants. It is now routine practice to recommend dietary folate supplements from conception to the 12th week of pregnancy; such supplementation has been proven to reduce the incidence of neural tube defects.

Decreased Levels: Megaloblastic anemia, Infantile hyperthyroidism, Alcoholism, Malnutrition, Scurvy, Liver disease, B12 deficiency, dietary amino acid excess, adult Celiac disease, Tropical Sprue, Crohn's disease, Hemolytic anemias, Carcinomas, Myelofibrosis, vitamin B6 deficiency, pregnancy, Whipple's disease, extensive intestinal resection, and severe exfoliative dermatitis.

Note:

Certain drugs like Pyrimethamine, methotrexate, and trimethoprim are all folate antagonists i.e. they stop the action of the folic acid; phenytoin can decrease the intestinal absorption of folates, and ethanol both decreases absorption and increases excretion of folic acid.

To differentiate vitamin B12 & folate deficiency, measurement of Methylmalonic acid in urine & serum Homocysteine level is suggested.

NABL certificate
and scope

This test has been performed at
TATA 1MG HYDERABAD
Address: SCB Door No. 3-14-011, 1st Floor,
Patny Square, SP Road, Rasoolpura,
Secunderabad, Telangana - 500003

Dr. Vittal Sri Navya
MBBS, MD (Pathology)
Consultant Pathologist
Reg. No: 85499

Scan for
digital copy



PO No :PO3372032243-231



Customer Name	: Mr.ARVIND	Collected Via	: TATA IMG HYDERABAD
Age/Gender	: 41/Male	Referred By	: Dr.
Lab Visit ID	: HYD572178	Collection Date	: 25/Jun/2025 06:38AM
Barcode ID/Order ID	: D18948383 / 13257890	Report Date	: 26/Jun/2025 12:44AM
Sample Type	: Serum	Report Status	: Final Report

IMMUNOLOGY

COMPREHENSIVE PLATINUM FULL BODY CHECKUP WITH SMART REPORT

Test Name	Result	Unit	Bio. Ref. Interval	Method
Homocysteine				
Homocysteine	41.24	μmol/L	<15	CLIA

Comment:

Interpretation:

Increased levels are seen in deranged Vit B12 metabolism and form an independent marker for risk of thromboembolic episodes in coronary artery disease (CAD)

Clinical Utility:

- Determine risk for heart disease, stroke and peripheral arterial blood vessel disease.
- Identify vitamin B12 deficiency or folic acid deficiency.
- Identify homocystinuria

The recommended use of Homocysteine (HCY) to assess risk factor for CAD are

- It is specially useful in young CAD patients (<40 years)
- In known cases of CAD,high HCYlevels should be used as a prognostic marker for CAD events and mortality.
- CAD patients with HCY levels >15 umol/L belong to high risk group.
- Increased HCY levels with low vitamin concentrations should be handled as a potential vitamin deficiency case .

High values of HCY are found in dietary deficiency of folic acid, vitamin B6, or vitamin B12, homocystinuria, chronic liver and renal failure,post menopausal state , hypothyroidism, Alzheimer's disease, various neoplastic disease like cancers of ovary or breast and Acute lymphoblastic leukemia, drugs (anti-anticonvulsants, antibiotics, theophylline, birth control pills, and tamoxifen), alcoholism, smoking or tobacco usage.

Low values may be caused by some medicines or vitamins such as folic acid, vitamin B12, or niacin.

- Please note test values may vary depending on the assay method used.
- CLIA-Chemiluminescent Immunoassay

NABL certificate
and scope

This test has been performed at

TATA 1MG HYDERABAD

Address: SCB Door No. 3-14-011, 1st Floor,
Patny Square, SP Road, Rasoolpura,
Secunderabad, Telangana - 500003

Dr. Vittal Sri Navya
MBBS, MD (Pathology)
Consultant Pathologist
Reg. No: 85499

Scan for
digital copy

Page 26 of 28



PO No :PO3372032243-231



Customer Name	: Mr.ARVIND	Collected Via	: TATA IMG HYDERABAD
Age/Gender	: 41/Male	Referred By	: Dr.
Lab Visit ID	: HYD572178	Collection Date	: 25/Jun/2025 06:38AM
Barcode ID/Order ID	: D18948385 / 13257890	Report Date	: 25/Jun/2025 03:40PM
Sample Type	: Urine	Report Status	: Final Report

CLINICAL PATHOLOGY

COMPREHENSIVE PLATINUM FULL BODY CHECKUP WITH SMART REPORT

Test Name	Result	Unit	Bio. Ref. Interval	Method
Urine Routine & Microscopy				
Urine Routine & Microscopy				
Colour	Pale Yellow		Pale Yellow	Manual
Appearance	Slightly Hazy		Clear	Manual
Specific gravity	1.020		1.003 - 1.035	pKa change
pH	6.0		4.6 - 8.0	Double Indicator
Glucose	Negative		Negative	GOD-POD
Protein	Negative		Negative	Protein Error Principle
Ketones	Negative		Negative	Nitroprusside
Blood	Negative		Negative	Peroxidase
Bilirubin	Negative		Negative	Diazonium
Urobilinogen	Normal		Normal	Ehrlich
Leucocyte Esterase	Positive		Negative	Pyrrole
Nitrite	Positive		Negative	P-arsanilic acid
Pus cells	7-8	/hpf	0-5	Microscopy
Red Blood Cells	Nil	/hpf	0-2	Microscopy
Epithelial cells	2-3	/hpf	Few	Microscopy
Casts	Nil	/lpf	Nil	Microscopy
Crystals	Nil		Nil	Microscopy
Yeast	Nil		Nil	Microscopy
Bacteria	Present		Nil	Microscopy

Comment:

•Note: Pre-test condition to be observed while submitting the sample-first void, mid stream urine, collected in a clean, dry, sterile container is recommended for routine urine analysis, avoid contamination with any discharge from vaginal, urethra, perineum, Avoid prolonged transit time & undue exposure to sunlight.

•During interpretation, points to be considered are Negative nitrite test does not exclude the urinary tract infections. Trace proteinuria can be seen with many physiological conditions like prolonged recumbency, exercise, high protein diet. False positive reactions for bile pigments, proteins, glucose and nitrites can be caused by peroxidase like activity by disinfectants, therapeutic dyes, ascorbic acid and certain drugs. • Urine microscopy is done in centrifuged urine specimens

*** End Of Report ***

NABL certificate and scope



This test has been performed at
TATA 1MG HYDERABAD
Address: SCB Door No. 3-14-011, 1st Floor,
Patny Square, SP Road, Rasoolpura,
Secunderabad, Telangana - 500003

Dr. Vittal Sri Navya
MBBS, MD (Pathology)
Consultant Pathologist
Reg. No: 85499

Scan for digital copy



Page 27 of 28

State of the Art Lab Facilities

Delhi • Gurgaon • Jaipur • Lucknow • Patna • Dehradun • Ahmedabad • Mumbai • Pune • Bangalore
Hyderabad • Chennai • Kolkata • Bhubaneswar • Angul



PO No :PO3372032243-231



Customer Name	: Mr.ARVIND	Collected Via	: TATA IMG HYDERABAD
Age/Gender	: 41/Male	Referred By	: Dr.
Lab Visit ID	: HYD572178	Collection Date	: 25/Jun/2025 06:38AM
Barcode ID/Order ID	: D18948385 / 13257890	Report Date	: 25/Jun/2025 03:40PM
Sample Type	: Urine	Report Status	: Final Report

CLINICAL PATHOLOGY

COMPREHENSIVE PLATINUM FULL BODY CHECKUP WITH SMART REPORT

Test Name	Result	Unit	Bio. Ref. Interval	Method
-----------	--------	------	--------------------	--------

Conditions of Laboratory Testing & Reporting:

Test results released pertain to the sample, as received. Laboratory investigations are only a tool to facilitate in arriving at a diagnosis and should be clinically correlated by the interpreting clinician. Result delays may happen because of unforeseen or uncontrollable circumstances. Test report may vary depending on the assay method used. Test results may show inter-laboratory variations. Test results are not valid for medico-legal purposes. Please mail your queries related to test results to Customer Care mail ID care@1mg.com

Disclaimer: Results relate only to the sample received. Test results marked "BOLD" indicate abnormal results i.e. higher or lower than normal. All lab test results are subject to clinical interpretation by a qualified medical professional. This report cannot be used for any medico-legal purposes. Partial reproduction of the test results is not permitted. Also, TATA 1mg Labs is not responsible for any misinterpretation or misuse of the information. The test reports alone may not be conclusive of the disease/condition, hence clinical correlation is necessary. Reports should be vetted by a qualified doctor only.

TATA 1mg Labs

NABL certificate
and scope

This test has been performed at
TATA 1MG HYDERABAD
Address: SCB Door No. 3-14-011, 1st Floor,
Patny Square, SP Road, Rasoolpura,
Secunderabad, Telangana - 500003

Dr. Vittal Sri Navya
MBBS, MD (Pathology)
Consultant Pathologist
Reg. No: 85499

Scan for
digital copy

ENSURING ACCURACY IN EVERY SINGLE REPORT

Following a 3-step review process:



Advanced systems & cutting-edge technology analyze results with precision

Experienced lab experts and technicians conduct comprehensive reviews

Each report undergoes rigorous medical scrutiny & is signed off by a doctor

Have concerns regarding the report?

Reach out to our care team at **care@1mg.com** or **chat with us**

CONNECT NOW >



Accurate Testing, Assured Quality: Diagnostics Precision



Stringent quality control

Measures meeting international norms of safety guidelines



Cutting-edge technology

Robust healthcare systems equipped with calibrated & well-maintained machinery



Experienced lab staff

30+ medical professionals with a collective experience of 200+ years



Verified test procedures

Highly standardized test procedures following CLSI* guidelines



External assessments

Thorough third-party assessment by authorized experts



Trained phlebotomists

Ensuring smooth sample collection experience & pre-analytical precision

Claim FREE doctor consultation

Consult top doctors from the comfort of your home

SCHEDULE NOW >



Watch how we take care of your sample



Get Second Opinion



Order Medicines



Book Lab Tests



Explore Financial Support & more



EXPLORE NOW >

Need Help?



1800-102-1618

Running Low on Vitamin D?

Choose clinically proven **DePURA**^{by Sanofi™}, ready-to-drink¹

Vitamin D3 with 34% better absorption²



34% Better Absorption^{*2}



Palatable Taste¹



Sugar Free¹

*With DePURA, 34% higher Vitamin D3 levels are achieved as compared to conventional formulation.
Reference: 1. Data on File 2. Marwaha, Raman K., et al. "Efficacy of micellized vs. fat-soluble vitamin D3 supplementation in healthy school children from Northern India." Journal of Pediatric Endocrinology and Metabolism 29.12.

LIMITED OFFER

20% OFF
ON TATA 1MG

SHOP NOW!

use code:
DEP20

MAT-IN-2500204-v1.0.02/25

Are you getting enough OMEGA 3 to protect your heart?

Because 80%^ of people don't!

Choose Tata 1mg Salmon Omega 3



Reduces
Triglyceride Levels



Improves
Good Cholesterol



Reduces
Inflammation



^ Stat KD, et al. progress in lipid research 2016
* Compared with competing products

* Reference from NCBI published article on comparative bioavailability
Optimal requirement of Omega-3 as mentioned by National Institutes of Health, Dietary Supplement Label Database 2015.

Why Choose Us?

50% Higher Absorption^{*}
with Triglyceride form

50% Lower Saturated Fats^{*}



Every batch
Tested and Certified

Upto 30% off

Shop Now