

AGAPORA Image Prompt Instructions

AGAPORA IMAGE PROMPT INSTRUCTIONS

Use this after RBGIA has already computed each egg/chick outcome.
The image API must not calculate genetics.

1. Collect from EACH outcome:
 - SPECIES
 - BASE COLOR
 - VISUAL MUTATION
 - SPLIT AND HIDDEN GENES
2. Insert those four fields into the prompt template.
3. Send the composed prompt to the image API (Pollinations).
4. Save the prompt text to a PDF. Save the returned picture on that egg card.

DRAW: species + base color + visual mutations.
DO NOT DRAW: split and hidden genes.

PROMPT TEMPLATE

AGAPORA OFFSPRING IMAGE PROMPT

This image is a visual representation of ONE computed egg/chick outcome.
Do not invent genetics. Do not add extra mutations. Do not draw hidden genes.

SPECIES: {SPECIES} ({SCIENTIFIC_NAME})
SEX: {SEX}
BASE COLOR: {BASE COLOR}
VISUAL MUTATIONS (DRAW THESE): {VISUAL MUTATION}
SPLIT AND HIDDEN GENES (DO NOT DRAW): {SPLIT AND HIDDEN GENES}

INSTRUCTIONS:

- Draw one realistic adult {SPECIES} ({SCIENTIFIC_NAME}) only.
- Use BASE COLOR as the ground color of the plumage.
- Draw VISUAL MUTATIONS as visible plumage. If none, keep wild-type markings for that species.
- Split and hidden genes must stay invisible. The bird looks normal at those loci.
- Dark eyes unless a visual mutation already requires otherwise.
- Clean studio photograph, single bird, no text, no labels, no extra birds.

EXAMPLE (Fischer cock, green, visual SL Greywing, hidden Opaline and Dilute)

AGAPORA OFFSPRING IMAGE PROMPT

This image is a visual representation of ONE computed egg/chick outcome.
Do not invent genetics. Do not add extra mutations. Do not draw hidden genes.

SPECIES: Fischer's lovebird (Agapornis fischeri)

AGAPORA Image Prompt Instructions

SEX: Male / Cock

BASE COLOR: Green

VISUAL MUTATIONS (DRAW THESE): SL Greywing

SPLIT AND HIDDEN GENES (DO NOT DRAW): Opaline, Dilute

INSTRUCTIONS:

- Draw one realistic adult Fischer's lovebird (*Agapornis fischeri*) only.
- Use BASE COLOR as the ground color of the plumage.
- Draw VISUAL MUTATIONS as visible plumage. If none, keep wild-type markings for that species.
- Split and hidden genes must stay invisible. The bird looks normal at those loci.
- Dark eyes unless a visual mutation already requires otherwise.
- Clean studio photograph, single bird, no text, no labels, no extra birds.

SETUP (OpenRouter image generation)

1. In backend/.env set OFFSPRING_IMAGE_ENABLED=true
2. OFFSPRING_IMAGE_PROVIDER=openrouter
3. Set OPENROUTER_API_KEY from <https://openrouter.ai/keys>
4. php artisan storage:link
5. Restart Laravel, then Analyze a pair.
6. Each egg collects SPECIES, BASE COLOR, VISUAL MUTATION, SPLIT AND HIDDEN GENES.
7. AGAPORA composes the prompt, calls OpenRouter, stores the PNG, and writes this PDF.
8. If OpenRouter credits run out, Hugging Face is used as fallback.