# STAT 200:

# Introduction to Data Science with



Descriptive Statistics and Visualizing Distributions

San Diego State University

000011

**Department | Mathematics and Statistics Division** | Statistics and Data Science

# **Descriptive Statistics**


# **Descriptive Statistics**

The first step in any data analysis is to gain an understanding of the data itself

head(murders, 10)					
	А	data.frai	me: 10 × 5		
	state	abb	region	population	total
	<chr></chr>	<chr></chr>	<fct></fct>	<dbl></dbl>	<db1></db1>
1	Alabama	AL	South	4779736	135
2	Alaska	AK	West	710231	19
3	Arizona	AZ	West	6392017	232
4	Arkansas	AR	South	2915918	93
5	California	CA	West	37253956	1257
6	Colorado	со	West	5029196	65
7	Connecticut	СТ	Northeast	3574097	97
8	Delaware	DE	South	897934	38
9	District of Columbia	DC	South	601723	99
10	Florida	FL	South	19687653	669

# **Descriptive Statistics**

- The first step in any data analysis is to gain an understanding of the data itself
- We do this by loading our data into some program (e.g., R, Excel) and exploring it various attributes
  - Number of observations
  - Number of variables
  - Identify errors in data entry
  - Identify missing values
  - etc.



- Once we have an initial understanding of our data, we can then perform basic data analysis through both
  - 1. Descriptive statistics
  - 2. Data visualizations

- Once we have an initial understanding of our data, we can then perform basic data analysis through both
  - 1. Descriptive statistics
  - 2. Data visualizations
- What are descriptive statistics?

- Once we have an initial understanding of our data, we can then perform basic data analysis through both
  - 1. Descriptive statistics
  - 2. Data visualizations
- What are descriptive statistics?
  - Measurements that summarize a given dataset

- Once we have an initial understanding of our data, we can then perform basic data analysis through both
  - 1. Descriptive statistics
  - 2. Data visualizations
- What are descriptive statistics?
  - Measurements that summarize a given dataset
- We use different descriptive statistics and data visualizations for different types of data

- Once we have an initial understanding of our data, we can then perform basic data analysis through both
  - 1. Descriptive statistics
  - 2. Data visualizations
- What are descriptive statistics?
  - Measurements that summarize a given dataset
- We use different descriptive statistics and data visualizations for different types of data
  - Numeric data

- Once we have an initial understanding of our data, we can then perform basic data analysis through both
  - 1. Descriptive statistics
  - 2. Data visualizations
- What are descriptive statistics?
  - Measurements that summarize a given dataset
- We use different descriptive statistics and data visualizations for different types of data
  - Numeric data
  - Categorical data

#### What is numeric data?

head(murders, 10)					
	А	data.frai	me: 10 × 5		
	state	abb	region	population	total
	<chr>&gt;</chr>	<chr></chr>	<fct></fct>	<dbl></dbl>	<db1></db1>
1	Alabama	AL	South	4779736	135
2	Alaska	AK	West	710231	19
3	Arizona	AZ	West	6392017	232
4	Arkansas	AR	South	2915918	93
5	California	CA	West	37253956	1257
6	Colorado	со	West	5029196	65
7	Connecticut	СТ	Northeast	3574097	97
8	Delaware	DE	South	897934	38
9	District of Columbia	DC	South	601723	99
10	Florida	FL	South	19687653	669


#### What is numeric data?

Data consisting of numbers that are either discrete or continuous

head(murders, 10)						
	A data.frame: 10 × 5					
	state	abb	region	population	total	
	<chr></chr>	<chr></chr>	<fct></fct>	<dbl></dbl>	<db1></db1>	
1	Alabama	AL	South	4779736	135	
2	Alaska	AK	West	710231	19	
3	Arizona	AZ	West	6392017	232	
4	Arkansas	AR	South	2915918	93	
5	California	CA	West	37253956	1257	
6	Colorado	со	West	5029196	65	
7	Connecticut	СТ	Northeast	3574097	97	
8	Delaware	DE	South	897934	38	
9	District of Columbia	DC	South	601723	99	
10	Florida	FL	South	19687653	669	


#### What is numeric data?

- Data consisting of numbers that are either discrete or continuous
- Discrete data consist of numeric values that are distinct and countable, typically integer-valued data
  - e.g., population
  - e.g., number of students in a classroom



#### What is numeric data?

- Data consisting of numbers that are either discrete or continuous
- Discrete data consist of numeric values that are distinct and countable, typically integer-valued data
  - e.g., population
  - e.g., number of students in a classroom
- Continuous numeric data is quantitative data, typically represented by a fraction or decimal
  - e.g., temperature of 98.6 degrees Fahrenheit
  - e.g., time of 2.84 seconds




head	head(murders, 10)					
	А	data.frai	me: 10 × 5			
	state	abb	region	population	total	
	<chr>&gt;</chr>	<chr>&gt;</chr>	<fct></fct>	<dbl></dbl>	<db1></db1>	
1	Alabama	AL	South	4779736	135	
2	Alaska	AK	West	710231	19	
3	Arizona	AZ	West	6392017	232	
4	Arkansas	AR	South	2915918	93	
5	California	CA	West	37253956	1257	
6	Colorado	СО	West	5029196	65	
7	Connecticut	СТ	Northeast	3574097	97	
8	Delaware	DE	South	897934	38	
9	District of Columbia	DC	South	601723	99	
10	Florida	FL	South	19687653	669	

#### What is categorical data?


Data divided into distinct groups or categories

head	head(murders, 10)					
	А	data.fra	me: 10 × 5			
	state	abb	region	population	total	
	<chr></chr>	<chr></chr>	<fct></fct>	<dbl></dbl>	<db1></db1>	
1	Alabama	AL	South	4779736	135	
2	Alaska	AK	West	710231	19	
3	Arizona	AZ	West	6392017	232	
4	Arkansas	AR	South	2915918	93	
5	California	CA	West	37253956	1257	
6	Colorado	co	West	5029196	65	
7	Connecticut	СТ	Northeast	3574097	97	
8	Delaware	DE	South	897934	38	
9	District of Columbia	DC	South	601723	99	
10	Florida	FL	South	19687653	669	


- Data divided into distinct groups or categories
- Two types of categorical data
  - Nominal
  - Ordinal



- Data divided into distinct groups or categories
- Two types of categorical data
  - Nominal
  - Ordinal
- Nominal categorical data consist of categories that have no inherent ordering or ranking
  - e.g., states, region




- Data divided into distinct groups or categories
- Two types of categorical data
  - Nominal
  - Ordinal
- Nominal categorical data consist of categories that have no inherent ordering or ranking
  - e.g., states, region
- Ordinal categorical data consist of categories that have inherent ordering or ranking
  - e.g., education level (high school, bachelor's, master's, PhD)
  - e.g., mood (1-sad, 2-neutral, 3-happy)



#### What is categorical data?

- Data divided into distinct groups or categories
- Two types of categorical data
  - Nominal
  - Ordinal
- Nominal categorical data consist of categories that have no inherent ordering or ranking
  - e.g., states, region
- Ordinal categorical data consist of categories that have inherent ordering or ranking
  - e.g., education level (high school, bachelor's, master's, PhD)
  - e.g., mood (1-sad, 2-neutral, 3-happy)



Note that discrete numeric data can be considered ordinal!

- For numeric data, there are two common types of descriptive statistics
  - Measures of central tendency
  - Measures of variability (how things change or vary)

- For numeric data, there are two common types of descriptive statistics
  - Measures of central tendency
  - Measures of variability (how things change or vary)

- Measures of central tendency
  - Mean (average)
  - Median (50<sup>th</sup> percentile, 0.50 quantile)
  - Mode (value that occurs most often)

- For numeric data, there are two common types of descriptive statistics
  - Measures of central tendency
  - Measures of variability (how things change or vary)

- Measures of central tendency
  - Mean (average)
  - Median (50<sup>th</sup> percentile, 0.50 quantile)
  - Mode (value that occurs most often)

- Measures of variability
  - Range
  - Variance
  - Standard deviation
  - Quantiles/Percentiles

- For numeric data, there are two common types of descriptive statistics
  - Measures of central tendency
  - Measures of variability (how things change or vary)

- Measures of central tendency
  - Mean (average)
  - Median (50<sup>th</sup> percentile, 0.50 quantile)
  - Mode (value that occurs most often)

- Measures of variability
  - Range
  - Variance
  - Standard deviation
  - Quantiles/Percentiles

These were likely covered in your high school courses, but I will review them here!


# **Descriptive Statistics:**

Measures of Central Tendency

Consider the following data on the amount of emphysema in the lungs

Patient	Percentage Emphysema
1	5
2	23
3	2
4	14
5	17
6	14

- Consider the following data on the amount of emphysema in the lungs
- We can represent the percentage emphysema on the number line



Patient	Percentage Emphysema
1	5
2	23
3	2
4	14
5	17
6	14

- Consider the following data on the amount of emphysema in the lungs
- We can represent the percentage emphysema on the number line




The mean is a measure of the center of these observations

mean = sum of values / number of values

Patient	Percentage Emphysema
1	5
2	23
3	2
4	14
5	17
6	14

- Consider the following data on the amount of emphysema in the lungs
- We can represent the percentage emphysema on the number line




The mean is a measure of the center of these observations

mean = sum of values / number of values

What is the mean of the variable percentage emphysema?

Patient	Percentage Emphysema
1	5
2	23
3	2
4	14
5	17
6	14

- Consider the following data on the amount of emphysema in the lungs
- We can represent the percentage emphysema on the number line



• The mean is a measure of the center of these observations

What is the mean of the variable percentage emphysema?

$$(5 + 23 + 2 + 14 + 17 + 14) / 6 = 12.5\%$$

Patient	Percentage Emphysema
1	5
2	23
3	2
4	14
5	17
6	14

- Consider the following data on the amount of emphysema in the lungs
- We can represent the percentage emphysema on the number line




The mean is a measure of the center of these observations

What is the mean of the variable percentage emphysema?

$$(5 + 23 + 2 + 14 + 17 + 14) / 6 = 12.5\%$$


Patient	Percentage Emphysema
1	5
2	23
3	2
4	14
5	17
6	14

- Consider the following data on the amount of emphysema in the lungs
- We can represent the percentage emphysema on the number line



Patient	Percentage Emphysema
1	5
2	23
3	2
4	14
5	17
6	14


- Consider the following data on the amount of emphysema in the lungs
- We can represent the percentage emphysema on the number line



 The median is the value such that half the data is above and half the data is below

Patient	Percentage Emphysema
1	5
2	23
3	2
4	14
5	17
6	14


- Consider the following data on the amount of emphysema in the lungs
- We can represent the percentage emphysema on the number line



- The median is the value such that half the data is above and half the data is below
- To find the median
  - 1. Rank the data 2, 5, 14, 14, 17, 23

Patient	Percentage Emphysema
1	5
2	23
3	2
4	14
5	17
6	14


- Consider the following data on the amount of emphysema in the lungs
- We can represent the percentage emphysema on the number line



- The median is the value such that half the data is above and half the data is below
- To find the median
  - 1. Rank the data 2, 5, 14, 14, 17, 23
  - 2. Find the "middle" number
    - If there is an even number of data points, average the two middle numbers

Patient	Percentage Emphysema
1	5
2	23
3	2
4	14
5	17
6	14


- Consider the following data on the amount of emphysema in the lungs
- We can represent the percentage emphysema on the number line



- The median is the value such that half the data is above and half the data is below
- To find the median
  - 1. Rank the data 2, 5, 14, 14, 17, 23
  - 2. Find the "middle" number
    - If there is an even number of data points, average the two middle numbers
- The median is 14
  - Half the data points are greater than 14 and half are less than 14


Patient	Percentage Emphysema
1	5
2	23
3	2
4	14
5	17
6	14

- Consider the following data on the amount of emphysema in the lungs
- We can represent the percentage emphysema on the number line



Patient	Percentage Emphysema
1	5
2	23
3	2
4	14
5	17
6	14


- Consider the following data on the amount of emphysema in the lungs
- We can represent the percentage emphysema on the number line



• The mode is the value that appears most often

Patient	Percentage Emphysema
1	5
2	23
3	2
4	14
5	17
6	14


- Consider the following data on the amount of emphysema in the lungs
- We can represent the percentage emphysema on the number line



- The mode is the value that appears most often
- To find the mode
  - 1. Count the occurrence of each value
  - 2. Choose the value with the largest occurrence

Patient	Percentage Emphysema
1	5
2	23
3	2
4	14
5	17
6	14


- Consider the following data on the amount of emphysema in the lungs
- We can represent the percentage emphysema on the number line



- The mode is the value that appears most often
- To find the mode
  - 1. Count the occurrence of each value
  - 2. Choose the value with the largest occurrence
- What is the mode of percentage emphysema?

Patient	Percentage Emphysema
1	5
2	23
3	2
4	14
5	17
6	14

- Consider the following data on the amount of emphysema in the lungs
- We can represent the percentage emphysema on the number line




- The mode is the value that appears most often
- To find the mode
  - 1. Count the occurrence of each value
  - 2. Choose the value with the largest occurrence
- What is the mode of percentage emphysema?
  - 14 since this value occurs 2 times
  - All other values occur less than 2 times

Patient	Percentage Emphysema
1	5
2	23
3	2
4	14
5	17
6	14

# **Descriptive Statistics:**

Measures of Variability


- Consider the following data on the amount of emphysema in the lungs
- We can represent the percentage emphysema on the number line



• The <u>range</u> is the maximum – minimum

Patient	Percentage Emphysema
1	5
2	23
3	2
4	14
5	17
6	14


- Consider the following data on the amount of emphysema in the lungs
- We can represent the percentage emphysema on the number line



- The <u>range</u> is the maximum minimum
- The range measures the dispersion (spread) of the data

Patient	Percentage Emphysema
1	5
2	23
3	2
4	14
5	17
6	14


- Consider the following data on the amount of emphysema in the lungs
- We can represent the percentage emphysema on the number line



- The <u>range</u> is the maximum minimum
- The range measures the dispersion (spread) of the data
- What is the range of percentage emphysema?

Patient	Percentage Emphysema
1	5
2	23
3	2
4	14
5	17
6	14

- Consider the following data on the amount of emphysema in the lungs
- We can represent the percentage emphysema on the number line




- The <u>range</u> is the maximum minimum
- The range measures the dispersion (spread) of the data
- What is the range of percentage emphysema?

range = 
$$max - min = 23 - 2 = 21\%$$

Patient	Percentage Emphysema
1	5
2	23
3	2
4	14
5	17
6	14


• <u>Variance</u> is another measure of dispersion

Patient	%emph	%emph - mean	(%emph – mean)^2
1	5	-7.5	56.25
2	23	10.5	110.25
3	2	-10.5	110.25
4	14	1.5	2.25
5	17	4.5	20.25
6	14	1.5	2.25
		Total:	301.5
		Total/(6-1):	60.3




- **Variance** is another measure of dispersion
- Unlike range, variance is calculated from all data points (more informative)

Patient	%emph	%emph - mean	(%emph – mean)^2
1	5	-7.5	56.25
2	23	10.5	110.25
3	2	-10.5	110.25
4	14	1.5	2.25
5	17	4.5	20.25
6	14	1.5	2.25
		Total:	301.5
		Total/(6-1):	60.3




- <u>Variance</u> is another measure of dispersion
- Unlike range, variance is calculated from all data points (more informative)
- Variance is the average sum of squared distances from the mean

Patient	%emph	%emph - mean	(%emph – mean)^2
1	5	-7.5	56.25
2	23	10.5	110.25
3	2	-10.5	110.25
4	14	1.5	2.25
5	17	4.5	20.25
6	14	1.5	2.25
		Total:	301.5
		Total/(6-1):	60.3




- **Variance** is another measure of dispersion
- Unlike range, variance is calculated from all data points (more informative)
- Variance is the average sum of squared distances from the mean
  - 1. Subtract the mean from each point

Patient	%emph	%emph - mean	(%emph – mean)^2
1	5	-7.5	56.25
2	23	10.5	110.25
3	2	-10.5	110.25
4	14	1.5	2.25
5	17	4.5	20.25
6	14	1.5	2.25
		Total:	301.5
		Total/(6-1):	60.3




- **Variance** is another measure of dispersion
- Unlike range, variance is calculated from all data points (more informative)
- Variance is the average sum of squared distances from the mean
  - 1. Subtract the mean from each point
  - 2. Square each difference

Patient	%emph	%emph - mean	(%emph – mean)^2
1	5	-7.5	56.25
2	23	10.5	110.25
3	2	-10.5	110.25
4	14	1.5	2.25
5	17	4.5	20.25
6	14	1.5	2.25
		Total:	301.5
		Total/(6-1):	60.3




- <u>Variance</u> is another measure of dispersion
- Unlike range, variance is calculated from all data points (more informative)
- Variance is the average sum of squared distances from the mean
  - 1. Subtract the mean from each point
  - 2. Square each difference
  - 3. Add the squared differences

Patient	%emph	%emph - mean	(%emph – mean)^2
1	5	-7.5	56.25
2	23	10.5	110.25
3	2	-10.5	110.25
4	14	1.5	2.25
5	17	4.5	20.25
6	14	1.5	2.25
		Total:	301.5
		Total/(6-1):	60.3



- **Variance** is another measure of dispersion
- Unlike range, variance is calculated from all data points (more informative)
- Variance is the average sum of squared distances from the mean
  - 1. Subtract the mean from each point
  - 2. Square each difference
  - 3. Add the squared differences
  - 4. Divide the sum by the number of points minus one

Patient	%emph	%emph - mean	(%emph – mean)^2
1	5	-7.5	56.25
2	23	10.5	110.25
3	2	-10.5	110.25
4	14	1.5	2.25
5	17	4.5	20.25
6	14	1.5	2.25
		Total:	301.5
		Total/(6-1):	60.3




# **Descriptive Statistics – Numeric Data – Standard Deviation**

- **Standard deviation** is the square root of the variance
- Since variance squares the units (%^2 in this case), we take square root to convert back to original units (%)

$$SD = \sqrt{Var} = \sqrt{60.3} = 7.77\%$$


Patient	%emph	%emph - mean	(%emph – mean)^2
1	5	-7.5	56.25
2	23	10.5	110.25
3	2	-10.5	110.25
4	14	1.5	2.25
5	17	4.5	20.25
6	14	1.5	2.25
		Total:	301.5
		Total/(6-1):	60.3
		$SD = \sqrt{60.3}$ :	7.77



What happens to the variance when points are further from the mean?


- A) Variance is larger
- B) Variance is smaller
- C) Variance stays the same

Patient	%emph	%emph - mean	(%emph – mean)^2
1	5	-7.5	56.25
2	23	10.5	110.25
3	2	-10.5	110.25
4	14	1.5	2.25
5	17	4.5	20.25
6	14	1.5	2.25
		Total:	301.5
		Total/(6-1):	60.3
		$SD = \sqrt{60.3}$ :	7.77




 Recap: The median is the value such that half the data is above and half the data is below

Patient	Percentage Emphysema
1	5
2	23
3	2
4	14
5	17




- Recap: The median is the value such that half the data is above and half the data is below
- i.e. 50% of data falls below the median and 50% falls above the median

Patient	Percentage Emphysema
1	5
2	23
3	2
4	14
5	17




- Recap: The median is the value such that half the data is above and half the data is below
- i.e. 50% of data falls below the median and 50% falls above the median
- Therefore, other names for the median are the
  - 50% percentile
  - 0.5 quantile

Patient	Percentage Emphysema
1	5
2	23
3	2
4	14
5	17




- Recap: The median is the value such that half the data is above and half the data is below
- i.e. 50% of data falls below the median and 50% falls above the median
- Therefore, other names for the median are the
  - 50% percentile
  - 0.5 quantile
- The p<sup>th</sup> percentile is the value where
  - p% of data falls below the p<sup>th</sup> percentile
  - (1-p)% of data fall above the p<sup>th</sup> percentile

Patient	Percentage Emphysema
1	5
2	23
3	2
4	14
5	17




- Two common percentiles are the
  - 25<sup>th</sup> percentile or "lower quartile"
  - 75<sup>th</sup> percentile or "upper quartile"

Patient	Percentage Emphysema
1	5
2	23
3	2
4	14
5	17




- Two common percentiles are the
  - 25<sup>th</sup> percentile or "lower quartile"
  - 75<sup>th</sup> percentile or "upper quartile"
- For the emphysema table on the right,
  - What is the lower quartile?

Patient	Percentage Emphysema
1	5
2	23
3	2
4	14
5	17




- Two common percentiles are the
  - 25<sup>th</sup> percentile or "lower quartile"
  - 75<sup>th</sup> percentile or "upper quartile"
- For the emphysema table on the right,
  - What is the lower quartile? 5

Patient	Percentage Emphysema
1	5
2	23
3	2
4	14
5	17




- Two common percentiles are the
  - 25<sup>th</sup> percentile or "lower quartile"
  - 75<sup>th</sup> percentile or "upper quartile"
- For the emphysema table on the right,
  - What is the lower quartile? 5
    - That is, 25% of the data falls below 5
    - 75% of the data falls above 5

Patient	Percentage Emphysema
1	5
2	23
3	2
4	14
5	17




- Two common percentiles are the
  - 25<sup>th</sup> percentile or "lower quartile"
  - 75<sup>th</sup> percentile or "upper quartile"
- For the emphysema table on the right,
  - What is the lower quartile? 5
    - That is, 25% of the data falls below 5
    - 75% of the data falls above 5
  - What is the upper quartile?

Patient	Percentage Emphysema
1	5
2	23
3	2
4	14
5	17




- Two common percentiles are the
  - 25<sup>th</sup> percentile or "lower quartile"
  - 75<sup>th</sup> percentile or "upper quartile"
- For the emphysema table on the right,
  - What is the lower quartile? 5
    - That is, 25% of the data falls below 5
    - 75% of the data falls above 5
  - What is the upper quartile? 17

Patient	Percentage Emphysema
1	5
2	23
3	2
4	14
5	17



- Two common percentiles are the
  - 25<sup>th</sup> percentile or "lower quartile"
  - 75<sup>th</sup> percentile or "upper quartile"
- For the emphysema table on the right,
  - What is the lower quartile?
    - That is, 25% of the data falls below 5
    - 75% of the data falls above 5
  - What is the upper quartile? 17
    - That is, 75% of the data falls below 17
    - 25% of the data falls above 17


Patient	Percentage Emphysema
1	5
2	23
3	2
4	14
5	17



- Two common percentiles are the
  - 25<sup>th</sup> percentile or "lower quartile"
  - 75<sup>th</sup> percentile or "upper quartile"
- For the emphysema table on the right,
  - What is the lower quartile? 5
    - That is, 25% of the data falls below 5
    - 75% of the data falls above 5
  - What is the upper quartile? 17
    - That is, 75% of the data falls below 17
    - 25% of the data falls above 17

Patient	Percentage Emphysema
1	5
2	23
3	2
4	14
5	17


Note there are variations for calculating percentiles



- Two common percentiles are the
  - 25<sup>th</sup> percentile or "lower quartile"
  - 75<sup>th</sup> percentile or "upper quartile"
- For the emphysema table on the right,
  - What is the lower quartile? 5
    - That is, 25% of the data falls below 5
    - 75% of the data falls above 5
  - What is the upper quartile? 17
    - That is, 75% of the data falls below 17
    - 25% of the data falls above 17

Patient	Percentage Emphysema
1	5
2	23
3	2
4	14
5	17


- Note there are variations for calculating percentiles
- For this class, we will use R to calculate percentiles for us



- Two common percentiles are the
  - 25<sup>th</sup> percentile or "lower quartile"
  - 75<sup>th</sup> percentile or "upper quartile"
- For the emphysema table on the right,
  - What is the lower quartile?
    - That is, 25% of the data falls below 5
    - 75% of the data falls above 5
  - What is the upper quartile? 17
    - That is, 75% of the data falls below 17
    - 25% of the data falls above 17

Patient	Percentage Emphysema
1	5
2	23
3	2
4	14
5	17

- Note there are variations for calculating percentiles
- For this class, we will use R to calculate percentiles for us
- I primarily want you to know how to interpret these values



# **Descriptive Statistics:**

**Categorical Variables** 

# **Descriptive Statistics – Categorical Data**

- Summarization of categorical data is much easier!
- We simply count the frequency of each category

# **Descriptive Statistics – Categorical Data**

- Summarization of categorical data is much easier!
- We simply count the frequency of each category
- The table of counts is referred as a
  - Frequency tableOR
  - Contingency table

<b>Descriptive Statistics –</b>	<b>Categorical Data</b>
---------------------------------	-------------------------

- Summarization of categorical data is much easier!
- We simply count the frequency of each category
- The table of counts is referred as a
  - Frequency table

OR

- Contingency table
- For example, we can categorize % emphysema into three ordinal categories
  - >0, ≤10
  - >10, ≤ 20
  - >20

Patient	Percentage Emphysema	Emphysema Category
1	5	>0, ≤10
2	23	>20
3	2	>0, ≤10
4	14	>10, ≤ 20
5	17	>10, ≤ 20
6	14	>10, ≤ 20

### **Descriptive Statistics – Categorical Data**

- Summarization of categorical data is much easier!
- We simply count the frequency of each category
- The table of counts is referred as a
  - Frequency tableOR
  - Contingency table
- For example, we can categorize % emphysema into three ordinal categories
  - >0, ≤10
  - >10, ≤ 20
  - >20
- We then count the number of observations per category

Patient	Percentage Emphysema	Emphysema Category
1	5	>0, ≤10
2	23	>20
3	2	>0, ≤10
4	14	>10, ≤ 20
5	17	>10, ≤ 20
6	14	>10, ≤ 20

#### Frequency Table

Emphysema Category	Frequency
>0, ≤10	2
>10, ≤ 20	3
>20	1

# Distributions and Histograms

#### **Measures of central tendency**

- Mean (average)
- Median (50<sup>th</sup> percentile, 0.50 quantile)
- Mode (value that occurs most often)

#### Measures of variability

- Range
- Variance
- Standard deviation
- Quantiles/Percentiles

 These descriptive statistics describe the attributes/shape of what we call the <u>data</u> distribution

#### **Measures of central tendency**

- Mean (average)
- Median (50<sup>th</sup> percentile, 0.50 quantile)
- Mode (value that occurs most often)

#### Measures of variability

- Range
- Variance
- Standard deviation
- Quantiles/Percentiles

- These descriptive statistics describe the attributes/shape of what we call the <u>data</u> distribution
- The data distribution describes how often values in your data occur

#### **Measures of central tendency**

- Mean (average)
- Median (50<sup>th</sup> percentile, 0.50 quantile)
- Mode (value that occurs most often)

#### Measures of variability

- Range
- Variance
- Standard deviation
- Quantiles/Percentiles

- These descriptive statistics describe the attributes/shape of what we call the <u>data</u> distribution
- The data distribution describes how often values in your data occur
- The frequency in which data occurs determines the shape of the distribution

#### **Measures of central tendency**

- Mean (average)
- Median (50<sup>th</sup> percentile, 0.50 quantile)
- Mode (value that occurs most often)

#### Measures of variability

- Range
- Variance
- Standard deviation
- Quantiles/Percentiles

Why do we care?

#### **Measures of central tendency**

- Mean (average)
- Median (50<sup>th</sup> percentile, 0.50 quantile)
- Mode (value that occurs most often)

#### Measures of variability

- Range
- Variance
- Standard deviation
- Quantiles/Percentiles

#### Why do we care?

 By summarizing the data distribution using descriptive statistics, we can

#### **Measures of central tendency**

- Mean (average)
- Median (50<sup>th</sup> percentile, 0.50 quantile)
- Mode (value that occurs most often)

#### Measures of variability

- Range
- Variance
- Standard deviation
- Quantiles/Percentiles

#### Why do we care?

- By summarizing the data distribution using descriptive statistics, we can
  - Better understand data within a group
    - e.g., salary, survival for cancer patients

#### Measures of central tendency

- Mean (average)
- Median (50<sup>th</sup> percentile, 0.50 quantile)
- Mode (value that occurs most often)

#### Measures of variability

- Range
- Variance
- Standard deviation
- Quantiles/Percentiles

#### Why do we care?

- By summarizing the data distribution using descriptive statistics, we can
  - Better understand data within a group
    - e.g., salary, survival for cancer patients
  - and compare data between groups
    - e.g., salary by gender, survival for different cancer treatments, etc.

#### Measures of central tendency

- Mean (average)
- Median (50<sup>th</sup> percentile, 0.50 quantile)
- Mode (value that occurs most often)

#### Measures of variability

- Range
- Variance
- Standard deviation
- Quantiles/Percentiles

#### Why do we care?

- By summarizing the data distribution using descriptive statistics, we can
  - Better understand data within a group
    - e.g., salary, survival for cancer patients
  - and compare data between groups
    - e.g., salary by gender, survival for different cancer treatments, etc.

Comparing groups using information about their data distributions is **statistics!** 

 Data distributions are most commonly visualized using histograms

- Data distributions are most commonly visualized using histograms
- Histograms describe how often values in your data occur

- Data distributions are most commonly visualized using histograms
- Histograms describe how often values in your data occur
- Consider the following data set:
  - You collect data on 9 students on the number of years a student has played an instrument

Person	Years Playing Music
1	8
2	6
3	5
4	7
5	6
6	7
7	8
8	7
9	9

- Data distributions are most commonly visualized using histograms
- Histograms describe how often values in your data occur
- Consider the following data set:
  - You collect data on 9 students on the number of years a student has played an instrument
  - How many times does 5 occur?


Person	Years Playing Music
1	8
2	6
3	5
4	7
5	6
6	7
7	8
8	7
9	9

- Data distributions are most commonly visualized using histograms
- Histograms describe how often values in your data occur
- Consider the following data set:
  - You collect data on 9 students on the number of years a student has played an instrument
  - How many times does 5 occur?
  - How many times does 7 occur?

Person	Years Playing Music
1	8
2	6
3	5
4	7
5	6
6	7
7	8
8	7
9	9


- Data distributions are most commonly visualized using histograms
- Histograms describe how often values in your data occur
- Consider the following data set:
  - You collect data on 9 students on the number of years a student has played an instrument
  - How many times does 5 occur?
  - How many times does 7 occur?
- The plot of these frequencies vs the data values is a <u>histogram</u>

Person	Years Playing Music
1	8
2	6
3	5
4	7
5	6
6	7
7	8
8	7
9	9




 But what if the data are continuous (decimals) and not discrete (integer-valued)?

Person	Years Playing Music
1	8.2
2	6.1
3	5.5
4	7.8
5	6.4
6	7.6
7	8.2
8	7.3
9	9.9




- But what if the data are continuous (decimals) and not discrete (integervalued)?
- Not very informative!

Person	Years Playing Music
1	8.2
2	6.1
3	5.5
4	7.8
5	6.4
6	7.6
7	8.2
8	7.3
9	9.9




- But what if the data are continuous (decimals) and not discrete (integer-valued)?
- Not very informative!
- In this case, we place the data in "bins" to visualize the distribution
  - e.g., 4-5, 5-6, 6-7

Years Playing Music	Bins
8.2	8-9
6.1	6-7
5.5	5-6
7.8	7-8
6.4	6-7
7.6	7-8
8.2	8-9
7.3	7-8
9.9	9-10
	Playing Music  8.2 6.1 5.5 7.8 6.4 7.6 8.2 7.3




- But what if the data are continuous (decimals) and not discrete (integer-valued)?
- Not very informative!
- In this case, we place the data in "bins" to visualize the distribution
  - e.g., 4-5, 5-6, 6-7
- The number of bins is chosen by the data scientist

Person	Years Playing Music	Bins
1	8.2	8-9
2	6.1	6-7
3	5.5	5-6
4	7.8	7-8
5	6.4	6-7
6	7.6	7-8
7	8.2	8-9
8	7.3	7-8
9	9.9	9-10




- But what if the data are continuous (decimals) and not discrete (integer-valued)?
- Not very informative!
- In this case, we place the data in "bins" to visualize the distribution
  - e.g., 4-5, 5-6, 6-7
- The number of bins is chosen by the data scientist
- The best number of bins is subjective

Person	Years Playing Music	Bins
1	8.2	8-9
2	6.1	6-7
3	5.5	5-6
4	7.8	7-8
5	6.4	6-7
6	7.6	7-8
7	8.2	8-9
8	7.3	7-8
9	9.9	9-10



- But what if the data are continuous (decimals) and not discrete (integer-valued)?
- Not very informative!
- In this case, we place the data in "bins" to visualize the distribution
  - e.g., 4-5, 5-6, 6-7
- The number of bins is chosen by the data scientist
- The best number of bins is subjective
- You may need to try a few times before acquiring a useful histogram


Person	Years Playing Music	Bins
1	8.2	8-9
2	6.1	6-7
3	5.5	5-6
4	7.8	7-8
5	6.4	6-7
6	7.6	7-8
7	8.2	8-9
8	7.3	7-8
9	9.9	9-10




 Thus far, we have defined a data distribution and visualized it using a histogram


- Thus far, we have defined a data distribution and visualized it using a histogram
- Data distributions can take on many, many different shapes

- Thus far, we have defined a data distribution and visualized it using a histogram
- Data distributions can take on many, many different shapes






Histogram of




- Thus far, we have defined a data distribution and visualized it using a histogram
- Data distributions can take on many, many different shapes


• However, the most common is a bell-shaped curve called the normal distribution  $\frac{1}{1-(x-y)^2}$ 

 $f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{\frac{-(x-\mu)^2}{2\sigma^2}}$ 






Histogram of




# Distributions and Descriptive Statistics


- Descriptive statistics describe different aspects of the distribution
- Central tendency Where is the center of the distribution?




- Descriptive statistics describe different aspects of the distribution
- Central tendency Where is the center of the distribution?




- Descriptive statistics describe different aspects of the distribution
- Central tendency Where is the center of the distribution?
  - Mean = 5.99, Median = 5.98, Mode = ~5.75




- Descriptive statistics describe different aspects of the distribution
- Central tendency Where is the center of the distribution?
  - Mean = 5.99, Median = 5.98, Mode = ~5.75
- What can we say about years playing music among students in our sample?




- Descriptive statistics describe different aspects of the distribution
- Central tendency Where is the center of the distribution?
  - Mean = 5.99, Median = 5.98, Mode = ~5.75
- What can we say about years playing music among students in our sample?
- Note: For the theoretical normal distribution, the mean = median = mode




- Descriptive statistics describe different aspects of the distribution
- Dispersion What is the range of years of music played?




- Descriptive statistics describe different aspects of the distribution
- Dispersion What is the range of years of music played?
  - 2 to 10 years




- Descriptive statistics describe different aspects of the distribution
- Dispersion What is the range of years of music played?
  - 2 to 10 years
- What can we say about this student?




- Descriptive statistics describe different aspects of the distribution
- Dispersion What is the range of years of music played?
  - 2 to 10 years
- What can we say about this student?
- Where is the student with the least amount of experience playing music?




- Imagine we collect data on two different groups of students (shown on the right)
- Which sample (1 or 2) has a smaller standard deviation in years of playing music?
  - A) Sample 1
  - B) Sample 2






- Imagine we collect data on two different groups of students (shown on the right)
- Which sample (1 or 2) has a smaller standard deviation in years of playing music?
  - A) Sample 1
  - B) Sample 2




Histogram of



When the distribution is close to symmetric the

mean ≈ median





When the distribution is close to symmetric the

mean ≈ median

• Left skewed distributions

mean < median







When the distribution is close to symmetric the


mean ≈ median


Left skewed distributions


mean < median

Right skewed distributions

mean > median










Data distributions can take on many, many different shapes!



# R Code Covered in Practice Assignment