

Ripples in Reality: The Theoretical Origin and Observational Confirmation of Gravitational Waves

Armaan Sachdeva

January 6, 2026

Abstract

For nearly a century, gravitational waves existed primarily as a mathematical consequence of Albert Einstein's General Theory of Relativity, a theoretical "shimmer" in the geometric fabric of the universe. This dissertation explores the journey of these waves from their formal derivation to their eventual direct detection, which has fundamentally altered our capacity to observe the cosmos.

The study begins with the mathematical origin of gravitational waves, demonstrating how the linearization of the Einstein Field Equations in the weak-field limit reveals a wave equation for spacetime perturbations. This derivation highlights critical mathematical concepts, including the application of the Lorenz gauge and the use of the trace-reversed metric perturbation. We further analyze the unique physical properties of these waves, specifically their transverse-traceless nature and the requirement of a time-varying quadrupole moment for their generation—a distinct departure from the dipole radiation found in electromagnetism.

Finally, the dissertation examines the observational evidence that transformed gravitational waves from a geometric prediction into a tool for precision physics. We discuss the transition from the indirect confirmation provided by the Hulse-Taylor binary pulsar to the direct detection era initiated by LIGO. By matching observed "chirp" waveforms to numerical relativity templates, we demonstrate how gravitational waves have enabled the first rigorous tests of General Relativity in the strong-field regime. This work concludes that the detection of these waves represents a shift in the astronomical paradigm: we have moved from a "silent" universe observed through light to a dynamic one we can now "hear" through the ripples of spacetime itself.

Contents

1	Introduction	2
2	The Mathematical origins of Gravitational Waves	3
2.1	The Einstein Field Equations	3
2.2	Linearized Gravity	3
2.2.1	The Connection and Curvature	4
2.3	Gauge Invariance and the Wave Equation	4
2.3.1	Trace-reversed Perturbation	4
2.3.2	The Lorenz Gauge	4
2.3.3	The Final Wave Equation	4
3	Properties and Waveforms of Gravitational Waves	6
3.1	The Transverse-Traceless (TT) Gauge	6
3.2	Physical Effect on Matter	6
3.3	Generation: The Quadrupole Formula	7
3.4	The "Chirp" from Binary Systems	8
4	Observational Tests of General Relativity	9
4.1	Indirect Evidence: The Hulse-Taylor Binary	9
4.2	Direct Tests with LIGO/Virgo	9
4.2.1	The Speed of Gravity	12
5	Conclusion	13
	References	13

Chapter 1

Introduction

On the 14th of September 2015, a "chirp" signal that had travelled through the cosmos for over a billion years was detected by the LIGO interferometers on Earth. This signal was designated GW150914. Despite the signal lasting only a fraction of a second, this detection marked a turning point in physics [2]. This discovery confirmed the existence of a gravitational wave: a ripple in the fabric of space and time itself, caused by the cataclysmic merger of two black holes. This event was not just the confirmation of the theory of relativity; it was the birth of a whole new field of astronomy, one that delves into the most violent and energetic events in the universe.

This prediction was first formulated by Einstein in his General Theory of Relativity (GR) in 1915 [11, 12]. GR changed the way we understand gravity, replacing the Newtonian concept of gravitation as a force with a description of geometric deformation in spacetime. Einstein realised that if spacetime is dynamic, then it should also be able to ripple. Just as an accelerating electric charge radiates electromagnetic waves, an accelerating mass system with a changing quadrupole moment must radiate gravitational waves—disturbances in spacetime curvature propagating at the speed of light. Thus, in 1916, he posited this exact claim, and almost 100 years later, it was finally confirmed.

This dissertation explores the theoretical underpinnings and observational implications of gravitational waves. Chapter 2 derives the existence of these waves directly from the Einstein Field Equations using linearized gravity. Chapter 3 examines the generation of these waves via the quadrupole formula. Chapter 4 analyzes the characteristic "chirp" waveform produced by compact binary coalescences. Finally, Chapter 5 discusses how recent detections have enabled rigorous new tests of General Relativity in the strong-field regime.

Chapter 2

The Mathematical origins of Gravitational Waves

2.1 The Einstein Field Equations

General Relativity is made up of a set of 10 coupled, non-linear partial differential equations known as Einstein's field equations. These can be written succinctly using the summation convention. They relate the geometry of spacetime to the matter which distorts it:

$$G_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu} \quad (2.1)$$

Where $G_{\mu\nu}$ is the Einstein Tensor, which describes the local curvature of spacetime. The Einstein tensor is defined by the contraction of the Riemann curvature tensor:

$$G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2} R g_{\mu\nu} \quad (2.2)$$

where $R_{\mu\nu} = R^\lambda_{\mu\lambda\nu}$ is the Ricci tensor (the trace of the Riemann tensor), and $R = g^{\mu\nu} R_{\mu\nu}$ is the Ricci scalar. This specific geometric combination is required because its covariant divergence is identically zero ($\nabla^\mu G_{\mu\nu} = 0$), ensuring compatibility with the conservation of energy-momentum ($\nabla^\mu T_{\mu\nu} = 0$). $T_{\mu\nu}$ is the stress energy tensor, which describes the distribution of matter and energy. The coupling constant $\kappa = \frac{8\pi G}{c^4}$ is approximately $2 \times 10^{-43} \text{ N}^{-1}$ in SI units. This constant implies that these equations are incredibly "stiff". This means one needs an enormous amount of energy-momentum $T_{\mu\nu}$ in order to see any noticeable curvature in local spacetime $G_{\mu\nu}$.

To put this stiffness into perspective, imagine a trampoline. Were the equations not stiff, placing any weight would cause the trampoline to deform significantly. However, because spacetime is stiff, one must imagine replacing the material of the trampoline with solid steel. Suddenly, one needs considerably more mass to make any sort of meaningful deformation [10].

2.2 Linearized Gravity

The coupled set of equations are almost impossible to solve by normal means. Therefore we have to employ an approximation, given by perturbation theory. We assume that we are far away from any strong gravitational sources- such as black holes- which allows us to treat spacetime as a flat metric using the Minkowski metric ($\eta_{\mu\nu}$) with some small perturbation ($h_{\mu\nu}$):

$$g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}, \quad \text{where } |h_{\mu\nu}| \ll 1 \quad (2.3)$$

With this new approximation of "Linearized gravity", we can neglect terms of order $O(h^2)$ and higher as they would have no meaningful contribution. This allows us to turn the non-linear physics of GR into a linear field theory on a flat spacetime background.

2.2.1 The Connection and Curvature

First, we compute the linearized Christoffel symbols. In the standard metric formulation, $\Gamma_{\mu\nu}^\rho = \frac{1}{2}g^{\rho\lambda}(\partial_\mu g_{\nu\lambda} + \partial_\nu g_{\mu\lambda} - \partial_\lambda g_{\mu\nu})$. Substituting our perturbed metric and keeping only first-order terms, this simplifies to:

$$\Gamma_{\mu\nu}^\rho \approx \frac{1}{2}\eta^{\rho\lambda}(\partial_\mu h_{\nu\lambda} + \partial_\nu h_{\mu\lambda} - \partial_\lambda h_{\mu\nu}) \quad (2.4)$$

From this, we construct the linearized Riemann curvature tensor. While the full expression is complex, the linearized Ricci tensor ($R_{\mu\nu} = R_{\mu\lambda\nu}^\lambda$) reduces to a sum of four second-derivative terms:

$$R_{\mu\nu} \approx \frac{1}{2}(\partial_\rho \partial_\mu h_\nu^\rho + \partial_\rho \partial_\nu h_\mu^\rho - \partial_\mu \partial_\nu h - \square h_{\mu\nu}) \quad (2.5)$$

where $h = h_\mu^\mu$ is the trace of the perturbation, and $\square = \partial^\lambda \partial_\lambda = -c^{-2}\partial_t^2 + \nabla^2$ is the d'Alembertian wave operator.

2.3 Gauge Invariance and the Wave Equation

The expression for $R_{\mu\nu}$ as seen above is not an elegant expression. However, GR contains a gauge symmetry which is analogous to that found in Electromagnetism. Just as we can adjust the electromagnetic potential $A_\mu \rightarrow A_\mu + \partial_\mu \Lambda$ without changing the physical fields, we can adjust the coordinate system to simplify the form of $h_{\mu\nu}$ without changing the physical spacetime curvature.

2.3.1 Trace-reversed Perturbation

To clean up the equations, it is standard practice to introduce the trace-reversed perturbation, $\bar{h}_{\mu\nu}$:

$$\bar{h}_{\mu\nu} = h_{\mu\nu} - \frac{1}{2}\eta_{\mu\nu}h \quad (2.6)$$

The name "trace-reversed" comes from the fact that taking the trace of this new variable flips the sign of the original trace: $\bar{h} = -h$.

2.3.2 The Lorenz Gauge

We now choose a specific coordinate system known as the Lorenz gauge (or harmonic gauge). This imposes the condition:

$$\partial^\mu \bar{h}_{\mu\nu} = 0 \quad (2.7)$$

This is mathematically identical to the Lorenz gauge condition in Electromagnetism ($\partial^\mu A_\mu = 0$). By applying this condition, the first three terms in the linearized Ricci tensor cancel out or combine, leading to a dramatic simplification of the Einstein Tensor.

2.3.3 The Final Wave Equation

Substituting the trace-reversed variable and applying the Lorenz gauge condition to the linearized Einstein Field Equations, we arrive at the central result of this chapter:

$$\square \bar{h}_{\mu\nu} = -\frac{16\pi G}{c^4}T_{\mu\nu} \quad (2.8)$$

In a vacuum, where $T_{\mu\nu} = 0$, this becomes the homogeneous wave equation:

$$\left(-\frac{1}{c^2}\frac{\partial^2}{\partial t^2} + \nabla^2\right)\bar{h}_{\mu\nu} = 0 \quad (2.9)$$

Equation 2.9 is the "smoking gun" of the theory. It tells us that in empty space, perturbations to the spacetime metric ($\bar{h}_{\mu\nu}$) do not just sit still or decay instantly; they propagate as waves. Furthermore, the factor of $1/c^2$ in the d'Alembertian indicates that these waves travel exactly at the speed of light, c . This mathematical derivation confirms Einstein's 1916 prediction: gravity is not an instantaneous force, but a field that communicates changes via waves propagating through the geometry of the universe [16][9].

Chapter 3

Properties and Waveforms of Gravitational Waves

From the previous chapter we have established that the Einstein Field equations allow for wave solutions. Now we must put them into physical perspective. What do these waves look like? What kind of motion is able to generate them?

3.1 The Transverse-Traceless (TT) Gauge

In Section 2.3, we used the Lorenz gauge to simplify the wave equation. However, this simplification did not exhaust all of the gauge freedoms. The perturbation metric $h_{\mu\nu}$ is a symmetric 4×4 tensor; this tensor contains 10 independent components, which is far too many for a physical field describing a massless spin-2 particle (which should only have 2 degrees of freedom, analogous to the two polarizations of a photon).

In order to isolate the physical degrees of freedom we need to work in a new gauge: the Transverse-Traceless (TT) gauge. In this frame, using a wave propagating in the z -direction as an example, we transpose 4 additional constraints such that all the time components and the trace become 0 ($h_{0\mu} = 0$ and $h^\mu_\mu = 0$). The resulting wave tensor takes a sparse form:

$$h_{\mu\nu}^{TT} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & h_+ & h_\times & 0 \\ 0 & h_\times & -h_+ & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \cos(\omega(t - z/c)) \quad (3.1)$$

Here, the indices represent (t, x, y, z) . The solution reveals two independent polarization states: the "plus" mode (h_+) and the "cross" mode ($h_\times$) [17].

3.2 Physical Effect on Matter

Gravitational waves are described as "tidal" waves due to the fact that they stretch and squeeze spacetime in directions perpendicular to the direction they propagate. This can be visualized by considering a ring of free falling test particles in the x - y plane (Figure 3.1).

- The Plus Polarization (h_+): As the wave passes, it stretches the ring along the x -axis while simultaneously squeezing it along the y -axis, transforming the circle into an ellipse. Half a cycle later, the effect reverses.
- The Cross Polarization ($h_\times$): This mode acts identically but rotated by 45° , stretching along the line $y = x$ and squeezing along $y = -x$.




Figure 3.1: Visualization of the effect of gravitational waves on a ring of test particles. The dashed grey line represents the original circular distribution. The solid green ellipse shows the instantaneous deformation of space as the wave passes. h_+ stretches the ring along the axes, while $h_\times$ stretches it along the diagonals. Diagrams were created by me via tkiz package in LaTeX, modified from[13]

This is a distinct sign of a "quadrupolar" deformation pattern. An expansion in one direction is matched by a contraction in the orthogonal direction. This is the signature pattern which interferometers like LIGO look for. It is the reason that LIGO's arms are perpendicular; as one arm lengthens, the other shortens, doubling the phase difference of the laser light.

3.3 Generation: The Quadrupole Formula

Since the gravitational field itself carries energy, we can define an effective stress-energy tensor for the waves, known as the Isaacson tensor $t_{\mu\nu}$. In the short-wavelength approximation, the energy flux is proportional to the square of the time-derivative of the perturbation:

$$t_{\mu\nu} = \frac{c^4}{32\pi G} \langle \dot{h}_{ij} \dot{h}^{ij} \rangle \quad (3.2)$$

This non-linear property is what allows gravitational waves to transport energy and angular momentum away from the binary system, driving the orbital decay observed in the Hulse-Taylor pulsar.[16] [17]

In electromagnetism, the dominant form of radiation is dipole radiation, generated by a time-varying charge dipole ($\vec{d}$). Why is gravity different? We can analyze this via a multipole expansion of the source term $T_{\mu\nu}$:

1. Monopole Moment (Mass): The first derivative of mass is zero due to conservation of mass-energy ($\dot{M} = 0$). A spherically symmetric pulsation radiates nothing.
2. Dipole Moment: The first derivative of the mass dipole is momentum (P), and the second derivative is the rate of change of momentum ($\dot{P}$). Due to the conservation of linear momentum, an isolated system has $\dot{d} = 0$. There is no gravitational dipole radiation.

Thus, the lowest-order radiation comes from the quadrupole moment (Q_{ij}). Einstein derived the famous Quadrupole Formula, which states that the amplitude of the wave h scales with the second time derivative of the quadrupole moment $\ddot{Q}$:

$$h_{ij} \approx \frac{2G}{c^4 r} \ddot{Q}_{ij}^{TT} \quad (3.3)$$

This tells us that to generate gravitational waves, a system must not only accelerate but must do so in a non-spherically symmetric way. A spinning neutron star with a "mountain" on it will radiate; a perfectly smooth spinning sphere will not [12].

3.4 The "Chirp" from Binary Systems

The most potent sources of gravitational waves are compact binary systems (neutron stars or black holes) spiraling into one another. As the system orbits, it has a massive, rapidly changing quadrupole moment, radiating energy away in the form of GWs.




Figure 3.2: A schematic representation of the characteristic "chirp" waveform from a binary black hole merger. Diagram was created by me via tkiz package in LaTeX, a simplified version from the original ligo paper [2]

As energy is lost, the orbital radius r shrinks[18]. Kepler's laws dictate that as the radius shrinks, the orbital velocity and frequency must increase. This creates a runaway process. The resulting waveform is a "chirp": a sinusoidal wave that sweeps up in both amplitude and frequency effectively until the moment of merger. This waveform serves as a fingerprint for identifying the masses and spins of the progenitors [2, 23].

Chapter 4

Observational Tests of General Relativity

This chapter connects the theory to observation, detailing how the detections predicted in Chapter 3 have allowed for rigorous tests of Einstein’s theory.

4.1 Indirect Evidence: The Hulse-Taylor Binary

Experimental efforts to detect gravitational waves began in the 1960s with Joseph Weber’s resonant bar detectors. While Weber claimed detection in 1969 [22], these results could not be replicated, leading the field to shift toward interferometry. Before direct detection was possible, compelling indirect evidence for gravitational waves was found in the Hulse-Taylor binary pulsar (PSR B1913+16), discovered in 1974. This system consists of two neutron stars orbiting each other with a period of approximately 7.75 hours.

Over a span of thirty years, precise timing of the pulsar’s radio pulses revealed that the orbital period was decreasing. This orbital decay indicated that the system was losing energy. Weisberg and Taylor demonstrated that this energy loss matched the predictions of the General Relativistic quadrupole formula to within 0.2% [23]. This precise agreement was the first confirmation that gravitational radiation physically carries energy away from a system, earning Hulse and Taylor the 1993 Nobel Prize in Physics.

4.2 Direct Tests with LIGO/Virgo

The detection of strains on the order of $h \sim 10^{-21}$ requires overcoming significant noise sources that mask the signal. At low frequencies (< 10 Hz), the sensitivity is limited by seismic noise from ground vibrations, which LIGO mitigates using sophisticated multi-stage pendulums. At intermediate frequencies, thermal noise in the mirror coatings dominates. At high frequencies (> 200 Hz), the fundamental limit is photon shot noise—quantum fluctuations in the number of photons arriving at the photodetector. The ‘whitening’ process applied to the data in Figure 4.1 is crucial specifically to suppress these dominant noise lines and reveal the underlying astrophysical signal.[1]

The direct detection of GW150914 allowed for tests of General Relativity in the “strong-field” regime—where gravitational fields are intense and velocities approach the speed of light, a regime inaccessible to solar system experiments [1].

To validate the quadrupole formalism derived in Chapter 3, We compared the output of a Newtonian numerical integration to the open data for GW150914. As shown in Figure 4.2, the frequency evolution of the Newtonian model matches the observed signal closely during




Figure 4.1: Leading-order Newtonian simulation of a binary black hole inspiral. The simulation integrates the energy loss predicted by the Quadrupole Formula. The bottom panel highlights the non-linear frequency evolution, which scales as $\dot{f} \propto f^{11/3}$, creating the runaway feedback loop characteristic of the inspiral phase. Note that the Newtonian approximation diverges at $t = 0$ as it does not account for the ringdown phase or higher-order Post-Newtonian terms.[20]




Figure 4.2: Comparison of the numerical Newtonian simulation (orange dashed) with the whitened strain data of GW150914 (blue solid). The simulation amplitude is rescaled for visual comparison. The Newtonian model accurately predicts the frequency evolution (chirp rate) during the inspiral phase ($t < -0.05\text{s}$). However, a phase drift becomes visible as the system approaches merger, illustrating the accumulation of relativistic effects not captured by the quadrupole approximation.[20]

the inspiral phase. The period of oscillation shortens at the same rate in both the theory and observation, validating the quadrupole energy loss formula derived in Eq 3.3.

However, two deviations are apparent. First, there is a noticeable phase drift between the curves as they approach $t = 0$; this is expected, as the Newtonian approximation fails to account for higher-order Post-Newtonian (PN) terms that affect the orbital phase[8]. Second, at $t = 0$, the real signal undergoes a smooth merger and ringdown[19][14], whereas the singular Newtonian model effectively cuts off. This comparison neatly isolates the domain of validity for linearized gravity.

We can further validate the source parameters by estimating the "Chirp Mass" ($\mathcal{M}$) directly from the signal's frequency evolution shown in Figure 4.1. Rewriting Eq 3.3 in terms of frequency f and its time derivative $\dot{f}$, we obtain:

$$\mathcal{M} = \frac{c^3}{G} \left(\frac{5}{96} \pi^{-8/3} f^{-11/3} \dot{f} \right)^{3/5} \quad (4.1)$$

From the simulation data in Figure 4.1, at time $t_1 = -0.10\text{s}$, the frequency is approximately $f_1 \approx 45\text{ Hz}$. Over the interval $\Delta t = 0.05\text{s}$, the frequency rises to $f_2 \approx 58\text{ Hz}$ at $t_2 = -0.05\text{s}$. This yields an approximate frequency derivative of $\dot{f} \approx 260\text{ Hz/s}$.

Substituting these values into the equation (using a central frequency $f \approx 51.5\text{ Hz}$) yields an estimated chirp mass of:

$$\mathcal{M} \approx 27.3 M_{\odot} \quad (4.2)$$

This estimate is consistent with the LIGO collaboration's reported source-frame chirp mass of roughly $28 M_{\odot}$ [2][3], confirming that the signal indeed originates from a stellar-mass black hole binary and demonstrating the power of the quadrupole formalism to extract physical parameters from the waveform.




Figure 4.3: Multi-messenger observation of the binary neutron star merger GW170817. Top: Gamma-ray count rate detected by the Fermi satellite (GRB 170817A). Bottom: Time-frequency map of the gravitational wave strain detected by LIGO. Note the ~ 1.7 second delay between the merger time ($t = 0$) and the arrival of the gamma-ray burst, constraining the speed of gravity to match the speed of light. [15][4]

The derivation of a $\approx 30M_{\odot}$ component mass was astrophysically significant. Prior to GW150914, the most massive known stellar-mass black holes were X-ray binaries below $20M_{\odot}$. The existence of such heavy black holes implies they formed in a low-metallicity environment where stellar winds are weak, allowing the progenitor stars to retain most of their mass before collapsing. This detection thus provided the first observational evidence for the existence of 'heavy' stellar-mass black holes.[2]

4.2.1 The Speed of Gravity

One of the most fundamental predictions of GR is that gravitational waves propagate at the speed of light, c . This was dramatically tested with the detection of GW170817, a binary neutron star merger. This event produced both a gravitational wave signal and a short gamma-ray burst (GRB 170817A).

The gamma rays arrived at Earth just 1.7 seconds after the gravitational wave signal ended, despite having traveled a distance of 130 million light-years. This near-simultaneous arrival constrained the difference between the speed of gravity (v_g) and the speed of light (c) to be extremely small [4]:

$$-3 \times 10^{-15} \leq \frac{v_g - c}{c} \leq 7 \times 10^{-16} \quad (4.3)$$

This result effectively rules out many alternative theories of gravity that predict a mass for the graviton or different propagation speeds.[5]

Chapter 5

Conclusion

The direct detection of gravitational waves represents the culmination of a century of theoretical and experimental endeavour. From the first derivation of the linearized wave equation in 1916 to the landmark observations of the LIGO-Virgo-KAGRA network, the journey has been one of moving from mathematical abstraction to physical reality.

This dissertation has shown how these "ripples in reality" are a natural consequence of a dynamic, geometric description of spacetime. By linearizing the Einstein Field Equations, we recovered a wave operator that predicts the propagation of metric perturbations at the speed of light. We have explored how the specific, quadrupolar nature of these waves, which arises from the conservation laws of mass and momentum, serves as a distinct fingerprint of General Relativity, separate from electromagnetic radiation.

The observational confirmation provided by binary black hole and neutron star mergers has done more than just validate Einstein's prediction; it has subjected General Relativity to its most rigorous tests in the previously inaccessible strong-field regime. The near-perfect matching of observed waveforms to Numerical Relativity templates, alongside the precise constraints on the speed of gravity, has left General Relativity as the undisputed champion of gravitational theories.

Looking forward, we are entering an era where gravitational waves are no longer just an object of study, but a tool for exploration. The transition from electromagnetic to multi-messenger astronomy is akin to the transition from silent films to "talkies" in cinema history; we are no longer just seeing the universe, we are truly "hearing" the collision of its most compact and energetic objects [21]. With future detectors such as the space-based LISA mission[7], we will expand our sensitivity to lower frequencies, allowing us to listen to the background hum of the early universe and the mergers of supermassive black holes[6]. The ripples Einstein predicted a century ago have finally reached us, and in doing so, they have changed the way we perceive the cosmos forever.

Bibliography

- [1] J. Aasi et al. Advanced ligo. *Classical and Quantum Gravity*, 32(7):074001, 2015.
- [2] B. P. Abbott et al. Observation of gravitational waves from a binary black hole merger. *Phys. Rev. Lett.*, 116:061102, Feb 2016.
- [3] B. P. Abbott et al. Properties of the binary black hole merger gw150914. *Physical Review Letters*, 116(24):241102, 2016.
- [4] B. P. Abbott et al. Gravitational waves and gamma-rays from a binary neutron star merger: Gw170817 and grb 170817a. *Astrophys. J. Lett.*, 848(2):L12, 2017.
- [5] B. P. Abbott et al. Gw170817: Observation of gravitational waves from a binary neutron star inspiral. *Physical Review Letters*, 119(16):161101, 2017.
- [6] Gabriella Agazie et al. The nanograv 15 yr data set: Evidence for a gravitational-wave background. *The Astrophysical Journal Letters*, 951(1):L8, 2023.
- [7] Pau Amaro-Seoane et al. Laser interferometer space antenna. *arXiv preprint arXiv:1702.00786*, 2017.
- [8] Luc Blanchet. Gravitational radiation from post-newtonian sources and inspirals. *Living Reviews in Relativity*, 17(1):2, 2014.
- [9] Hermann Bondi, MGJ van der Burg, and AWK Metzner. Gravitational waves in general relativity. vii. waves from axi-symmetric isolated systems. *Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences*, 269(1336):21–52, 1962.
- [10] Sean M Carroll. *Spacetime and Geometry: An Introduction to General Relativity*. Addison Wesley, San Francisco, 2004.
- [11] A. Einstein. Näherungsweise integration der feldgleichungen der gravitation. *Sitzungsberichte der Königlich Preußischen Akademie der Wissenschaften (Berlin)*, pages 688–696, 1916.
- [12] A. Einstein. Über gravitationswellen. *Sitzungsberichte der Königlich Preußischen Akademie der Wissenschaften (Berlin)*, pages 154–167, 1918.
- [13] Maximiliano Isi. Parametrizing gravitational-wave polarizations. *Zenodo (CERN European Organization for Nuclear Research)*, 08 2022.
- [14] Roy P. Kerr. Gravitational field of a spinning mass as an example of algebraically special metrics. *Physical Review Letters*, 11(5):237, 1963.
- [15] Briley Lewis. Meet the aas keynote speakers: Dr. brian metzger, 12 2019.
- [16] Michele Maggiore. *Gravitational Waves: Volume 1: Theory and Experiments*. Oxford University Press, 2008.

- [17] Charles W. Misner, Kip S. Thorne, and John Archibald Wheeler. Gravitation. W. H. Freeman, San Francisco, 1973.
- [18] Philip C Peters and J Mathews. Gravitational radiation from point masses in a keplerian orbit. *Physical Review*, 131(1):435, 1963.
- [19] Frans Pretorius. Evolution of binary black hole spacetimes. *Physical Review Letters*, 95(12):121101, 2005.
- [20] Armaan Sachdeva. Newtonian binary black hole simulation code. <https://github.com/BBCnewslondon/Chirp>, 2026.
- [21] B. S. Sathyaprakash and Bernard F. Schutz. Physics, astrophysics and cosmology with gravitational waves. *Living Reviews in Relativity*, 12(1):2, 2009.
- [22] Joseph Weber. Evidence for discovery of gravitational radiation. *Physical Review Letters*, 22(24):1320, 1969.
- [23] Joel M. Weisberg and Joseph H. Taylor. Relativistic binary pulsar b1913+16: Thirty years of observations and analysis. *ASP Conf. Ser.*, 328:25, 2005. arXiv:astro-ph/0407149.