



[Unit 2 Nonlinear Classification,](#)
[Linear regression, Collaborative](#)
[Course](#) > [Filtering \(2 weeks\)](#)
8. Regularization

> [Lecture 5. Linear Regression](#) >

Audit Access Expires May 11, 2020

You lose all access to this course, including your progress, on May 11, 2020. Upgrade by Mar 25, 2020 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

8. Regularization

Ridge Regression




Video[Download video file](#)**Transcripts**[Download SubRip \(.srt\) file](#)[Download Text \(.txt\) file](#)

Regularization: extreme case 1

0/1 point (graded)

As in the video above, define the loss function

$$J_{n,\lambda}(\theta, \theta_0) = \frac{1}{n} \sum_{t=1}^n \frac{(y^{(t)} - \theta \cdot x^{(t)} - \theta_0)^2}{2} + \frac{\lambda}{2} \|\theta\|^2$$

where λ is the regularization factor.

In the figure above, the blue dots are the training examples. If we increase λ to ∞ , where does $f(x) = \theta \cdot x + \theta_0$ converge to?

☒ line 1 ✓

☐ line 2

☐ line 3


Solution:

If we increase λ to ∞ , minimizing J is equivalent to minimizing $\|\theta\|$. Thus θ will have to be a zero vector. Thus $f(x) = \theta \cdot x + \theta_0$ becomes $f(x) = \theta_0$, a horizontal line. Thus f converges to line 1.

You have used 2 of 2 attempts

i Answers are displayed within the problem


Regularization: Extreme case 2

1/1 point (graded)

As in the problem above,

$$J_{n,\lambda}(\theta, \theta_0) = \frac{1}{n} \sum_{t=1}^n \frac{(y^{(t)} - \theta \cdot x^{(t)} - \theta_0)^2}{2} + \frac{\lambda}{2} \|\theta\|^2$$

where λ is the regularization factor.



In the figure above, the blue dots are the training examples. If we decrease λ to 0, where does $f(x) = \theta \cdot x + \theta_0$ converge to?

☐ line 1

☒ line 2

☐ line 3



Solution:

If we decrease λ to zero, minimizing J is equivalent to minimizing

$\frac{1}{n} \sum_{t=1}^n \frac{(y^{(t)} - \theta \cdot x^{(t)} - \theta_0)^2}{2}$, which is the "fit." Thus f converges to line 2.

Submit

You have used 1 of 2 attempts

 Answers are displayed within the problem

Discussion





[Hide Discussion](#)

Topic: Unit 2 Nonlinear Classification, Linear regression, Collaborative Filtering (2 weeks):Lecture 5. Linear Regression / 8. Regularization

[Add a Post](#)

Show all posts

by recent activity

- | | | |
|---|---|---|
|  | what is the $\eta * \lambda * \theta$ do in the update | 3 |
|  | Why does the derivative of square norm of theta is $2 * \theta$
When calculating the gradient of the regularization term, the gradient/derivative of λ... | 3 |
|  | [Staff] Typo mistake with derivation on the board
I think the professor forgot to include the summation sign in the derivation (when she took t... | 5 |
|  | why θ_0 is not regularized?
hi what's the reason that θ_0 is not included in the regularization component of the loss f... | 3 |

[Learn About Verified Certificates](#)

© All Rights Reserved