



[Unit 2 Nonlinear Classification,](#)
[Linear regression, Collaborative](#)

[Course](#) > [Filtering \(2 weeks\)](#)

> [Lecture 5. Linear Regression](#) >

9. Closing Comment

Audit Access Expires May 11, 2020

You lose all access to this course, including your progress, on May 11, 2020.

Upgrade by Mar 25, 2020 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

9. Closing Comment

Closing Comment



Video[Download video file](#)**Transcripts**[Download SubRip \(.srt\) file](#)[Download Text \(.txt\) file](#)**(Optional) Equivalence of regularization to a Gaussian Prior on Weights****(Optional) Equivalence of regularization to a Gaussian Prior on Weights**

The regularized linear regression can be interpreted from a probabilistic point of view. Suppose we are fitting a linear regression model with n data points $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$. Let's assume the ground truth is that y is linearly related to x but we also observed some noise ϵ for y :

$$y_t = \theta \cdot x_t + \epsilon$$

where $\epsilon \sim \mathcal{N}(0, \sigma^2)$.

Then the likelihood of our observed data is

$$\prod_{t=1}^n \mathcal{N}(y_t | \theta x_t, \sigma^2).$$

Now, if we impose a Gaussian prior $\mathcal{N}(\theta | 0, \lambda^{-1})$, the likelihood will change to

$$\prod_{t=1}^n \mathcal{N}(y_t | \theta x_t, \sigma^2) \mathcal{N}(\theta | 0, \lambda^{-1}).$$

Take the logarithm of the likelihood, we will end up with

$$\sum_{t=1}^n -\frac{1}{2\sigma^2}(y_t - \theta x_t)^2 - \frac{1}{2}\lambda \|\theta\|^2 + \text{constant}.$$

Try to derive this result by yourself. Can you conclude that maximizing this loglikelihood equivalent to minimizing the regularized loss in the linear regression? What does larger λ mean in this probabilistic interpretation? (Think of the error decomposition we discussed.)







[Hide](#)

Discussion

[Hide Discussion](#)

Topic: Unit 2 Nonlinear Classification, Linear regression, Collaborative Filtering (2 weeks):Lecture 5. Linear Regression / 9. Closing Comment

[Add a Post](#)

| Show all posts | by recent activity |
|--|--------------------|
|  Help clarify the notation in the optional problem | 2 |
|  Great Lecture! Amazing. A difficult mathematical equation expressed in simple words. Great teacher! | 13 |
|  [STAFF] What does larger λ mean in this probabilistic interpretation? (Think of the error decomposition we discussed.) How it is related to the error decomposition is unclear to me | 2 |
|  Question regarding Gaussian weights (Optional Section) | 8 |
|  Why is minimizing the norm of theta increasing the error? Minimizing the norm of theta increases the error? or perhaps did I get it wrong from the lect... | 2 |
|  Question about Gaussian prior I want to know whether the first initialize of theta to be 0 caused the prior $N(0, 1/\lambda)$ or Not. If it... | 2 |



The probabilistic or Bayesian interpretation is very interesting

1

That explains why we choose the mean square loss and quadratic regularization, which earlie...

Learn About Verified Certificates

© All Rights Reserved