

Percolation Phase Transitions on Dynamically Grown Graphs

Braden Hoagland
Advised by Rick Durrett

July 8, 2021

Background

Dynamically Grown Graphs
Percolation

Basic Results

Continuity of the Phase Transition
Scaling Behavior

2-Choice Rules

Examples
Uniform Scaling

Future Directions

Background

Dynamically grown graphs and percolation

Dynamically Grown Graphs

Start with a graph with n vertices and no edges.

Every $t = 1/n$ units of time, add edges to the graph by sampling m vertices i.i.d. and following some fixed rule.

Dynamically Grown Graphs

Start with a graph with n vertices and no edges.

Every $t = 1/n$ units of time, add edges to the graph by sampling m vertices i.i.d. and following some fixed rule.

Let $n \rightarrow \infty$.

Percolation

A *giant component* is a cluster comprising a finite fraction εn of the graph.

Percolation is the emergence of a giant component.

Percolation can have lots of different qualitative behaviors.

Explosive Percolation

For simple rules, the giant component might emerge in a predictable, linear manner.

If a rule prioritizes adding together smaller clusters, the giant component's emergence is delayed and happens very quickly (seemingly discontinuous). This is called *explosive percolation*.

Basic Results

Continuity of the phase transition and scaling behavior

Continuity of the Phase Transition

ℓ -vertex rule: choose ℓ vertices i.i.d., and you're only required to add an edge if all ℓ of them are in distinct clusters.

Riordan, R., and Warnke, L. (2012): all ℓ -vertex rules have a continuous phase transition. We just can't see it in simulation because the coefficients are insanely small.

Continuity of the Phase Transition

ℓ -vertex rule: choose ℓ vertices i.i.d., and you're only required to add an edge if all ℓ of them are in distinct clusters.

Riordan, R., and Warnke, L. (2012): all ℓ -vertex rules have a continuous phase transition. We just can't see it in simulation because the coefficients are insanely small.

Their proof is by contradiction, so it doesn't give us much quantitative information about the clusters' behavior.

Scaling Behavior

For rules with continuous phase transitions, we see *scaling behavior*.

Let $\delta = t - t_c$ and let $P(s, t)$ be the probability that a randomly chosen vertex has cluster size s at time t . Then near t_c , there are constants τ and σ such that

$$P(s) = s^{1-\tau} f(s\delta^{1/\sigma}).$$

Scaling behavior lets us express several important quantities as powers of δ .

Scaling Behavior

For rules with continuous phase transitions, we see *scaling behavior*.

Let $\delta = t - t_c$ and let $P(s, t)$ be the probability that a randomly chosen vertex has cluster size s at time t . Then near t_c , there are constants τ and σ such that

$$P(s) = s^{1-\tau} f(s\delta^{1/\sigma}).$$

Scaling behavior lets us express several important quantities as powers of δ .

From now on, we assume scaling behavior.

Scaling Behavior

Let S be the size of the giant component, and let

$$\chi_k(t) = \sum_s s^k P(s, t).$$

Then

$$S \approx \delta^\beta, \quad \chi_1(t) \approx \delta^{-\gamma}, \quad \frac{\chi_k(t)}{\chi_{k-1}(t)} \approx \delta^{-\Delta}$$

These unknowns are called *critical exponents*.

Scaling Relations

Goal: determine all critical exponents in terms of one unknown.

Why is this useful?

Scaling Relations

Goal: determine all critical exponents in terms of one unknown.

Why is this useful?

What kinds of rules can we do this for?

2-Choice Rules

Generalizing rules with useful properties

2-Choice Rules

Pick two groups of vertices i.i.d.

Select one vertex from each and add an edge between them.

$\phi_i(s) = \mathbb{P}(\text{vertex chosen from group } i \text{ has cluster size } s).$

Erdős Rényi

Pick two random vertices and add an edge between them.

Erdős Rényi

Pick two random vertices and add an edge between them.

Percolation occurs after $t_c = 1/2$.

$\beta = 1$, so S grows linearly near t_c .

da Costa

Introduced by da Costa to disprove Achlioptas' discontinuity conjecture.

Pick two groups of vertices, both of size m . Pick the vertex with the smallest cluster size from the two groups and add an edge between them.

Same as Erdős Rényi when $m = 1$. As $m \rightarrow \infty$,

$$\beta \rightarrow 0, \quad t_c \rightarrow 1.$$

Uniform Scaling

For any 2-choice rule, the quantity $\partial_t S$ has a simple form that can be explicitly calculated.

Near t_c , $\partial_t S$ will look like

$$\delta^a + \delta^b + \delta^c + \dots$$

We say that a rule *scales uniformly* if $a = b = c = \dots$

Uniform Scaling

For any 2-choice rule, the quantity $\partial_t S$ has a simple form that can be explicitly calculated.

Near t_c , $\partial_t S$ will look like

$$\delta^a + \delta^b + \delta^c + \dots$$

We say that a rule *scales uniformly* if $a = b = c = \dots$

All 2-choice rules **almost** scale uniformly.

Uniform Scaling

Theorem

For any 2-choice rule, there will be two dominating terms of $\partial_t S$ with the same order. If some extra technical conditions hold, then the rule scales uniformly.

Uniform Scaling

Consequences:

- ▶ For all 2-choice rules, we can solve for all critical exponents, as well as the growth rate of the average cluster size, in terms of β .
- ▶ For a large family of 2-choice rules, we can do this algorithmically.

Future Directions

Future Directions

- ▶ Interaction between the groups?
- ▶ When does scaling behavior actually occur?
- ▶ What about n -choice rules?