

MATH 531 HOMEWORK 6

BRADEN HOAGLAND

Scalar Multiplication. *Prove scalar multiplication is a continuous function from $\mathbb{R} \times \mathcal{V} \rightarrow \mathcal{V}$, for \mathcal{V} a normed vector space and $\mathbb{R} \times \mathcal{V}$ equipped with the product metric space structure.*

Given open $U \subset \mathcal{V}$, we must show that $\cdot^{-1}(U)$ is open. Let $(k, v) \in \cdot^{-1}(U)$, i.e. $kv \in U$, then there is an $\varepsilon > 0$ such that $D(kv, \varepsilon) \subset U$. If we find δ such that $\cdot(D((k, v), \delta)) \subset D(kv, \varepsilon)$, then we are done. Since $\mathbb{R} \times \mathcal{V}$ is equipped with the product metric, if $(k', v') \in D((k, v), \delta)$, then

$$d_1((k, v), (k', v')) = |k - k'| + \|v - v'\| < \delta.$$

Since both of these terms are nonnegative, this implies $|k - k'| < \delta$ and $\|v - v'\| < \delta$. Then by the triangle inequality,

$$\begin{aligned} d_2(kv, k'v') &= \|kv - k'v'\| \\ &= \|kv - k'v + k'v - k'v'\| \\ &\leq \|v\||k - k'| + |k'|\|v - v'\|. \end{aligned}$$

Since $|k'| = |k' - k + k| \leq \delta + |k|$, this becomes

$$d_2(kv, k'v') < \|v\|\delta + (\delta + |k|)\delta.$$

Surely we can let $\delta \leq 1$, since δ need not be “optimal” in any sense. Then we have

$$d_2(kv, k'v') < \|v\| + |k| + \delta.$$

So if we take $\delta = \min\{1, \varepsilon - \|v\| - |k|\}$, then $\cdot(D((k, v), \delta)) \subset D(kv, \varepsilon)$, which implies that $\cdot^{-1}(U)$ is open. Thus scalar multiplication is a continuous function on a normed vector space.

Page 182, 4.1.3. *Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ be continuous. Show $A = \{(x, y) \in \mathbb{R}^2 \mid 0 \leq f(x, y) \leq 1\}$ is closed.*

Since f is continuous, we know that for any closed set $F \subset \mathbb{R}$, $f^{-1}(F)$ is closed in \mathbb{R}^2 . Thus it suffices to find a closed set $B \subset \mathbb{R}$ such that $f^{-1}(B) = A$.

Let $B = \{z \in \mathbb{R} \mid 0 \leq z \leq 1\}$, then we claim that B is closed in \mathbb{R} and that $f^{-1}(B) = A$. Since $B^c = \{z < 0\} \cup \{z > 1\}$ is clearly open (it is the finite intersection of open sets), B is closed by definition. Also by definition, $f^{-1}(B) = \{(x, y) \mid f(x, y) \in B\} = \{(x, y) \mid 0 \leq f(x, y) \leq 1\} = A$.

Page 184, 4.2.4. Let $A, B \subset \mathbb{R}$, and suppose $A \times B \subset \mathbb{R}^2$ is connected.

- (1) Prove that A is connected.
- (2) Generalize to metric spaces.

- (1) Define $f : A \times B \rightarrow A$ by $f((a, b)) = a$. We claim that f is continuous. Let $U \subset A$ be open, then we must show $f^{-1}(U) = \{(a, b) \mid a \in U\}$ is also open. Let $(a, b) \in f^{-1}(U)$, and let $D(a, \varepsilon) \subset U$ (we know this ball exists for some $\varepsilon > 0$ since U is open). We now claim that $D((a, b), \varepsilon) \subset f^{-1}(U)$.

If $(a', b') \in D((a, b), \varepsilon)$, then since $A \times B$ is equipped with the product metric, $|a - a'|^2 \leq |a - a'|^2 + |b - b'|^2 < \varepsilon^2$. This implies $a' \in D(a, \varepsilon) \subset U$, so $(a', b') \in f^{-1}(U)$. Thus $f^{-1}(U)$ is open and, subsequently, f is continuous. Since the image of a connected set under a continuous map is itself connected, we have that A is connected.

- (2) To generalize this to metric spaces, note that the proof only uses a property of \mathbb{R}^n when defining the ε -balls around a and (a, b) . Let $A \times B$ be equipped with the product metric, d_A be the metric for A , and d_B be the metric for B . Then instead of

$$|a - a'|^2 \leq |a - a'|^2 + |b - b'|^2 < \varepsilon^2,$$

we write

$$d_A(a, a') \leq d_A(a, a') + d_B(b, b') < \varepsilon.$$

The rest of the proof is valid without any changes, so the result holds for general metric spaces.

Page 184, 4.2.5. Let $A, B \subset \mathbb{R}$, and suppose $A \times B \subset \mathbb{R}^2$ is open. Must A be open? _____

Yes. Let $a \in A$ and $b \in B$, then there is an $\varepsilon > 0$ such that $D((a, b), \varepsilon) \subset A \times B$. Consider $D(a, \varepsilon)$. For $a' \in D(a, \varepsilon)$, we have $|a' - a| + |b - b| = |a' - a| < \varepsilon$, which implies $(a', b) \in D((a, b), \varepsilon) \subset A \times B$. Thus $a' \in A$, so $D(a, \varepsilon) \subset A$, so A is open.

Page 191, 4.4.3. Let $f : K \subset \mathbb{R}^n \rightarrow \mathbb{R}$ be continuous on a compact set K and let $M = \{x \in K \mid f(x) \text{ is the maximum of } f \text{ on } K\}$. Show that M is a compact set. _____

Since K is compact and f is continuous, K contains all x' such that $f(x') = \sup_x f(x)$. Thus $M = f^{-1}(\{\sup f(x)\})$. The set $\{\sup f(x)\}$ is a single point in \mathbb{R} , so it is closed. Since f is continuous, M must then be closed too. Furthermore, $M \subset K$ and K is bounded (since it is compact), so M must also be bounded. Then by the Heine-Borel theorem, M is compact in \mathbb{R}^n .

Page 193, 4.5.1. What happens when you apply the method used in Example 4.5.4 to quadratic polynomials? To quintic polynomials? _____

Let $f(x) = x^2 + 10$ be a quadratic polynomial. It is clear that $f(x)$ is always positive, so it need not have a real root. Thus the method from the example fails.

Let $f(x) = a_1x^5 + a_2x^4 + a_3x^3 + a_4x^2 + a_5x + a_6$ for $a_1 > 0$. Then we can rewrite this as

$$f(x) = a_1x^5 \left(1 + \frac{a_2}{a_1x} + \frac{a_3}{a_1x^2} + \frac{a_4}{a_1x^3} + \frac{a_5}{a_1x^4} + \frac{a_6}{a_1x^5} \right).$$

The sum in parentheses tends to 1 as $x \rightarrow \infty$, so $f(x) > 0$ for some large and positive x . Similarly, $f(x) < 0$ for some large and negative x . Since \mathbb{R} is connected and f is continuous, by the intermediate value theorem we know that f has a real root.

In general, the largest exponent in a polynomial being odd guarantees that we can find points satisfying $f(x) < 0$ and $f(x) > 0$, in which case the method from the example is valid. As seen with the quadratic example, when the largest exponent is even, this is not always possible.

Page 193, 4.5.2. Let $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be continuous. Let $\Gamma = \{(x, f(x)) \mid x \in \mathbb{R}^n\}$ be the graph of f in $\mathbb{R}^n \times \mathbb{R}^m$. Prove that Γ is closed and connected. Generalize your result to metric spaces. _____

Γ is the image of the map $g : \mathbb{R}^n \rightarrow \mathbb{R}^n \times \mathbb{R}^m$ defined by $g(x) = (x, f(x))$. We claim that g is continuous. Fix $x \in \mathbb{R}^n$ and $\varepsilon > 0$, then we wish to show that the preimage of $D((x, f(x)), \varepsilon)$ under g is open. Since f is continuous, for any $\varepsilon > 0$ we can find δ such that $|x - y| < \delta$ implies $|f(x) - f(y)| < \varepsilon/2$. Let $\delta' = \min\{\varepsilon/2, \delta\}$, then for any $y \in D(x, \delta')$ we have

$$|x - y| + |f(x) - f(y)| < \varepsilon/2 + \varepsilon/2 = \varepsilon.$$

Since $\mathbb{R}^n \times \mathbb{R}^m$ is equipped with the product metric, this shows that $D(x, \delta')$ is a subset of $g^{-1}(D((x, f(x)), \varepsilon))$. Thus g is continuous. Since g is continuous and \mathbb{R}^n is connected, Γ must also be connected.

Now we show that Γ is closed. Let $\{(x_n, f(x_n))\}_{n=1}^\infty$ converge, i.e. $(x_n, f(x_n)) \rightarrow (x, y)$ for some $x \in \mathbb{R}^n$, $y \in \mathbb{R}^m$. The map f is given to be continuous, so $\lim_{n \rightarrow \infty} f(x_n) = f(x)$. Since by assumption, $\lim_{n \rightarrow \infty} f(x_n) = y$, this implies $f(x) = y$. Thus our sequence converges to $(x, f(x))$, which clearly lies in Γ . This shows that Γ is closed.

The proof of connectedness relied on \mathbb{R}^n being connected. If we replace \mathbb{R}^n with any connected metric space, the result still holds. The proof of closedness did not rely on the structure of \mathbb{R}^n at all, so this result holds in any metric space. This gives us a more general proposition:

Let X and Y be metric spaces, and let $f : X \rightarrow Y$ be continuous. Then $\Gamma = \{(x, f(x)) \mid x \in X\}$ is closed. If X is connected, then Γ is connected.

Page 193, 4.5.3. Let $f : [0, 1] \rightarrow [0, 1]$ be continuous. Prove that f has a fixed point. _____

Since $[0, 1]$ is connected and f is continuous, the intermediate value theorem holds. Define another continuous function $g : [0, 1] \rightarrow [-1, 1]$ by $g(x) = f(x) - x$, then note that $g(0) = f(0) \geq 0$ and $g(1) = f(1) - 1 \leq 0$. If $g(0) = 0$, then $f(0) = 0$ and we have a fixed point. Similarly, if $g(1) = 0$, then $f(1) = 1$ and we have a fixed point. So assuming that these trivial cases are false, we have $g(0) < 0 < g(1)$. Then by the intermediate value theorem, there is some $z \in [0, 1]$ such that $g(z) = f(z) - z = 0$, which implies $f(z) = z$, so we have a fixed point.

Page 194, 4.5.4. *Let $f : [a, b] \rightarrow \mathbb{R}$ be continuous. Show that the range of f is a bounded closed interval.* _____

The interval $[a, b]$ is compact, so $f([a, b])$ contains its minimum and maximum, denoted by $f(c)$ and $f(d)$, respectively. Then since $[a, b]$ is also connected, by the intermediate value theorem, the image also contains all points between $f(c)$ and $f(d)$. Thus the image is $f([a, b]) = [f(c), f(d)]$, a closed and bounded interval in \mathbb{R} .

Page 194, 4.5.5. *Prove that there is no continuous map taking $[0, 1]$ onto $(0, 1)$.* _____

Let $f : [0, 1] \rightarrow (0, 1)$ be continuous. Then since $[0, 1]$ is compact, so is $f([0, 1]) \subset (0, 1)$, i.e. $f([0, 1]) = [a, b]$ for some $a > 0$, $b < 1$. But since $a > 0$, this means $a/2$, which is in $(0, 1)$, is not in $f([0, 1])$. Thus f is not onto.