

Initial Date: 07/19/2023

Revised Date:

Section 9-23R

Epinephrine

Pharmacological Category: Sympathomimetic agent

Routes: IV/IO/IM, Nebulized

Indications:

1. Anaphylaxis
2. Bradycardia
3. Respiratory distress
4. Hypotension
5. Cardiac arrest

Expected effects:

1. Decreased wheezing
2. Increased BP
3. Increased HR

Notes:

1. This protocol does NOT apply to Epi Auto Injector (see Epi Auto Injector Protocol)
2. Note that epinephrine is not utilized in the pediatric bradycardia protocol

Preparing PUSH DOSE Epinephrine:

1. Prepare (epinephrine 10 mcg/mL)
 - a. Combine 1 mL of 1 mg/10 mL epinephrine in 9mL NS

Preparing EPINEPHRINE INFUSION:

1. Prepare (epinephrine 2 mcg/mL, recommended concentration)
 - a. Add 1 mg of epinephrine 1:1,000 (1 mL of 1 mg/mL) to a 500 mL bag of normal saline.
 - b. Final concentration is 2 mcg/mL.
 - c. Where a different concentration is prepared, rates must be recalculated for the concentration in use.
 - d. Apply a completed medication-added label showing drug, total dose, final concentration, time, and preparer.
 - e. Epinephrine infusion must be delivered via an IV infusion pump.

Dosing: SHOCK

Indication: Hypotension unresponsive to fluid bolus administration

Adults administer:

1. PUSH DOSE epinephrine 10-20 mcg (1-2 mL of 10 mcg/mL) IV/IO. Repeat every 3-5 minutes. Titrate to SBP > 90 mm/Hg.

Pediatrics administer:

1. PUSH DOSE epinephrine utilizing MI MEDIC cards
2. If MI MEDIC cards are not available administer: PUSH DOSE epinephrine 1 mcg/kg (0.1 mL of epinephrine 10 mcg/mL) IV/IO. Maximum single dose 10 mcg (1 mL). Repeat every 3-5 minutes.

Dosing: ANAPHYLAXIS/ALLERGIC REACTION

MCA Name: Berrien County

MCA Board Approval Date: 8/14/26

MCA Implementation Date: PENDING

MDHHS Approval: PENDING

MDHHS Reviewed 2023

Initial Date: 07/19/2023

Revised Date:

Section 9-23R

Indication: Anaphylaxis/Severe Allergic Reaction

Adults administer:

1. Epinephrine (1mg/mL) 0.3 mg (0.3 mL) IM. May repeat one time after 3-5 minutes if patient remains hypotensive. Maximum of 2 doses total of epinephrine (including epi pen).

Pediatrics administer EPI IM:

1. EPI IM according to MI MEDIC cards
2. If MI MEDIC cards are not available administer: for child weighing < 30 kg, epinephrine (1mg/mL) 0.15 mg (0.15 mL) IM; for child weighing > 30 kg, epinephrine (1mg/mL) 0.3 mg (0.3 mL) IM. May repeat one time after 3-5 minutes if patient remains hypotensive. Maximum of two IM doses total (including epi pen).

Dosing: ADULT RESPIRATORY DISTRESS

Indication: Impending respiratory failure and unable to tolerate nebulizer therapy

Adults administer EPI IM:

1. Epinephrine (1mg/mL) 0.3 mg (0.3 mL) IM

Dosing: CRASHING ADULT/IMPENDING ARREST

Indication: Patient in whom cardiac or respiratory arrest appears imminent and hypotension is unresponsive to fluid bolus administration

Adults administer:

1. PUSH DOSE epinephrine 10-20 mcg (1-2 mL of 10 mcg/mL) IV/IO. Repeat every 3-5 minutes. Titrate to SBP > 90 mm/Hg.

Dosing: PEDIATRIC RESPIRATORY DISTRESS, FAILURE OR ARREST

Indication: Pediatric patient presents with stridor at rest without suspected airway obstruction.

Pediatrics administer EPI IM:

1. EPI IM according to MI MEDIC cards
2. If MI MEDIC cards are not available administer: child weighing < 30 kg, epinephrine (1 mg/mL) 0.15 mg (0.15 mL) IM; child weighing > 30 kg, epinephrine (1 mg/mL) 0.3 mg (0.3 mL) IM

Pediatrics administer NEBULIZED EPI (severe respiratory distress):

1. Epinephrine (1 mg/1 mL) 5 mg nebulized

Dosing: ADULT CARDIAC ARREST

Indication: Cardiac arrest

Adults administer:

1. Epinephrine (1 mg/10 mL) 1 mg IV/IO every 3 to 5 minutes

Dosing: PEDIATRIC CARDIAC ARREST

Indication: Cardiac arrest

Pediatrics administer:

1. Epinephrine according to MI MEDIC cards.
2. If MI MEDIC cards are not available administer epinephrine (1 mg/10 ml), 0.01 mg/kg (0.1 ml/kg). Max dose 1 mg (10 mL). Repeat every 3-5 minutes

Initial Date: 07/19/2023

Revised Date:

Section 9-23R

Dosing: ADULT BRADYCARDIA

Indication: Patients with persistent symptomatic bradycardia

Adults administer:

1. PUSH DOSE epinephrine 10-20 mcg (1-2 mL of 10 mcg/mL) IV/IO. Repeat every 3-5 minutes. Titrate to SBP > 90 mm/Hg.

Dosing: ADULT CHF/CARDIOGENIC SHOCK

Indication: If SBP is below 100 mmHG treat for cardiogenic shock

Adults administer:

1. PUSH DOSE epinephrine 10-20 mcg (1-2 mL of 10 mcg/mL) IV/IO. Repeat every 3-5 minutes. Titrate to SBP > 90 mm/Hg.

Dosing: ADULT ROSC

Indication: Hypotension unresponsive to fluid bolus administration

Adults administer:

1. PUSH DOSE epinephrine 10-20 mcg (1-2 mL of 10 mcg/mL) IV/IO. Repeat every 3-5 minutes. Titrate to SBP > 90 mm/Hg.

Dosing: PEDIATRIC BRADYCARDIA

Indication: If pulse remains < 60, despite oxygenation & ventilation

Pediatrics administer:

1. Epinephrine according to MI MEDIC cards.
2. If MI MEDIC cards are not available administer epinephrine (1 mg/10 mL) 0.01 mg/kg (0.1 mL/kg) IV/IO up to 1 mg (10 mL). Repeat every 3-5 minutes.

Dosing: PEDIATRIC ROSC

Indication: Hypotension unresponsive to fluid bolus administration

Pediatrics administer:

1. PUSH DOSE epinephrine according to MI MEDIC cards, titrating to age appropriate SBP per MI MEDIC cards.
2. If MI MEDIC cards are not available administer PUSH DOSE epinephrine 1 mcg/kg (0.1 mL of epinephrine 10 mcg/mL) IV/IO. Maximum single dose 10 mcg (1 mL). Repeat every 3-5 minutes. Titrate to SBP > 70 mmHG + (2 x age in years) up to 100 mmHg.

Dosing: EPINEPHRINE INFUSION

Indication: Hypotension persisting after appropriate fluid resuscitation where push dose epinephrine is indicated and repeated dosing is required or anticipated. Administer per Epinephrine Infusion-Procedure Protocol (Section 7-32). Requires an IV infusion pump, Critical Care Paramedic privileges under Section 8-17, and MCA approved training in vasopressor infusion administration.

Adults administer:

1. Epinephrine infusion 2 mcg/mL. Begin at 2 mcg/min (60 mL/hr) IV/IO. Titrate in 2 mcg/min increments every 3 to 5 minutes. Usual range 2 to 10 mcg/min (60 to 300 mL/hr). Titrate to SBP > 90 mm/Hg. Contact Medical Control before exceeding 10 mcg/min.

Initial Date: 07/19/2023

Revised Date:

Section 9-23R

Pediatrics administer:

1. MI MEDIC cards do not carry epinephrine infusion dosing. Refer to the pediatric rate table in Epinephrine Infusion-Procedure Protocol (Section 7-32).
2. Begin at 0.1 mcg/kg/min IV/IO, not to exceed the adult starting dose of 2 mcg/min (60 mL/hr). Titrate in 0.1 mcg/kg/min increments every 3 to 5 minutes, each increment not to exceed 2 mcg/min. Titrate to SBP > 70 mmHG + (2 x age in years) up to 100 mmHg. Contact Medical Control before exceeding 1 mcg/kg/min or the adult ceiling of 10 mcg/min (300 mL/hr), whichever is reached first.
3. Do not administer push dose epinephrine and an epinephrine infusion concurrently except as a single bridging dose while the infusion is being prepared.

Used in the Following Protocols

Shock (Section 1 General Treatment)
Anaphylaxis/Allergic Reaction (Section 1 General Treatment)
Respiratory Distress (Section 3 Adult Treatment)
Crashing Adult/Impending Arrest (Section 3 Adult Treatment)
Pediatric Respiratory Distress, Failure or Arrest (Section 4 Obstetrics and Pediatrics)
General Cardiac Arrest (Section 5 Adult Cardiac)
Pediatric Cardiac Arrest – General (Section 6 Pediatric Cardiac)
Bradycardia (Section 5 Adult Cardiac)
Pulmonary Edema/Cardiogenic Shock (Section 5 Adult Cardiac)
Pediatric Bradycardia (Section 6 Pediatric Cardiac)
Return of Spontaneous Circulation (ROSC)-Adult (Section 3 Adult Treatment)
Peds ROSC (Section 4 Obstetrics and Pediatrics)
Epinephrine Infusion (Section 7 Procedures)