>36 kilograms (> 80lbs) Adult >14 Years

CARDIAC RESUSCITATION

BLACK

Normal Vitals: HR: 60-100, RR: 12-20, Systolic BP: 100-140 mmHg, Blood Glucose > 60 mg/dl

Resuscitation Medications - (confirm concentation is as specified)	**Dose**	**Volume**
Epinephrine (1 mg/10 mL prefilled syringe) IV/IO Q 3-5 min **for arrest/bradycardia**	1 mg	**10 mL**
Amiodarone (150 mg/3 mL) IV/IO **for shock resistant V-Fib**	300 mg	**6 mL**
Lidocaine (100 mg/5 mL) IV/IO for shockable V-Fib (or wide-complex tachycardia)	100 mg	**5 mL**
Amiodarone (150 mg/3 mL) IV for stable wide-complex tachy or symptomatic irregular narrow tachy. **Over 10 minutes**	150 mg	**3 mL**
Diltiazam (25 mg/5 mL) IV for symptomatic irregular narrow tachycardia.	20 mg	**4 mL**
Verapamil (10 mg/4 mL) IV for symptomatic irregular narrow tachycardia.	5 mg	**2 mL**
Atropine (1 mg/10 mL) IV/IO **for bradycardia, every 3-5 min to a max of 3 mg**	1 mg	**10 mL**
Adenosine (6 mg/2 mL) IV/IO **1st Dose.** For SVT (HR > 150)	6 mg	**2 mL**
Adenosine (6 mg/2 mL) IV/IO **2nd Dose.** For SVT (HR > 150)	12 mg	**4 mL**
Epinephrine IV/IO (1 mg/10 mL) **Push Dose - Dilute 1 mL with 9 mL Normal Saline** = 10 mcg/1 mL	10 -20 mcg	**1 - 2 mL (Diluted)**
Electrical Therapy	**Initial[1]**	**Repeat[1]**
V-Fib or Pulseless V-Tach: **Defibrillation**	**120-200 J**	**Maximum**
Unstable, wide irregular tachycardia. Heart rate > 150 bpm: **Defibrillation**	**120-200 J**	**Maximum**
Unstable, narrow complex tachycardia. Heart rate > 150 bpm: **Synchronized Cardioversion**	**100 J**	**≥ 100 J[2]**
Narcotic Overdose	**Dose**	**Volume**
Naloxone IV/IM (2 mg/2 mL)	2 mg	**2 mL**
Naloxone IN (2 mg/2 mL) **Divide dose equally between both nostrils**	2 mg	**2 mL IN**
Fluid Bolus		
Normal Saline **1000 mL IV/IO** - **May repeat one time, PRN**		

[1]Based on biphasic, use manufacturer's recommended energy

[2]If no response to first shock, increase energy in a stepwise manner by 20-50 J

>36 kilograms (> 80lbs) Adult >14 Years (Black)

CONDITIONS/MEDICATIONS

BLACK

Normal Vitals: HR: 60-100, RR: 12-20, Systolic BP: 100-140 mmHg, Blood Glucose > 60 mg/dl

Condition	Medication - (confirm concentration is as specified)	Dose	Volume
Wheezing	Albuterol Nebulized (2.5 mg/3 mL)	2.5 mg	**3 mL**
	Ipratropium Bromide Nebulized (0.5 mg/2.5 mL if wheezing)	500 mcg	**2.5 mL**
Anaphylaxis	Diphenhydramine IM/IV/IO (50 mg/mL)	50 mg	**1 mL**
	Prednisone PO (50 mg tablet)	50 mg	**1 tablet PO**
	Methylprednisolone IV/IO/IM (125 mg/2mL)	125 mg	**2 mL**
Anaphylaxis/ Profound Distress	Epinephrine IM (1 mg/mL) ***or*** 1 Epinephrine Autoinjector IM (**Severe symptoms only**)	0.3 mg	**0.3 mL**
Seizure (Sedation)	Midazolam IM (5 mg/mL) **Give first if no IV**	10 mg	**2 mL IM**
	Midazolam IV (5 mg/mL)	5 mg	**1 mL**
Fever/Pain	Ibuprofen PO (100 mg/5 mL)	600 mg	**30 mL PO**
	Acetaminophen PO (160 mg/5 mL) 15 mg/kg, max dose 1 g	800 mg	**25 mL PO**
Hypoglycemia (<60 mg/dL)	D50% (25 g/50 mL) **Give Slow IV**	25 g	**50 mL (D50%)**
	D10% (25 g/250 mL)	25 g	**250 mL**
	Glucagon IM (1 mg/mL)	1 mg	**1 mL IM**
Pain Management	Fentanyl IV (100 mcg/2 mL) **For IN, divide dose equally between nostrils**	50-100 mcg	**1-2 mL**
	Ketamine IV/IO (500mg/5 mL) **Mix dose in 100 mL NS, administer over 10 minutes**	10-20 mg	**0.1-0.2 mL in 100 mL NS**
	Ketamine IN (500 mg/5 mL) Divide dose equally between both nostrils	30-50 mg	**0.3-0.5 mL**
	Ketorolac IV/IM (15 mg/ 1 mL)	15 mg	**1 mL IV/IM**
	Morphine IV/IO/IM (10 mg/mL)	4-5 mg	**0.4-0.5 mL**
Procedural Sedation	Midazolam (5 mg/mL) 1-5 mg titrated **SLOWLY**	1-5 mg	**0.2 - 1 mL**
	Diazepam (10 mg/2mL) 5-10 mg titrated **SLOWLY**	5-10 mg	**1 - 2 mL**
	Fentanyl IV (100 mcg/2 mL) 50-100 mcg titrated **SLOWLY**	50-100 mcg	**1 - 2 mL**
	Ketamine IV (500 mcg/5 mL) 1.5 mg/kg	75-150 mg	**0.75-1.5 mL**
Delirium w/Severe Agitation	Ketamine IM (500 mg/5 mL) 4 mg/kg, MAX dose 500 mg **FOR SEVERE AGITATION ONLY**	200-400 mg	**2-4 mL IM**
Open Fracture	Cefazolin (1000 mg/10 mL) Mix 1 gm of medication per 10 mL NS. **Administer IV SLOWLY.**	2 g	**20 mL**
	Cefazolin (1000 mg/10 mL) Mix as above. **Add specific dose to 100 mL NS.**	2 g	**20 mL in 100 mL NS**
	Ceftriaxone (1000 mg/10 mL) Mix 1 gm of medication per 10 mL NS. **Administer IV SLOWLY.**	2 g	**20 mL**
	Ceftriaxone (1000 mg/10 mL) Mix as above. **Add specific dose to 100 mL NS.**	2 g	**20 mL in 100 mL NS**

MDHHS Michigan Department of Health & Human Services | Bureau of Emergency Preparedness, EMS and Systems of Care

MICHIGAN EMSC State Partnership Program

MI-MEDIC is provided by the Michigan Department of Health and Human Services (MDHHS), Bureau of Emergency Preparedness, EMS and Systems of Care. MI-MEDIC was developed and edited in partnership with Western Michigan University Homer Stryker MD School of Medicine, Division of EMS and Disaster Medicine.

This project is supported by the Health Resources and Services Administration (HRSA) of the US Department of Health and Human Services (HHS) under the EMSC Partnership Grant. This information or content and conclusions are those of the author and should not be construed as the official position or policy of, nor should any endorsements be inferred by HRSA, HHS, or the US Government.

If you have questions or comments about MI-MEDIC, contact: EMS@Michigan.Gov

FST5 Do you need resources related to mental health challenges?
Call 1-833-34-STRONG or go to www.fst5.org

MI-MEDIC by MDHHS, WMed is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License.
Based on a work at www.michigan.gov/ems.