

Initial Date: 07/19/2023

Revised Date:

Section: 9-38R

Ondansetron

Pharmacological Category: Antiemetic

Indications:

1. Nausea and vomiting

Routes: IV/IM; ODT (for patients ≥ 30 kg)

Contraindications:

1. Patients with Phenylketonuria (PKU)

Precautions:

1. Do not administer ODT to patients that are actively vomiting

Expected effects:

1. Diminished nausea

Side effects:

1. Headache
2. Dry mouth
3. Drowsiness

Notes:

1. Orally Disintegrating Tablet (ODT) is an MCA optional medication and may not be available.

Dosing: NAUSEA & VOMITING

Indication: Nausea & vomiting

Adults administer:

1. Ondansetron ODT 4mg if not actively vomiting and ODT is available.
2. Ondansetron 4mg IV/IM if patient is actively vomiting, vomited post ODT administration, or ODT is not available.
3. May administer a second dose of ondansetron 4 mg (IV/IM only). Total dose (including ODT) not to exceed 8 mg.

MCA Name Berrien County

MCA Board Approval 2/9/24

MCA Implementation Date 2/12/24

MDHHS Approval: 7/19/23

MDHHS Reviewed 2023

Page 1 of 2

Initial Date: 07/19/2023

Revised Date:

Section: 9-38R

Pediatrics administer:

1. Ondansetron according to MI MEDIC cards
2. If MI MEDIC cards are not available administer:
 - a. Pediatrics $\geq$ 30 kg that is not actively vomiting and ODT is available administer:
 - i. Ondansetron 4 mg ODT
 - b. Pediatrics < 30 kg, or if the patient is actively vomiting, or if the patient vomited post OD administration, or ODT is not available, administer:
 - i. Ondansetron 0.1 mg/kg IV/IM, maximum dose of 4 mg.
 - c. May repeat ondansetron 0.1 mg/kg IV/IM, maximum dose of 4 mg. Total dose (including ODT) may not exceed 8 mg.

Used in the Following Protocol(s):

Nausea & Vomiting (Section 1 General Treatment)