

Initial Date: 07/19/2023
Revised Date: 08/11/2023

Section: 9-18R

Ceftriazone

Pharmacological Category: Antibiotic, Cephalosporin (Third Generation)

Indications:

1. Open fractures
2. Partial/complete amputations
3. Major soft tissue injuries (e.g., mangled extremity).

Contraindications:

1. Patients $\leq$ 2 months old (any administration of ceftriazone)
2. Infusion <7 years of age (volume for infusion is larger than allowable fluid bolus).
3. Allergies to cefepime (Maxipime) or cefotaxime (Claforan)

Side effects:

1. Rapid administration can result in tachycardia, restlessness, diaphoresis, and palpitations, pain at injection site.

Notes:

Slow IV push dilution of ceftriazone

1. Dilute 2 gm ceftriazone with 20 mL NS:
 - a. Inject two 10 mL flushes into one 2 gm vial of ceftriazone**OR**
 - b. Inject one 10 mL flush into each 1 gm vial of ceftriazone.
2. Resulting concentration is 100 mg/mL

Infusion dilution of ceftriazone

1. Add ceftriazone dosage (slow IV push dilution) to 100 mL bag of NS:
 - a. Adults: add 20 mL (2 gm of slow IV push dilution) to 100 mL bag of NS
 - b. Pediatrics > 7 years of age: volume of diluted ceftriazone added to 100 mL bag of NS will be calculated weight-based dosage.

Dosing: SOFT TISSUE AND ORTHOPEDIC INJURIES

Indication: Partial/complete amputations, major soft tissue injuries (e.g., mangled extremity) and open fractures.

Adults administer:

1. Ceftriazone Slow IVP: 2gm (slow IV push dilution), slow IVP over 3-5 minutes
- OR**
2. Ceftriazone Infusion: 2gm (slow IV push dilution) added to a 100 mL bag of NS.
Infuse over 15-30 minutes.

Pediatrics

1. Pediatrics > 2 months old ceftriazone slow IV push administer:
 - a. Ceftriazone (slow IV push dilution) according to MI MEDIC cards.

Michigan
MEDICATION SECTION
CEFTRIAXONE

Initial Date: 07/19/2023

Revised Date: 08/11/2023

Section: 9-18R

ii. If MI MEDIC cards are not available administer ceftriaxone (slow IV push dilution) 50 mg/kg slow IVP over 3-5 minutes. Maximum dose 2 gm.

OR

2. Pediatrics ≥ 7 years of age ceftriaxone infusion administer:

a. Ceftriaxone infusion according to MI MEDIC cards

i. If MI MEDIC cards are not available administer ceftriaxone (slow IV push dilution) 50 mg/kg added to 100 mL bag of NS. Max dose 2 gm. Infuse over 15-30 minutes.

Used in the Following Protocol(s):

Soft Tissue and Orthopedic Injuries (Section 2 Trauma and Environmental)