



# Infectious Disease & Wilderness Medicine

John C. Bennison<sup>1</sup> · Johnathan Glauser<sup>2</sup>

Accepted: 3 June 2024 / Published online: 24 August 2024

© The Author(s), under exclusive licence to Springer Science+Business Media, LLC, part of Springer Nature 2024

## Abstract

**Purpose of Review** The recognition, treatment, and management of infectious processes is critical for individuals traveling into the wilderness. Particularly in a resource-limited environment, individuals exposed to infectious disease in the wilderness can experience sequelae not commonly observed in the hospital-based urban clinical environment. In the United States, several infectious pathogens are well known to contribute to the development of infection in the wilderness setting. The purpose of this review is to discuss the treatable infectious diseases encountered by individuals venturing into the wilderness of the United States of America. Diligent preparation stands as the cornerstone for safe and rewarding outdoor adventures. By performing critical analysis of current literature, we seek to offer a comprehensive guide on the identification and management of infectious agents that pose a risk to the outdoors people of the United States.

**Recent Findings** Current literature regarding infectious disease in the wilderness consistently highlights the diversity of pathogens encountered in the wild. However, the most common clinical manifestation acquired in the wilderness is diarrhea (Auerbach in *Wilderness Medicine* (7th ed.). Elsevier, 2017 [1], Victora et al. in *Bull. World Health Organ.* 78:(10), 1246, 2000 [2]). Frequently transmitted through fecal–oral route, contaminated water sources are frequently implicated in these cases (Auerbach in *Wilderness Medicine* (5th ed). Elsevier, 2018 [3], [https://www.cdc.gov/healthywater/drinking/travel/backcountry\\_water\\_treatment.html](https://www.cdc.gov/healthywater/drinking/travel/backcountry_water_treatment.html), [www.cdc.gov/eid](http://www.cdc.gov/eid)). Tick-borne illnesses also represent a major avenue by which wilderness explorers become infected (Openshaw et al. in *Am J Trop Med Hyg.* 83(1), 174–182, 2010 [5]). Lyme disease and Rocky Mountain Spotted Fever are two of the most common diseases wilderness travelers face (Rosenberg et al. in *Morbidity and Mortality Weekly Report*, 68(9), 1–11, 2019 [6], <https://www.cdc.gov/rmsf/index.html>). Zoonotic infections are spread when an infectious pathogen is introduced to a human via an animal host. Individuals can become infected by direct contact with the animal host, their bodily fluids, or their environment. Some ticks and mosquitoes are considered “vectors” and directly transmit the infectious agent (Auerbach in *Wilderness Medicine* (7th ed.). Elsevier, 2017 [1], Harms et al. in *J Infect.* 82(1), 98–104, 2021 [7], <https://www.cdc.gov/rmsf/index.html>, Openshaw et al. in *Am J Trop Med Hyg.* 83(1), 174–182, 2010 [5]).

**Summary** Certain infections are prevalent in people travelling to rural United States. This review concentrates on the management of common treatable infections acquired in wilderness areas within the United States.

**Keywords** Emergency medicine · Infectious disease · Wilderness medicine · Water purification · Tick born illness · Water borne illness · Lyme disease · Rocky mountain spotted fever · Giardiasis · E. Coli · Travelers diarrhea · Cryptosporidium · Leptospirosis

## Introduction

### Water Purification

Water purification in the wilderness is crucial for ensuring safe drinking water and preventing spread of waterborne illnesses. It is important to try collecting water from the cleanest source available such as a spring, flowing river, rainwater, or melted snow. Next, allowing sediment to settle will increase the effectiveness of any purification method.

✉ John C. Bennison  
jbennison@metrohealth.org

<sup>1</sup> Department of Emergency Medicine, Metrohealth, 2500 Metro Health Drive, Cleveland, OH 44109, USA

<sup>2</sup> Emergency Medicine Faculty, MetroHealth Medical Center, Case Western Reserve University, Cleveland, OH 44109, USA

If possible, utilizing a combination of the following methods provides the best protection against waterborne illness [4, 9]. Water purification can be summarized to 3 primary methods; boiling, filtration, and disinfection.

Boiling water is the most effective method for killing the various bacteria, viruses, and protozoa responsible for waterborne illness. Water should be brought to a boil for at least 1 min. At altitude greater than 6,500 feet, water should be boiled for a total of 3 min [4, 9].

Filtration involves passing the water through a physical barrier that removes most bacteria and protozoa. However, filtration has limited effectiveness against viruses as they are typically small enough to pass through the filter. Only filters with a chemical disinfectant matrix will be effective against some viruses [4, 9].

Disinfection utilizes chemical agents or ultraviolet light to kill or inactivate microorganisms. Chemical agents such as chlorine, iodine, and chlorine dioxide are commonly used. These agents are mixed in the water source, creating a reaction which kills microorganisms. These chemicals can leave behind an unpleasant taste and require longer wait time for safe drinking. Ultraviolet light works by damaging the DNA of these organisms rendering them innocuous. Ultraviolet light works almost immediately and doesn't leave behind a foul taste; however it can be limited by the clarity of the water [4, 9].

## Lyme Disease

Lyme disease is the most common vector borne zoonotic infection in the United States [6]. Lyme disease is caused by the spirochete bacteria, *Borrelia Burgdorferi*. This bacteria is transmitted to humans by the *Ixodes scapularis* tick, also known as the deer tick or the black legged tick. Because the disease is spread via this host, peak transmission occurs during the warmer months; (May through August) [6].

The disease takes its name from Old Lyme, Connecticut. This is the town from which a group of cases were initially investigated during the 1970's by researchers from Yale. Since that time, Lyme disease has been reported internationally. However, it has been reported that 96% of cases are localized to just 14 states in the northeast United States [6, 11]. Therefore, geographic location and recent travel are crucial to making the diagnosis.

The clinical manifestations of Lyme disease begin 3–30 days following the tick bite. The initial symptoms to be aware of are fevers, chills, myalgia, arthralgia, lymphadenopathy, and the characteristic “bull’s eye” rash. This rash known as Erythema migrans (EM) is a red circular rash that develops around the site of the tick bite. The rash will gradually expand over a few days and then develop central clearing giving the characteristic appearance. EM is present in 70–80% of Lyme disease cases [6, 12]. Delayed

disseminated manifestations occur weeks to months after a tick bite and include dermatologic, neurologic, cardiac, and rheumatologic pathologies [6, 12]. These latter manifestations are unlikely to be encountered by the wilderness explorer within the acute timeframe. That said, it is not uncommon for patients to present with findings such as facial palsy or heart block who do not recall a specific tick bite.

The most effective means of protection against Lyme disease is preventing tick contact in the first place. Physical barriers such as long clothing and insect repellent such as DEET (N,N-diethyl-meta-toluamide) are effective measures [13, 14]. DEET is a chemical originally created by the US military, became commercially available in 1956 and is considered among the most broad spectrum and efficacious insect repellents. The US Centers for Disease Control and Prevention recommend the use of DEET in prevention of Lyme disease. The safety and efficacy of DEET has been reviewed on several occasions by the US environmental protection agency [13]. With proper use and application, the safety profile of DEET is promising with mild skin reactions being most common. The US Food and Drug Administration has deemed DEET safe for use in 2nd and 3rd trimester pregnant women [15]. The American Academy of Pediatrics does not recommend use of DEET in children < 2 years of age. Children > 2 years of age should use products with a maximum DEET concentration of 33% [13].

Prompt treatment of Lyme disease often leads to rapid and complete resolution [6, 7]. If a tick is attached to the skin, removal of the tick is advised as soon as possible. Using a pair of fine tipped tweezers, grasp the tick as close to the surface of skin as possible and gently pull upward. Attempt to remove all parts of the tick from the skin. After tick removal, thoroughly clean the bite area with rubbing alcohol or soap and water.

If EM develops, antibiotic therapy is advised [7]. Dosage may be adjusted depending on patient's age, medical history, pregnancy, and allergies. Doxycycline, amoxicillin, or cefuroxime axetil are recommended by the CDC as first-line agents. Doxycycline dosing is 100 mg orally twice daily for adults; 4.4 mg/kg per day in two divided doses (maximum 100 mg per dose) for children. The duration of therapy is 10 days [6–8].

Prophylactic antibiotic therapy should be considered on a case-by-case basis [7, 10]. One resource advises use of a single dose of doxycycline for nonpregnant adults and children who meet a set criteria; these include attached tick identified as a deer tick, attachment for > 36 h, prophylaxis within 72 h of tick removal, bite occurring within a highly endemic area, and doxycycline not being contraindicated [10]. This prophylactic dose of doxycycline is 200 mg for adults and 4.4 mg/kg up to maximum dose of 200 mg for children given as a single dose. For those who cannot take doxycycline, an alternative antibiotic is not recommended.

## Rocky Mountain Spotted Fever

Rocky Mountain Spotted Fever (RMSF) is caused by the bacterium *Rickettsia rickettsii*, a coccobacillary, obligate, intracellular organism [16]. This bacterium is spread by three tick species: the American dog tick in the eastern United States, the Rocky Mountain wood tick in the central mountain states, and the brown dog tick in the Southwestern United States [17]. Although cases have been reported in each of the lower 48 states, more than 60% of cases have been reported in North Carolina, Oklahoma, Arkansas, Tennessee, and Missouri. Peak transmission occurs between May and August [5].

RMSF is a rapidly progressive disease and can be deadly without early administration of antibiotics. Most people report sudden onset fever and headache approximately 3–12 days following a bite from an infected tick [18]. A characteristic rash will develop in approximately 90% of patients during their illness. The rash is described as small, flat, pink macules on the extremities, which may progress to include the trunk, palms of the hands and soles of the feet [18]. Development of a petechial rash is an ominous finding and is associated with severe disease. Delayed treatment can progress to more severe illness, including neurological deficits, respiratory compromise, necrosis of tissue surrounding the bite, and multiorgan system dysfunction [17, 18].

Once diagnosis of RMSF is suspected, prompt treatment with doxycycline 100 mg twice daily for adults and 2.2 mg/kg/dose twice per day (max 200 mg) for children should be initiated [10]. Doxycycline is most effective at preventing severe complications if started within the first 5 days of illness. The minimum course of treatment is 5–7 days; however, the CDC recommends treating with doxycycline for at least 3 days after the fever subsides and there is evidence of clinical improvement [7, 17].

## Giardiasis

*Giardia* is the most common protozoal intestinal parasite worldwide and is more prevalent in developing areas with poor sanitation [19]. Giardiasis is also known as “backpacker’s diarrhea” because of the common association with hikers exposed to contaminated water [20]. *Giardia* is transmitted via the fecal–oral route and all age groups may be affected [21]. Most cases occur during the summer months, as this coincides with increased outdoor activity and potential exposures, such as camping or the use of communal water sources. Water is a primary source of infection in community outbreaks, usually in small water systems that use untreated or inadequately treated surface water [28].

Clinical symptoms consist of diarrhea, abdominal cramping, and bloating. Stools have a characteristic foul smell and greasy appearance. Limited cases can go on to develop

chronic symptoms including worsening abdominal symptoms, malnutrition, and weight loss [19, 20].

Disease prevention is critical and is promoted by implementation of strict handwashing and avoidance of ingesting contaminated water. Treatment of symptomatic patients includes the use of nitroimidazoles such as tinidazole (2 g once), metronidazole (250 mg three times per day for 5 days), or nitazoxanide (500 mg twice daily for 5 days) [19].

## Escherichia Coli

Overall, the most common cause of traveler’s diarrhea is Enterotoxigenic *Escherichia coli* (ETEC) [2]. *E. coli* are normal inhabitants of the human gastrointestinal tract. However, when they acquire additional genetic material they can become pathogenic [22]. ETEC specifically express fimbriae which allow for intestinal attachment. ETEC also produces secretory toxins which manifest clinically as watery diarrhea, which is the hallmark symptom of this bacterium [23]. ETEC outbreaks are associated with unsafe preparations and storage of food and water. ETEC has a relatively short incubation period of 1–3 days. The illness is self limited, lasting 1–5 days [24].

The treatment of this acute diarrhea focuses on the correction of fluid and electrolyte losses. The majority of these cases can be treated with oral rehydration salts [25]. The World Health Organization and United Nations Children’s Fund recommends a reduced osmolality solution containing 2.6 g sodium chloride, 2.9 g sodium citrate, 1.5 g potassium citrate, and 13.5 g glucose per one liter of clean water [26]. For those individuals exhibiting signs of severe volume loss (lethargy, inability to drink, weak radial pulses) IV fluid replacement with a balanced crystalloid solution (ex. Lactated Ringer’s) is advised [27]. Antimicrobial therapy is not recommended for acute watery diarrhea, as most cases resolve spontaneously [9, 26].

## Cryptosporidium

Cryptosporidiosis, caused by *C. parvum* and *C. hominis*, are common intestinal parasites. Reporting is infrequent because patients may be asymptomatic and symptoms when present typically resolve within two weeks in immunocompetent patients. *Cryptosporidium* infections occur typically in summer and fall, related to drinking water contamination, animal contact, use of recreational water facilities and travel. Ammonia or formalin have been proposed to purify drinking water. Cryptosporidiosis, as with other waterborne infections, has been associated with non-wilderness exposures. It has been reported that 43 million US residents are served by private wells or domestic water systems that are not regulated by the Environmental Protection Agency Safe Drinking Water Act [9]. For our purposes, enteric pathogens from

ingesting untreated water can be introduced via stormwater runoff, animal waste, sewage or septic system malfunction, or swimming in lakes and rivers. Infectious *Cryptosporidium* oocysts have tough outer shells which confer extreme chlorine tolerance [9]. *Cryptosporidium* infections have been reported to cause of the order of 211,000 exposures annually in the United States resulting in approximately 750 hospitalizations. As a classical waterborne pathogen, *Cryptosporidium* is predominantly sourced from animals such as cattle, with high prevalence rates in wastewater and higher detection rates and concentrations during warmer seasons [28].

Symptoms ranging from mild watery diarrhea to severe enteritis with possible biliary tract involvement may occur. The latter may be manifest by elevated alkaline phosphatase, or an enlarged gall bladder with a thickened wall, and dilated biliary ducts on ultrasound. Joint pain, weight loss, abdominal pain and fatigue may be presenting complaints [19].

Diagnosis is most reliably made by microscopy/immunofluorescence, enzyme immunoassays or polymerase chain reaction. Routine stool ova and parasite examinations may not detect the oocysts so, if suspected, the laboratory should be notified in order to use the correct staining techniques [21].

Treatment is, for the most part, supportive with hydration. Antidiarrheal agents such as loperamide or even tincture of opium have been advocated. Spontaneous recovery typically occurs within one week. Parasitic cure also occurs within a few weeks to months [19]. If symptoms persist beyond two weeks, Nitazoxanide is approved by the US Food and Drug Administration for treatment. A typical dosage for an adult is 500 mg twice daily for three days [29]. Patients should be warned that the drug may cause nausea, headache, and bright yellow urine. Patients should avoid public pools for two weeks following diarrhea.

Patients with advanced HIV, for example CD4 counts of less than 50 cells/microliter, may need immune restoration with antiretroviral therapy (ART) before symptoms can be eradicated and may require nitazoxanide therapy for many weeks. While nitazoxanide is the only FDA approved therapy for treatment of cryptosporidiosis, other agents have been proposed for treatment, notably paromomycin [29]. ART and management of infections in immunocompromised patients is beyond the scope of this review.

## Leptospirosis

Leptospirosis is a zoonotic waterborne disease caused by spirochetes of the genus *Leptospira*. Adolph Weil described it in 1886 from kidney tissues, and the lethal disseminated disease has been known as Weil's Disease [30]. In Weil's description, patients had icterus, renal failure, conjunctivitis, and an enlarged spleen [31]. It is highly prevalent in the tropics, and particularly in Southeast Asia, and may be the most widely spread zoonotic infection in the world [32]. Flooding

from hurricanes or monsoons increase the risk of infection from contaminated water.

While rats are the major carriers of leptospires, leptospirosis has been well-documented in dogs, cattle, swine, and horses, among others. Rats excrete very high concentrations of organisms. It is endemic in countries with humid or subtropical climate, but now is also seen in temperate regions due to climate change, human migration, and poor sanitation or waste disposal. There are at least nine species of pathogens associated with human leptospirosis, with many more serotypes and hundreds of serovars of strains. While unusual in recreational travelers within the United States, leptospirosis accounts for over 1 million infections and approximately 60,000 deaths annually worldwide [33].

Transmission of *Leptospira* is generally via direct transmission by contaminated water or soil/carrier mammals. Therefore in tropical regions farmers, agricultural workers, rodent catchers, animal caretakers, fishermen, gardeners and health care professionals are at risk. For the wilderness traveler in the US, skin abrasions and cuts or entry through mucous membranes or conjunctivae are common exposure risks from ponds, rivers, lakes, puddles, or water dams contaminated with rat urine. This applies especially to people who participate in surfing, caving, canoeing, or any freshwater sport. The bacterium has been reportedly able to enter through intact skin as well [32].

Leptospirosis shows two phases of infection. There is a mild anicteric phase, which generally is self-limiting without treatment. The incubation period is approximately 1–2 weeks. Approximately 90% recovery from anicteric leptospirosis has been reported [30]. Leptospirosis can present with protean manifestations: weakness, cough, fever, vomiting, chills, shortness of breath, vomiting and headache. There may be conjunctivitis and uveitis, myalgias, and prostration.

One problem with the diagnosis of leptospirosis is that the signs and symptoms of the disease overlap with those of other diseases which occur in warm humid climates. As well, there may be co-infection with dengue, malaria, chikungunya, hepatitis, or typhus. Patients diagnosed with leptospirosis must have these illnesses considered as well, although some are rare for the wilderness traveler within the US [33].

Recovery takes place in the majority of people. However, in some cases, the infection will re-appear after 2–3 days and cause severe infection with involvement of the kidney, liver, lungs, myocarditis, uveitis and meningitis. There may be myocarditis, encephalomyelitis, intracranial bleeding, or transverse myelitis, and pancreatitis [31]. The icteric phase or Weil's disease ensues when the leptospires infect hepatocytes, causing anuria and oliguria. Leptocytes can be found in the urine from 7–30 days of infection [30]. Pulmonary hemorrhage may occur with or without renal failure or jaundice. Anemia, thrombocytopenia, elevated transaminases, rhabdomyolysis CSF pleocytosis may be present.



Diagnosis may be suspected in some one with a history of risk exposure, perhaps with prostration, jaundice, conjunctivitis, oliguria recently arrived from a camping trip to Hawaii. Blood, urine, and cerebrospinal fluid are typical body fluids used for the diagnosis of leptospiral infection. Leptospire can be found in the blood initially (within the first week), and later in the urine.

Leptospira may be seen under dark-field microscopy (DFM) without staining. DFM can demonstrate leptospire in the blood and CSF in the early bacteremic phase [33].

Blood culture for leptospirosis requires samples during the first week of illness. Culture of leptospire can be performed using a special medium, but takes time for growth.

There are several serological techniques available to check for antibodies against *Leptospira*. The gold standard is the microscopic agglutination test (MAT), but there are commercially available ELISA-based test kits to detect IgG and IgM antibodies. MAT is very sensitive in the early stages of infection, with a titer value of > 400 considered a positive test. ELISA is used to detect IgM antibody in the early stage of infection [30]. ELISA tests become positive after the 6th day of disease onset [32].

Polymerase chain reaction (PCR) has been deemed to be a rapid diagnostic tool on admission, and may be positive even with very low levels of leptospiral DNA [32, 34]. One drawback of serologic testing is the inability to differentiate earlier or past infections. It is suggested to coordinate testing with one's laboratory capabilities.

Prevention includes avoiding contacts with water bodies which are likely to contain urine from rodents, cattle, pigs, or other domesticated animals. Chemoprophylaxis with penicillin or doxycycline has been proposed [30].

As noted above, many cases of leptospirosis are subclinical or resolve spontaneously. Cephalosporins such as ceftriaxone, penicillin, doxycycline, or azithromycin have proven effective. Mild outpatient cases can be managed with doxycycline or azithromycin. Pregnant women can be treated with amoxicillin or azithromycin. For mild disease and no evidence of organ involvement, doxycycline 100 mg twice daily for 7 days is suggested [31].

Patients with severe disease (Weil's) will require more intensive care and supportive therapy. Penicillin G 1.5 million units every 6 h or ceftriaxone 1 g twice daily may be used, with azithromycin or doxycycline for patients allergic to penicillin.

No vaccine is currently available [31]. Vaccine development has been difficult due to the high number of strains—literally in the hundreds. As well, the renal carrier state can last for months or even years, and leptospiral cells can survive in distilled water for months, making the disease difficult to eradicate [30].

**Author Contributions** JB and JG wrote and reviewed the entire manuscript.

**Data Availability** No datasets were generated or analysed during the current study.

## Declarations

**Conflict of Interest** The authors declare that they have no conflict of interests.

**Human and Animal Rights and Informed Consent** This article does not contain any studies with human or animal subjects performed by any of the authors.

## References

1. Auerbach PS. Wilderness Medicine. 7th ed. Elsevier; 2017. **Comprehensive resource on wilderness medicine including infectious diseases and treatment protocols in the wilderness.**
2. Victora CG, Bryce J, Fontaine O, Monasch R. Reducing deaths from diarrhoea through oral rehydration therapy. Bull World Health Organ. 2000;78(10):1246.
3. Auerbach PS. Field Guide to Wilderness Medicine. 5th ed. Elsevier; 2018.
4. Centers for Disease Control and Prevention. Backcountry Water Treatment. 2023. Retrieved from [https://www.cdc.gov/healthywater/drinking/travel/backcountry\\_water\\_treatment.html](https://www.cdc.gov/healthywater/drinking/travel/backcountry_water_treatment.html). **Overview on water treatment methods crucial for an outdoor traveler.**
5. Openshaw JJ, Swerdlow DL, Krebs JW. Rocky Mountain spotted fever in the United States, 2000–2007: Interpreting contemporary increases in incidence. Am J Trop Med Hyg. 2010;83(1):174–82. <https://doi.org/10.4269/ajtmh.2010.09-0752>.
6. Rosenberg R, Lindsey NP, Fischer M, Gregory CJ, Hinckley AF, Mead PS, Paz-Bailey G, et al. Lyme disease — United States, 2001–2018. Morb Mortal Wkly Rep. 2019;68(9):1–11.
7. Harms MG, Hofhuis A, Sprong H, Bennema SC, Ferreira JA, Fonville M, Docters van Leeuwen A, Assendelft WJJ, Van Weert HCPM, Van Pelt W, Van den Wijngaard CC. A single dose of doxycycline after an Ixodes ricinus tick bite to prevent Lyme borreliosis: An open-label randomized controlled trial. J Infect. 2021;82(1):98–104.
8. Lantos PM, Rumbaugh J, Bockenstedt LK, Falck-Ytter YT, Agüero-Rosenfeld ME, Auwaerter PG, Zemel LS. Clinical Practice Guidelines by the Infectious Diseases Society of America (IDSA), American Academy of Neurology (AAN), and American College of Rheumatology (ACR): 2020 Guidelines for the Prevention, Diagnosis, and Treatment of Lyme Disease. Arthritis & Rheumatology. 2021;73(1):12.
9. Gerdes ME, Miko S, Kunz JM, et al. Estimating Waterborne Infectious Disease Burden by Exposure Route, United States 2014. Emerging Infectious Diseases 2023;29(7). Retrieved from [www.cdc.gov/eid](http://www.cdc.gov/eid)
10. Nadelman RB, Nowakowski J, Fish D, Falco RC, Freeman K, McKenna D, Welch P, Marcus R, Agüero-Rosenfeld ME, Dennis DT, Wormser GP, on behalf of the Tick Bite Study Group 2001. Prophylaxis with single-dose doxycycline for the prevention of Lyme disease after an Ixodes scapularis tick bite. N Engl J Med. 2001;345(2):79.
11. Centers for Disease Control and Prevention. Lyme disease --- United States, 2000. 2020 Retrieved November 2020, from <https://www.cdc.gov/mmwr/preview/mmwrhtml/mm5102a3.htm>

12. Centers for Disease Control and Prevention. Lyme disease data tables: most recent year. 2020. Retrieved November 2020, from <https://www.cdc.gov/lyme/datasurveillance/tables-recent.html>
13. American Academy of Pediatrics. Insect repellents. 2024. Retrieved 26 May 2024, from <https://www.aap.org/en/patient-care/environmental-health/promoting-healthy-environments-for-children/insect-repellents/#:~:text=When%20used%20on%20children%2C%20insect,the%20US%20Environmental%20Protection%20Agency>.
14. Chen LH, Wilson ME, Schlagenhauf P. Prevention of tickborne diseases. In: CDC Yellow Book 2020: Health Information for International Travel. Oxford University Press: 2020. p. 293–296.
15. Snodgrass HL, Nelson DC, Weeks MH. Dermal penetration and potential for placental transfer of the insect repellent, N, N-diethyl-m-toluamide. *Am Ind Hyg Assoc J*. 1982;43:747–53.
16. Snowden J, Simonsen KA. Rocky Mountain Spotted Fever (*Rickettsia rickettsii*). 2023. Retrieved 26 May 2024, from <https://www.ncbi-nlm-nih-gov.ccmmain.ohionet.org/books/NBK430881/>
17. Centers for Disease Control and Prevention. Rocky Mountain Spotted Fever (RMSF). 2022. Retrieved from <https://www.cdc.gov/rmsf/index.html>
18. Heymann, D. L. (Ed.). Control of Communicable Diseases Manual (20th ed.). American Public Health Association. 2015.
19. Pyzocha N, Cuda A. Common Intestinal Parasites. *Am Fam Physician*. 2023;108(5):487–94.
20. Centers for Disease Control and Prevention. Giardiasis - Resources for Health Professionals. 2020. Retrieved from [https://www.cdc.gov/parasites/giardia/health\\_professionals/index.html](https://www.cdc.gov/parasites/giardia/health_professionals/index.html)
21. Haque R. Human Intestinal Protozoa. *Gastroenterol Clin North Am*. 2007;36(1):135–46. <https://doi.org/10.1016/j.gtc.2006.12.002>.
22. Cassels FJ, Wolf MK. Colonization factors of diarrheagenic *E. coli* and their intestinal receptors. *J Ind Microbiol*. 1995;15(3):214.
23. Sears CL, Kaper JB. Enteric bacterial toxins: mechanisms of action and linkage to intestinal secretion. *Microbiol Rev*. 1996;60(1):167.
24. Centers for Disease Control and Prevention. Preparing for Travel: Traveler's Diarrhea. In: CDC Yellow Book 2024: Health Information for International Travel. 2024. Retrieved from <https://wwwnc-cdc.gov.ccmmain.ohionet.org/travel/yellowbook/2024/preparing/travelers-diarrhea>. **Reliable resource for healthcare professionals and travelers, providing practical guidance on Traveler's Diarrhea.**
25. Alam NH, Majumder RN, Fuchs GJ. Efficacy and safety of oral rehydration solution with reduced osmolality in adults with cholera: a randomised double-blind clinical trial. *CHOICE study group The Lancet*. 1999;354:296.
26. World Health Organization. The treatment of diarrhoea, a manual for physicians and other senior health workers (4th revision). Geneva, Switzerland: World Health Organization. 2005. **Guidelines on oral rehydration and fluid management.**
27. Hahn S, Kim S, Garner P. Reduced osmolality oral rehydration solution for treating dehydration caused by acute diarrhoea in children. *Cochrane Database of Systematic Reviews*. 2002;CD002847.
28. Su Y, Huang F, Liang B, et al. Occurrence, transmission and risks assessment of pathogens in aquatic environments accessible to humans. *J Environ Manage*. 2024;354:1–13.
29. Sparks H, Nair G, Castellanos-Gonzalez A, White AC Jr. Treatment of Cryptosporidium: What we know, Gaps, and the Way Forward. *Current Tropical Medicine Reports*. 2015;2:181–7.
30. Karpagam KB, Ganesh B. Leptospirosis: a neglected tropical zoonotic infection of public health importance- an updated review. *Eur J Clin Microbiol Infect Dis*. 2020;39:835–46.
31. Rajapakse S. Leptospirosis: Clinical aspects. *Clin Med*. 2022;22(1):14–7.
32. Verma V, Goyal M, Kala D, et al. Recent advances in the diagnosis of leptospirosis. *Frontiers in Bioscience, Landmark* 2020;1655–1681.
33. Md-Lasim A, Shafwati F, Abdul-Halim M, et al. Leptospirosis and coinfection: Should we be concerned? *Int J Environ Res Public Health*. 2021;18:9411. <https://doi.org/10.3390/ijerph18189411>.
34. Woods K, Nic-Fhogartaigh C, Arnold C, et al. A comparison of two molecular methods for diagnosing leptospirosis from three different sample types in patients presenting with fever in Laos. *Clin Microbiol Infect*. 2021;24:1017. <https://doi.org/10.1016/j.cmi.2021.10.017>.

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.

## Terms and Conditions

Springer Nature journal content, brought to you courtesy of Springer Nature Customer Service Center GmbH (“Springer Nature”).

Springer Nature supports a reasonable amount of sharing of research papers by authors, subscribers and authorised users (“Users”), for small-scale personal, non-commercial use provided that all copyright, trade and service marks and other proprietary notices are maintained. By accessing, sharing, receiving or otherwise using the Springer Nature journal content you agree to these terms of use (“Terms”). For these purposes, Springer Nature considers academic use (by researchers and students) to be non-commercial.

These Terms are supplementary and will apply in addition to any applicable website terms and conditions, a relevant site licence or a personal subscription. These Terms will prevail over any conflict or ambiguity with regards to the relevant terms, a site licence or a personal subscription (to the extent of the conflict or ambiguity only). For Creative Commons-licensed articles, the terms of the Creative Commons license used will apply.

We collect and use personal data to provide access to the Springer Nature journal content. We may also use these personal data internally within ResearchGate and Springer Nature and as agreed share it, in an anonymised way, for purposes of tracking, analysis and reporting. We will not otherwise disclose your personal data outside the ResearchGate or the Springer Nature group of companies unless we have your permission as detailed in the Privacy Policy.

While Users may use the Springer Nature journal content for small scale, personal non-commercial use, it is important to note that Users may not:

1. use such content for the purpose of providing other users with access on a regular or large scale basis or as a means to circumvent access control;
2. use such content where to do so would be considered a criminal or statutory offence in any jurisdiction, or gives rise to civil liability, or is otherwise unlawful;
3. falsely or misleadingly imply or suggest endorsement, approval, sponsorship, or association unless explicitly agreed to by Springer Nature in writing;
4. use bots or other automated methods to access the content or redirect messages
5. override any security feature or exclusionary protocol; or
6. share the content in order to create substitute for Springer Nature products or services or a systematic database of Springer Nature journal content.

In line with the restriction against commercial use, Springer Nature does not permit the creation of a product or service that creates revenue, royalties, rent or income from our content or its inclusion as part of a paid for service or for other commercial gain. Springer Nature journal content cannot be used for inter-library loans and librarians may not upload Springer Nature journal content on a large scale into their, or any other, institutional repository.

These terms of use are reviewed regularly and may be amended at any time. Springer Nature is not obligated to publish any information or content on this website and may remove it or features or functionality at our sole discretion, at any time with or without notice. Springer Nature may revoke this licence to you at any time and remove access to any copies of the Springer Nature journal content which have been saved.

To the fullest extent permitted by law, Springer Nature makes no warranties, representations or guarantees to Users, either express or implied with respect to the Springer nature journal content and all parties disclaim and waive any implied warranties or warranties imposed by law, including merchantability or fitness for any particular purpose.

Please note that these rights do not automatically extend to content, data or other material published by Springer Nature that may be licensed from third parties.

If you would like to use or distribute our Springer Nature journal content to a wider audience or on a regular basis or in any other manner not expressly permitted by these Terms, please contact Springer Nature at

[onlineservice@springernature.com](mailto:onlineservice@springernature.com)