

Concentrations of Selected Elements in Liver Tissue of Grey Wolves (*Canis lupus*) from Serbia

Srdan Subotić¹ · Željka Višnjić-Jeftić² · Aleksandra Penezić¹ · Duško Ćirović¹

Received: 9 August 2017 / Accepted: 2 November 2017 / Published online: 8 November 2017
© Springer Science+Business Media, LLC, part of Springer Nature 2017

Abstract The grey wolf (*Canis lupus*) is a large carnivore species and a top predator in the ecosystems that it inhabits. Considering its role in food webs, wolves may be exposed to high concentrations of potentially harmful elements. Therefore liver samples from 28 legally hunted wolves were analyzed for concentrations of 16 elements using inductively coupled plasma optical emission spectrometry. The Mann–Whitney *U* test showed a significant difference between the genders only for Li, and there were no differences between individuals caught in different years. The majority of statistically significant correlations between element levels were positive, except for three cases. Compliance with several criteria for suitable bioindicator organisms imply that wolves may serve for monitoring environmental contamination.

Keywords Ecotoxicology · Wolf · ICP-OES · Element concentration

Exposure to non-essential elements, such as arsenic (As) or cadmium (Cd), can cause different negative effects on health (mutagenic, carcinogenic or teratogenic effects, immune suppression, growth inhibition, impaired reproduction) (Gall et al. 2015). Moreover, these elements are toxic even in trace amounts (Yilmaz et al. 2010). Furthermore, some essential elements needed in biological systems, such as iron (Fe) or

zinc (Zn), can show toxicity at high concentrations (Yilmaz et al. 2010).

The concentration of elements in soil depends primarily on the geological composition of parent material (Shan et al. 2013), although some elements are transported atmospherically and settled by wet or dry deposition (Rodriguez et al. 2012). Many plants are able to bioaccumulate elements and some species have the ability to hyperaccumulate various elements (Mann et al. 2011), and thus can become one of the first links of element transfer in the terrestrial food web (Gall et al. 2015).

The grey wolf (*Canis lupus* Linnaeus, 1758) is the most widespread large carnivore in Europe (Randi 2011), and represents one of the top predators in terrestrial ecosystems (Sergio et al. 2008). Conservation success, protective legislation, and supportive public opinion in the last few decades have led to the recovery of European populations (Chapron et al. 2014), so several areas have been recolonized (Stronen et al. 2013). In European Union countries, wolves are under strict legal protection (Kaczensky et al. 2012). In Serbia, this carnivore has a relatively continuous range, which encompasses most forested areas of western, southwestern, southern, southeastern and eastern parts of the country (Paunović et al. 2008). With the exception of a small isolated population in southern Banat, there is no protection of this species in Serbia, due to its stable population trend and a population size estimated at 700–800 individuals (Paunović et al. 2008; Djan et al. 2014).

The aims of this study are: (a) to evaluate the possibility of using the grey wolf, as the top predator, for monitoring of environmental contamination; (b) to determine the levels of toxic elements and trace metals in wolf liver; (c) to explore differences between the genders and correlations with regard to element concentrations; (d) to compare our findings with the results of other similar studies.

✉ Srdan Subotić
ssubotic@bio.bg.ac.rs

¹ Faculty of Biology, University of Belgrade, Studentski trg 16, 11000 Belgrade, Serbia

² Institute for Multidisciplinary Research, University of Belgrade, Kneza Višeslava 1, 11000 Belgrade, Serbia

Materials and Methods

Sjenica is located on the Pešter plateau, in the mountain area of south-western Serbia. The study area comprises 1056 km² of average altitude 1000 masl. Because of the surrounding mountains and poorly developed infrastructure, this region is scarcely populated. Vegetation is dominated by pastures and meadows, and the old conifer forests cover only about 20% of the investigated area. The primary commercial activity is still the traditional way of animal husbandry (Institute for nature conservation of Serbia 2013). Notable is the presence of the “Štavalj” coal mine inside the sampling area. The mine has a small production (70,000 t annually) of hard brown coal (Turković et al. 2012). With the exception of this mine, industrial production is not developed (Institute for nature conservation of Serbia 2013).

In co-operation with the local hunting association from Sjenica, the carcasses of legally hunted wolves were obtained during the winter season. Out of 28 animals, 16 were females (average length 112.95 ± 3.62 cm; average weight 28.30 ± 4.69 kg) and 12 were males (average length 119.75 ± 7.09 cm; average weight 34.38 ± 5.21 kg). Nine individuals were collected in 2011, 11 in 2012, 7 in 2013, and 1 in 2015. During post mortem analysis of these specimens, liver samples were taken, rinsed with distilled water and stored at -18°C prior to analysis by inductively coupled plasma optical emission spectrometry (ICP-OES). The liver was chosen as the target organ as it is the major detoxification site for contaminants, and it can accumulate inorganic substances (Becker 2000). Another reason for using liver tissue is to facilitate comparison of our findings with those from other toxicological studies performed on wolves.

Sample portions between 0.3 and 0.5 g (wet weight) were processed in a microwave digester (ETHOS EASY ADVANCED MICROWAVE DIGESTION SYSTEM 230V/50 Hz, Mileston, Italy), using 9 mL of 65% HNO₃ (Suprapur®, Merck) and 1 mL of 30% H₂O₂ (Suprapur®, Merck) at a Fresh liver temperature program. Potential presence of the analyzed elements in the reagents was resolved by digestion of blank samples. After cooling to room temperature, the digested samples were diluted with distilled water to a total volume of 25 mL and the concentrations of 16 elements (Al, B, Ba, Cd, Co, Cr, Cu, Fe, Hg, Li, Mn, Mo, Ni, Pb, Sr, and Zn) determined by ICP-OES (Spectro Genesis EOP II, Spectro Analytical Instruments DmbH, Kleve, Germany). The following wavelength lines were used: Al 394.401 nm, B 249.773 nm, Ba 233.527 nm, Cd 228.802 nm, Co 228.616 nm, Cr 205.552 nm, Cu 324.754 nm, Fe 259.941 nm, Hg 184.950 nm, Li 460.289 nm, Mn 259.373 nm, Mo 202.095 nm, Ni 231.604 nm, Pb 220.353 nm, Sr 460.733 nm and Zn 206.1919 nm. The quality of the analytical process was controlled through analysis of BCR-185R reference materials of bovine liver, for the

following elements: Cd, Cu, Mn, Pb, and Zn. IAEA-336 reference materials of lichen was used for the quality of the analytical process for the remaining elements, except for Li, Mo and Ni for which there were no reference materials. The concentrations found were within 90%–115% of the certified values for all measured elements. Concentrations of all elements were expressed as mg kg⁻¹ wet weight (ww).

The Shapiro–Wilk test was used to infer if the data obtained showed normal distribution. Due to non-normal distribution, we applied the Mann–Whitney *U* test to determine the significance of differences in the levels of elements in liver tissue between the genders. The Kruskal–Wallis test for *k* independent samples was used to examine differences in levels of the elements in individuals between sampling years (2011, 2012, and 2013). Finally, correlation analysis (bivariate correlations with Spearman's correlation coefficients) was used to check for significant relationships between the length and weight of wolves and element content.

Results and Discussion

The levels of Al and B were below the detection limit, whereas the other elements were above the detection threshold in all samples. Mean values and ranges for element levels in females and males, as well as in all individuals, are presented in Table 1. The highest concentrations were detected for Fe and Zn.

Most published data on the subject of toxic element concentrations in large mammals concern animals important in the human diet, such as cattle, deer, sheep, wild boar (Gall et al. 2015). Several papers have focused on these problems in tissues from carnivorous species but only a few for the wolf (Gamberg and Braune 1999; Gnamuš et al. 2000; Shore et al. 2001; Flaten et al. 2008; Hoffmann et al. 2010; Bilandžić et al. 2012; Hernández-Moreno et al. 2013; Vihnanek Lazarus et al. 2013; Lazarus et al. 2017).

We found lower level of Cu and higher levels of Hg and Pb than reported for samples taken in neighbouring Croatia (Bilandžić et al. 2012), while Cd levels were similar. These authors also observed that liver content of Cd was higher than for Pb, which is not the case in our study. Due to the uneven sample size, comparison with the results of their paper should be kept under scrutiny. However, also in Croatia but with a larger sample, Vihnanek Lazarus et al. (2013) reported a higher content of Pb than Cd in liver, which is corroborated in our study. In a comprehensive ecotoxicology study on carnivores in Croatia, Lazarus et al. (2017) found higher contents of Co, Cu, and Zn, but lower contents of Hg, Mo, and Pb in wolf liver, than we obtained in our study. Levels of Cd, Fe, and Mn were similar. Our results for Pb were lower and for Hg (minimum and maximum values)

Table 1 Concentrations of Ba, Cd, Co, Cr, Cu, Fe, Hg, Li, Mn, Mo, Ni, Pb, Sr, and Zn in wolf liver samples (mg kg⁻¹ ww) (mean value ± standard deviation; ranges are presented in brackets)

Element	Detection limits (mg L ⁻¹)	Females	Males	Total
Ba	0.000531	0.26 ± 0.62 (0.01–2.40)	0.32 ± 0.63 (0.01–2.14)	0.29 ± 0.61 (0.01–2.40)
Cd	0.000132	0.11 ± 0.08 (0.004–0.36)	0.07 ± 0.04 (0.003–0.11)	0.09 ± 0.07 (0.003–0.36)
Co	0.00024	0.02 ± 0.02 (0.01–0.06)	0.04 ± 0.04 (0.005–0.10)	0.03 ± 0.03 (0.005–0.10)
Cr	0.000366	0.15 ± 0.10 (0.08–0.49)	0.14 ± 0.03 (0.09–0.20)	0.15 ± 0.08 (0.08–0.49)
Cu	0.000588	5.26 ± 2.48 (2.52–11.37)	5.82 ± 4.90 (2.50–19.41)	5.50 ± 3.65 (2.50–19.41)
Fe	0.000562	143.77 ± 60.65 (49.92–292.31)	168.19 ± 62.25 (69.62–276.84)	154.24 ± 62.69 (49.92–292.31)
Hg	0.00553	0.16 ± 0.08 (0.08–0.38)	0.13 ± 0.07 (0.02–0.28)	0.15 ± 0.07 (0.02–0.38)
Li	0.042	1.00 ± 0.33* (0.50–1.45)	0.67 ± 0.26* (0.42–1.09)	0.86 ± 0.34 (0.42–1.45)
Mn	0.000403	3.94 ± 0.98 (2.82–6.24)	3.35 ± 0.64 (2.32–4.54)	3.69 ± 0.88 (2.32–6.24)
Mo	0.000784	0.85 ± 0.36 (0.41–1.52)	0.71 ± 0.22 (0.38–1.08)	0.79 ± 0.31 (0.38–1.52)
Ni	0.00114	0.04 ± 0.07 (0.004–0.31)	0.02 ± 0.01 (0.003–0.04)	0.03 ± 0.06 (0.003–0.31)
Pb	0.00343	0.49 ± 0.50 (0.16–2.18)	0.88 ± 0.99 (0.08–3.47)	0.66 ± 0.76 (0.08–3.47)
Sr	0.00138	0.50 ± 0.29 (0.19–1.07)	0.47 ± 0.21 (0.23–0.89)	0.49 ± 0.26 (0.18–1.07)
Zn	0.000391	24.01 ± 7.77 (13.22–39.28)	24.24 ± 6.52 (18.09–38.53)	24.11 ± 7.13 (13.22–39.28)

The values in the same row are significantly different (Mann–Whitney *U* test, *p* < 0.05)

were higher than Shore et al. (2001) found in wolf livers from Russia.

The study of Hernández-Moreno et al. (2013) showed higher contents of Cd and Pb in liver of the Iberian wolf (*C. lupus signatus*) than we found for the grey wolf, while Zn levels were similar. Higher concentrations of Cd, Co and Zn were reported in arctic wolves from Canada by Gamberg and Braune (1999). They also found lower concentration of Pb but similar concentrations of Hg when compared with our results. In a survey of wolves from Norway, Flaten et al. (2008) obtained higher levels of Cd, Cu, Mn, and Zn than we observed in our paper. They found lower levels of Ba, Hg, Mo, Pb, and Sr, while Co and Ni levels were similar to ours. For comparisons with these studies, we converted the results expressed, published in dry weight, to wet weight using 3.33 as the conversion factor, as the water content in wolf liver is about 70% (Gamberg and Braune 1999).

The Mann–Whitney test revealed a significant difference between the genders, only for Li concentration (females had a higher level). According to the Kruskal–Wallis test, there were no statistically significant differences between individuals sampled in different years.

Currently, there are few papers concerning differences in element concentration between the genders in wolves. Hernández-Moreno et al. (2013) also observed no significant differences in liver element content between the genders but there was a difference in Cd concentration in kidney tissue. Element analysis in wolf kidneys, from the USA and Canada, showed no disparity between the genders, except for Fe concentration (Hoffmann et al. 2010). Lastly, Lazarus et al. (2017) found differences between the genders for concentrations of Cu, Fe, Mn and Tl in muscle tissue, for As, Ca and Tl concentration in liver tissue, and for Tl in kidney tissue. Other carnivores also exhibited no difference between genders, with regard to element concentration, which is shown by Millán et al. (2008) and Pérez-López et al. (2016), in their studies on red foxes (*Vulpes vulpes*).

The majority of statistically significant correlations between length, weight and element levels in liver were positive. The only statistically significant negative correlations were between length and Ni concentration (*r* = −0.454, *p* < 0.05), between weight and Ni concentration (*r* = −0.389, *p* < 0.05) and between the concentrations of Co and Mo (*r* = −0.449, *p* < 0.05).

A good bioindicator organism should meet several criteria such as: (1) abundance and wide distribution to enable continual sampling and comparison with other research; (2) a long lifespan in order to compare accumulation at various life stages; (3) a suitable target tissue for analysis; (4) easy sampling; (5) an important position in the food chain (Zhou et al. 2008). When considering these criteria, the wolf is a favorable model organism for monitoring element contamination in its home range. It is relatively abundant in Serbia (Djan et al. 2014), it is widespread in Europe (Randi 2011), has a life span of around 13 years (Mech 1988; Holyan et al. 2005), and it is one of the top predators in the food chain (Sergio et al. 2008). Moreover, samples of target tissues are easily obtainable during annual wolf hunts.

This is a first study of element concentration in tissues of the grey wolf and only the second (Ćirović et al. 2015) for a canid species from Serbia. Our results are meaningful due to an adequate sample size from a relatively small area, and the number of detected elements. Also this is the first study that considers correlations between element levels in wolf tissues. As a species, the wolf, could potentially be used for environmental monitoring in Serbia, but more work is needed, especially to determine the levels of toxic and trace elements in the wolf's prey, as well as in other carnivore species.

Acknowledgements We acknowledge the support provided by Projects No. TR31009 and 173045, funded by Ministry of Education, Science and Technological Development of the Republic of Serbia. The authors would like to thank dr Miroslav Nikolić and dr Ljiljana Kostić-Kravljanić for their help during heavy metal analysis, and to anonymous reviewers for providing helpful comments and suggestions that improved the quality of the paper.

References

- Becker PR (2000) Concentration of chlorinated hydrocarbons and heavy metals in Alaska arctic marine mammals. *Mar Pollut Bull* 40:819–829
- Biladžić N, Deždek D, Sedak M, Đokić M, Šimić B, Rudan N, Brstilo M, Lisicin T (2012) Trace elements in tissues of wild carnivores and omnivores in Croatia. *Bull Environ Contam Toxicol* 88:94–99
- Chapron G, Kaczensky P, Linnell JDC, von Arx M et al (2014) Recovery of large carnivores in Europe's modern human-dominated landscapes. *Science* 346:1517–1519
- Ćirović D, Gizajewska A, Jovanović V, Penezić A, Milenković M, Vujošević M, Blagojević J (2015) Concentration of selected trace elements in golden jackal (*Canis aureus* L., 1758) population from Serbia. *Acta Zool Bulg* 67:409–414
- Djan M, Maletić V, Trbojević I, Popović D, Veličković N, Burazerović J, Ćirović D (2014) Genetic diversity and structuring of the grey wolf population from the Central Balkans based on the mitochondrial DNA variation. *Mamm Biol* 79:277–282
- Flaten TP, Andersen LH, Holsen AMH, Ottemo VG, Pedersen HC, Steinnes E, Lierhagen S, Jenssen BM (2008) A nationwide survey of trace elements in lynx, wolverines, wolves, and brown bears in Norway. *Cell Biol Toxicol* 24(Suppl 1):6–7
- Gall JE, Boyd RS, Rajakaruna N (2015) Transfer of heavy metals through terrestrial food webs: a review. *Environ Monit Assess* 187:201. <https://doi.org/10.1007/s10661-015-4436-3>
- Gamberg M, Braune BM (1999) Contaminant residue levels in arctic wolves (*Canis lupus*) from Yukon Territory. *Can Sci Total Environ* 243:329–338
- Gnamuš A, Byrne AR, Horvat M (2000) Mercury in the soil-plant-deer-predator food chain of a temperate forest in Slovenia. *Environ Sci Technol* 34:3337–3345
- Hernández-Moreno D, de la Casa Resino I, Fidalgo LE, Llana L, Soler Rodríguez F, Pérez-López M, López-Beceiro A (2013) Noninvasive heavy metal pollution assessment by means of Iberian wolf (*Canis lupus signatus*) hair from Galicia (NW Spain): a comparison with invasive samples. *Environ Monit Assess* 185:10421–10430
- Hoffmann SR, Blunck SA, Petersen KN, Jones EM et al (2010) Cadmium, copper, iron, and zinc concentration in kidneys of grey wolves, *Canis lupus*, from Alaska, Idaho, Montana (USA) and the Northern Territories (Canada). *Bull Environ Contam Toxicol* 85:481–485
- Holyan J, Boyd DK, Mack CM, Pletscher DH (2005) Longevity and productivity of three wolves, *Canis lupus* in the wild. *Can Field Nat* 119:447–448
- Institute for nature conservation of Serbia (2013) Special nature reserve “Peštersko polje”, conservation proposal. Belgrade, p 127
- Kaczensky P, Chapron G, von Arx M, Huber D, Andrén H, Linnell J (2012) Status, management and distribution of large carnivores—bear, lynx, wolf and wolverine—in Europe, part 1. European Commission, p 72
- Lazarus M, Sekanović A, Orct T, Reljić S, Kusak J, Jurasović J, Huber D (2017) Apex predatory mammals as bioindicator species in environmental monitoring of elements in Dinaric Alps (Croatia). *Environ Sci Pollut Res*. <https://doi.org/10.1007/s11356-017-0008-0>
- Mann RM, Vijver MG, Peijnenburg WJGM (2011) Metals and metalloids in terrestrial systems: bioaccumulation, biomagnification and subsequent adverse effects. In: Sánchez-Bayo F, van den Brink PJ, Mann RM (eds) Ecological impacts of toxic chemicals. Bentham e-Books, pp 43–62
- Mech LD (1988) Longevity in wild wolves. *J Mammal* 69:197–198
- Millán J, Mateo R, Taggart MA, López-Bao JV, Viota M, Monsalve L, Camarero PR, Blázquez E, Jiménez B (2008) Levels of heavy metals and metalloids in critically endangered Iberian lynx and other wild carnivores from Southern Spain. *Sci Total Environ* 399:193–201
- Paunović M, Ćirović D, Milenković M (2008) Status, management and conservation of large carnivores in Serbia. In Conference paper of Coexistence of large carnivores and humans: threat or benefit? pp 111–117
- Pérez-López M, Soler Rodríguez F, Hernández-Moreno D, Rigueira L, Fidalgo LE, López Beceiro A (2016) Bioaccumulation of cadmium, lead and zinc in liver and kidney of red fox (*Vulpes vulpes*) from NW Spain: influence of gender and age. *Toxicol Environ Chem* 98:109–117
- Randi E (2011) Genetics and conservation of wolves *Canis lupus* in Europe. *Mamm Rev* 41: 99–111
- Rodríguez JH, Wannaz ED, Salazar MJ, Pignata ML, Fangmeier A, Franzaring J (2012) Accumulation of polycyclic aromatic hydrocarbons and heavy metals in the tree foliage of *Eucalyptus rostrata*, *Pinus radiata* and *Populus hybridus* in the vicinity of a large aluminium smelter in Argentina. *Atmos Environ* 55:35–42
- Sergio F, Caro T, Brown D, Clucas B, Hunter J, Ketchum J, McHugh K, Hiraldo F (2008) Top predators as conservation tools: ecological rationale, assumptions, and efficacy. *Annu Rev Ecol Evol Syst* 39:1–19

- Shan Y, Tysklind M, Hao F, Ouyang W, Chen S, Lin C (2013) Identification of sources of heavy metals in agricultural soils using multivariate analysis and GIS. *J Soils Sediments* 13:720–729
- Shore RF, Casulli A, Bologov V, Wienburg CL, Afsar A, Toyne P, Dell'Omo G (2001) Organochlorine pesticide, polychlorinated biphenyl and heavy metal concentration in wolves (*Canis lupus* L. 1758) from north-west Russia. *Sci Total Environ* 280:45–54
- Stronen AV, Jędrzejewska B, Pertoldi C, Demontis D, Randi E, Niedziałkowska M, Pilot M, Sidorovich VE, Dykyy I, Kusak J, Tsingarska E, Kojola I, Karamanlidis AA, Ornicans A, Lobkov VA, Dumenko V, Czarnomska SD (2013) North-south differentiation and a region of high diversity in European wolves (*Canis lupus*). *PLoS ONE* 8(10):e76454
- Turković M, Bukumirović D, Radeka G, Mušović S (2012) Potentiality and prospects of underground coal mining in the Sjenica coal basin. *Min Eng* 2:115–124
- Vihnanek Lazarus M, Sekovanić A, Kljaković-Gašpić Z, Orčić T, Jurasović J, Kusak J, Reljić S, Huber Đ (2013) Cadmium and lead in grey wolf liver samples: optimisation of microwave-assisted digestion method. *Arh Hig Rada Toksikol* 64:395–403
- Yilmaz AB, Sangün MK, Yağlıoğlu D, Turan C (2010) Metals (major, essential to non-essential) composition of the different tissues of three demersal fish species from İskenderun Bay, Turkey. *Food Chem* 123:410–415
- Zhou Q, Zhang J, Fu J, Shi J, Jiang G (2008) Biomonitoring: an appealing tool for assessment of metal pollution in the aquatic ecosystem. *Anal Chim Acta* 606:135–150

Terms and Conditions

Springer Nature journal content, brought to you courtesy of Springer Nature Customer Service Center GmbH (“Springer Nature”).

Springer Nature supports a reasonable amount of sharing of research papers by authors, subscribers and authorised users (“Users”), for small-scale personal, non-commercial use provided that all copyright, trade and service marks and other proprietary notices are maintained. By accessing, sharing, receiving or otherwise using the Springer Nature journal content you agree to these terms of use (“Terms”). For these purposes, Springer Nature considers academic use (by researchers and students) to be non-commercial.

These Terms are supplementary and will apply in addition to any applicable website terms and conditions, a relevant site licence or a personal subscription. These Terms will prevail over any conflict or ambiguity with regards to the relevant terms, a site licence or a personal subscription (to the extent of the conflict or ambiguity only). For Creative Commons-licensed articles, the terms of the Creative Commons license used will apply.

We collect and use personal data to provide access to the Springer Nature journal content. We may also use these personal data internally within ResearchGate and Springer Nature and as agreed share it, in an anonymised way, for purposes of tracking, analysis and reporting. We will not otherwise disclose your personal data outside the ResearchGate or the Springer Nature group of companies unless we have your permission as detailed in the Privacy Policy.

While Users may use the Springer Nature journal content for small scale, personal non-commercial use, it is important to note that Users may not:

1. use such content for the purpose of providing other users with access on a regular or large scale basis or as a means to circumvent access control;
2. use such content where to do so would be considered a criminal or statutory offence in any jurisdiction, or gives rise to civil liability, or is otherwise unlawful;
3. falsely or misleadingly imply or suggest endorsement, approval, sponsorship, or association unless explicitly agreed to by Springer Nature in writing;
4. use bots or other automated methods to access the content or redirect messages
5. override any security feature or exclusionary protocol; or
6. share the content in order to create substitute for Springer Nature products or services or a systematic database of Springer Nature journal content.

In line with the restriction against commercial use, Springer Nature does not permit the creation of a product or service that creates revenue, royalties, rent or income from our content or its inclusion as part of a paid for service or for other commercial gain. Springer Nature journal content cannot be used for inter-library loans and librarians may not upload Springer Nature journal content on a large scale into their, or any other, institutional repository.

These terms of use are reviewed regularly and may be amended at any time. Springer Nature is not obligated to publish any information or content on this website and may remove it or features or functionality at our sole discretion, at any time with or without notice. Springer Nature may revoke this licence to you at any time and remove access to any copies of the Springer Nature journal content which have been saved.

To the fullest extent permitted by law, Springer Nature makes no warranties, representations or guarantees to Users, either express or implied with respect to the Springer nature journal content and all parties disclaim and waive any implied warranties or warranties imposed by law, including merchantability or fitness for any particular purpose.

Please note that these rights do not automatically extend to content, data or other material published by Springer Nature that may be licensed from third parties.

If you would like to use or distribute our Springer Nature journal content to a wider audience or on a regular basis or in any other manner not expressly permitted by these Terms, please contact Springer Nature at

onlineservice@springernature.com